VOL. 26. Ser. A. Title-page and Index. [Issued September 1939.]

THE REVIEW OF APPLIED ENTOMOLOGY.

SERIES A: AGRICULTURAL.

ISSUED BY THE IMPERIAL INSTITUTE OF ENTOMOLOGY.

LONDON:
THE IMPERIAL INSTITUTE OF ENTOMOLOGY,
41, QUEEN'S GATE, S.W. 7.

Price 6s. net.

All Rights Reserved.

IMPERIAL INSTITUTE OF ENTOMOLOGY.

Executive Council.

F. J. Du Toit, Chairman, Union of South Africa.

Sir Donald Fergusson, K.C.B., United Kingdom.

Dr. William Allen, Canada.

F. L. McDougall, C.M.G., Australia.

Nevill L. Wright, F.I.C., D.I.C., New Zealand.

J. M. Adams, F.R.C.Sc. (I), Eire.

D. James Davies, C.B.E., Newfoundland.

Shamaldhari Lal, India.

B. F. Wright, Southern Rhodesia.

J. A. Calder, C.M.G., Colonies, Protectorates and Mandated Territories.

Sir David Chadwick, K.C.M.G., C.S.I., C.I.E., Secretary.

Director.

Sir Guy A. K. Marshall, C.M.G., F.R.S.

Assistant Director and Editor. Dr. S. A. NEAVE, O.B.E.

Assistant Director and Superintendent of Farnham House Laboratory.
Dr. W. R. THOMPSON, F.R.S.

Head Office—British Museum (Natural History), Cromwell Road, London, S.W.7.

Publication Office-41, Queen's Gate, London, S.W.7.

INDEX OF AUTHORS.

A reference in heavy type indicates that a paper by the author has been abstracted.

Abbott, W. S., 553. Abraham, G., 250. Abraham, R., 52, 234, 250, 418. Acree, jr., F., 694. Aczél, M., 197. Adams, J. A., 224, 387. Ahlberg, O., 299. Ahmad, T., 451. Akiyama, T., 767. Allan, W., 366. Allen, H. W., 108. Allen, M. W., 121, 141, 496. Allen, N., 393, 645. Allman, S. L., 510. Alsterlund, J. F., 20. Amaral, J. F., 320. Ambros, W., 659, 715. Amono, E., 266. Amos, J. M., 442. Amotz, Y. Ben-, 280. Ananda Rau, S., 148. Anantanarayanan, K. P., 240, 360. Andersen, K. T., 248, 519. Anderson, H. W., 587. Anderson, L. D., 13. Andison, H., 483. Andre, F., 23, 225, 283. Andrés, C. González de, 700. Andrewartha, H. G., 111, 584, 703. Andrewartha, H. V., 318. Annand, P. N., 136. Anthon, E. W., 127. Aoyama, T., 605. App, B. A., **729.** Appel, O., **703.** Arant, F. S., **572, 691.** Araujo, R. L., **145.** Arkhangel'skaya, A. D., 301. Ashworth, J. T., 14, 723. Atwood, C. E., 538. Aufhammer, G., 521. Austin, M. D., 296. Autuori, M., 627. Ayyar, T. V. Ramakrishna, 89, 240, 360.

Back, E. A., 224, 227, 228, 367, 400, 568. Bailey, S. F., 128. Baker, A. D., **60, 623.** Baker, F. E., 441. Baker, W. A., **729**, **730**. Baker, W. L., **560**. Balachowsky, A., **259**, **303**, **415**, 559, 560, 696. Balch, R. E., 330, 487, 539. Balduf, W. V., 173. Ball, E. D., **282.** Ballard, E., 77, 514. Barber, G. W., 144, 304, 745. Barnes, D. F., 118, 573. Barnes, H. F., 32, 33. Barreda, L. de la, 248. Basheer, M., 603. Batra, H. N., 765. Batten, E. T., 609. Baudyš, E., 656. Beall, G., 486. Beard, R. L., 17, 725. Beaumont, A., 527. Bebbington, A. G., 366. Becker, W. B., 576. Beckwith, C. S., 508, 562. Bedford, H. W., 511. Beeson, C. F. C., 247, 764. Beffa, G. Della, **384, 440.** Belanovskii (Bilanovs'kii), I. D., 94, 302. Bell, A. F., 151. Bellio, G., **593.** Bel'skiĭ, B. I., 236. Belyavskaya, A. F., 306. Ben-Amotz, Y., 280. Benlloch Martínez, M., 697. Bennett, C. W., 445. Benson, H. J. Craufurd-, 405, 610, 616 Benton, C., 78. Berezhkov, R. P., 308. Berezina, S., 479. Berezina, V. M., **429, 431.** Berg, V. L., **440.** Berger, E. W., 587.

Berlese, A., 94. Berry, P. A., 745. Besse, R. S., 291. Betrem, J. G., 606, 607. Bev. M. Kamel, 682. Bhalla, Hem Raj, 180. Bhatia, H. L., 181. Bhatta, K. L., 360. Bianchi, F. A., 492, 493. Bigger, J. H., 391. Bigi, F., 592. Bilanovs'kiĭ (see Belanovskiĭ). Bishopp, F. C., 385. Bisset, G. A., **342**. Blanchard, E. E., **582**, **689**. Blanton, F. S., 288, 393. Blattny, C., 658. Bleton, C. A., 341, 501, 641. Bliss, C. I., 165, 633. Blöte, H. C., 268. Blunck, H., 250, 345, 346, 520. Boam, J. J., **616.** Bobb, M. L., **271, 506.** Bobibskaya, S. G., 425. Bodenheimer, F. S., 159, 636, 677. Bodine, J. H., 304. Bodkin, G. E., 316. Boell, E. J., 304. Boer, S. de, **599.** Boerger, A., **290.** Böhmel, W., **50.** Bohorquez, R., 700. Bondar, G., 501, 643. Bondy, F. F., 644. Borkhsenius, N. S., 304. Börner, C., 50, 601. Bosma, B., 465. Bosq, J. M., 696. Bottger, G. T., 729. Bottimer, L. J., 28. Bouet, G., 180. Bouhelier, R., 414. Bourgin, G. Viennot-, 259, 525. Bourne, A. I., **646.** Bovey, P., 99, 163, 299. Bovien, P., 49, 468. Boyce, A. M., 508. Boyce, H. R., 535. Bozzini, G., **598.** Bradley, W. G., **729.** Brahmachari, K., 499. Brammanis, L., 90, 721. Brande, J. van den, 193. Braun, R., 459. Bredo, H. J., 687.

Bréjoux, D., 619. Bremer, H., 39. Brémond, P., 416, 526. Briant, A. K., 175. Bridwell, J. C., 292. Brimblecombe, A. R., 451. Brindley, T. A., 26, 653. Brittin, G., 303. Britton, W. E., 14, 15, 722, 723, 726. Broekhuizen, S., 655. Bromley, S. W., 546, 560. Bronson, T. E., 740. Brooks, H. E., 297. Brown, A. W. A., 534. Brown, H. B., 722. Browne, F. G., 109, 110, 295, 453, 763. Bruel, W. E. Van den, 70, 313. Bruneteau, J., **526.** Bryant, G. E., **372.** Bryson, H. R., 292. Buckell, E. R., 676, 680, 686. Buckner, R. P., 117. Bueno, J. R. de la Torre-, 11. Bulger, J. W., 134. Burgess, A. F., 560. Burks, B. D., 573. Burnham, J. C., **144.** Busbey, R. L., 21, 695. Busvine, J. R., 633. Butovitsch, 611. Buxton, D. R., 679. Bwardwaj, N. K., 87. Bynum, E. K., 137. Bywaters, M. F., **166.**

Caguerre, J. B., 689.
Cahn, R. S., 616.
Caldwell, J. S., 367.
Caldwell, N. E. H., 211, 268.
Calfee, R. K., 368, 704.
Campbell, H. L., 30.
Campbell, L. W., 134.
Campbell, R. E., 569, 744.
Candioli, P., 594.
Cañizo, J. del, 697, 698.
Cann, F. R., 463.
Canto, J. P., 221.
Cantrell, I. J., 81.
Canzanelli, A., 153.
Cardoso, J. G. A., 682.
Caresche, L., 184, 185, 187, 188, 205.
Carl, J., 156.

Carlini, L., 189. Carlson, A. B., 218. Carpenter, E. L., 567, 741. Carrol, J., 496. Carruth, L. A., 564. Carter, W., 143. Carter, W. B., 116. Cartwright, K. St. G., 632. Cartwright, O. L., 643. Caserio, E., 411. Cassab, A., 457. Cassil, C. C., 368, 567, 695. Castro, A. Ruiz, 700, 701. Cavitt, H. S., 274. Chaboussou, 523. Chamberlain, K. F., 215. Chamberlin, F. S., **132, 144.** Chan (Mong Shi), **243.** 734. Chapin, E. A., 26. Chapman, A. J., 274. Chapman, P. J., 287. Chapman, R. N., 311. Chatten, F., 652. Chauvin, R., 440, 624. 603. Chernuishev, P. K., 74. Chevalier, G., 371. Chin (Meng-hsiao), 244. China, W. E., 8, 688. Chiu (Shin-foon), 109. Chopard, L., 629. Christensen, C., 43. Christie, J. R., 123. Christóbal, U. López, 289. Chu, C. V., 498.

Chandler, S. C., 544, 545, 652, 653, Charmoy, D. d'Emmerez de, 501. Cherian, M. C., 55, 408, 531, 602, Chu (Joo-tso), 244. Chugunin, Ya. V., 74. Chûjô, M., 336. Chun, N. M., 674. Ciferri, R., 592. Ciślik, W., 98. Claborn, H. V., 439. Clark, A. F., 58, 752. Clark, E. P., 120, 368, 694. Clark, J. C., 277. Clark, J. R., 386. Cleare, L. D., 221, 369, 491. Clemente, F. Gómez, 699, 702. Clinton, G. P., 749. Coaton, W. G. H., 48. Cochran, W. G., 630, 631. Cohen, M., 710.

(1967)

Colcord, M., 367. Collins, C. W., 551. Comas, J. Nonell, 696. Compere, H., 212. Compton, C. C., 20. Conklin, J. G., 20. Constantino, G., 70, 92, 93, 119. Cook, M. T., 116. Cooke, O. B., 14, 723. Corbett, G. H., **31, 434.** Cory, E. N., **541.** Costa Lima, A. da, 319, 704. Cottier, W., 45. Cotton, R. T., 64, 118, 321, 362, 568. Coutinho Saraiva, A., 315. Couturier, A., 595. Craig, R., 386. Crawford, J. C., 329, 490. Craufurd-Benson, H. J., 405, 610, 616. Creighton, J. T., 22. Cristóbal, U. López, 30, 290. Crosthwait, S. L., 444. Crowley, D. J., **171.** Cupples, H. L., 320, 388. Currie, G. A., 86. Cushman, R. A., 119, 215, 295. Cuthbertson, A., 89.

da Costa Lima, A., 319, 704. Daguerre, J. B., 146. Dahms, R. G., 63, 144. Daniels, L. B., 643. Darling, R. C. Maxwell-, 198. Davidson, J., 584, 695, 703. Davidson, R. H., 369, 746. Davidson, W. M., 104. Davies, W. M., 379, 709. Davis, A. C., 286, 579. Davis, E. W., 283. Davis, J. J., 126. Dawsey, L. H., 227. Dawson, J. C., 23. Daxer, H., 600. Dean, G. A., 118. Dean, R. W., 561. de Andrés, C. González, 700. Dearborn, F. E., 183, 284. de Boer, S., 599. de Charmoy, D. d'Emmerez, 501. Decker, G. C., 23, 283. Decker, P., 217. Defago, G., 91.

Cutright, C. R., 61, 399, 401.

de Fluiter, H. J., 182, 434. de la Barreda, L., 248. de la Torre-Bueno, J. R., 11. de la Vega, G. A. Kreibohm, 362, 704. del Cañizo, J., 697, 698. Delgado de Torres, D., 699. Della Beffa, G., 384, 440. De Long, D. M., 367. de Macedo, W., 483. d'Emmerez de Charmoy, D., 501. de Nardo, A., 469. de Ong, E. R., 282. de Peyerimhoff, P., 410, 688. de Santis, L., 290. Deshpande, V. G., 241, 247, 261. Deshusses, J., 689, 715. Deshusses, L., 689. Deslandes, J., 115. de Sornay, P., 317. de Torres, D. Delgado, 699. De Villiers, F. J., 681. De Vore, Ll., 772. Diakonoff, A., **529**, 529, **599**. Dick, J., **164.** Dietz, H. F., 551. Di Maria, G., 560. Dirks, C. O., 746. Dirsh, V. M., 356. Distant, W. L., 8. Ditman, L. P., 440. Djou, Y. W., 497, 693. Dobreanu, E., 192. Dobromuislov, D., 76. Dobroscky, I. D., 25. Doehlert, C. A., 508. Dohanian, S. M., 43. Domínguez, G., 697. Dominick, C. B., **549.** Donaldson, F. T., 609. Donohoe, H. C., 29, 118, 228, 285, 292, 573. Donovan, W. O'B., **752.** Dorst, H. E., **78, 283, 743.** Dotti, R., 593. Douglas, W. A., 218. Dowden, P. B., 576, 589, 745. Downes, W., 293. Dozier, H. L., 41, 286, 395. Drake, C. J., 23. Drake, N. L., 695. Dreyfus, L., 459. Driggers, B. F., 555, 737, 750. Duarte, A. J., 613. Dubrovenskii, V. F., 430.

Ducker, H. C., 367.

Dudley, jr., J. E., 740. Dumbleton, L. J., 56, 57, 266. Duncan, D. C., 772. Dunlap, A. A., 722. Dunnan, E. W., 277. Du Plessis, C., 676. Dupont, R., 204. Dustan, G. G., 536, 539. Dyk, A., 665. Dzevaltovskaya, N. G., 237.

Ebeling, W., **575.** Eckert, J. E., 129. Eckstein, F., 468, 599. Eckstein, K., 233. Edwards, E. E., 382. Edwards, W. H., 145, 220, 268, 490. Egorova, A. I., 474. Eguchi, M., 309. Eichler, W., **69.** Eĭdel'man, Z. M., 355, 482. Eidmann, H., 4, 154, 248, 346. Elgueta, P. N., 654. Ellenwood, C. W., 402. Elliott, H. G., 754. Ellisor, L. O., 225, 388. El Zoheiry, M. S., **685**. Emchuk, E. M., 159. Emery, W. T., 121. English, L. L., 690. Enser, K., 715. Erlanson, C. O., 749. Esaki, T., **55, 439.** Escherich, K., 496. Essig, E. O., 589. Ewing, H. E., 173. Ewing, K. P., 277. Evans, A. C., 632. Evans, J. W., 8, 208, 582. Everly, R. T., **590.** Eyer, J. R., 137, 281.

Faes, H., 70, 299.
Faggioli, D., 153.
Fahey, J. E., 320.
Fahringer, F., 660.
Fajans, E., 370.
Farrar, M. D., 652, 734.
Farský, O., 665.
Fassig, W. W., 30.
Faure, Jacobus C., 681, 683, 684, 687.
Fawcett, G. L., 581.
Felt, E. P., 215, 546, 692, 772.

Feng, C. L., 498. Fennah, R. G., 402, 751. Fenton, F. A., 136, 138, 139, 246. Ferrière, Ch., 610. Ferris, G. F., 101, 184, 470. Feytaud, J., 5, 314, 412, 594, 597. Fife, L. C., 41, 42. Fischer, A., 496. Fischer, W., 188, 600. Fisher, C. K., 118, 648. Fisher, R. C., 34, 380. Fisher, W. S., 214, 262, 329. Fisk, F. W., 389. Fitzgerald, D. Vesey-, 433, 611. Flachs, K., 304. Flanders, S. E., 287, 744. Fleming, W. E., 441. Fletcher, R. K., 278. Flint, W. P., 124, 391, 544, 652, 653, 734, 748. Floyd, E. H., 388. Fluiter, H. J. de, 182, 434. Foenander, E. C., 109. Foerster, H., 301. Fonseca, J. P., 319, 320, 502. Fontaine, M., 184. Forbes, S. A., 562. Förster, H., 496. Foury, A., 414. Fowler, E. D., 749. Fowler, R., 318. Fox, D. E., 691. Francke-Grosmann, H., 2, 252, 344, 762. Franklin, H. J., 645. Fransen, J. J., 253, 347, 348, 368, Franssen, C. J. H., 261, 262, 295, 765. Frappa, C., 66, 67, 68, 118, 167, 413, 416, 454. Frear, D. E. H., 488, 742. Frezzi, M. J., 581. Frickhinger, H. W., 120. Friend, R. B., 218, 547. Frison, T. H., 573. Froggatt, J. L., **435**, **495**. Frost, S. W., **26**, **135**. Fukai, K., 667. Fukaya, S., **310, 515.** Fukushi, T., 167, 265. Fullaway, D. T., 212, 692. Fuller, M. E., 629. Fulton, R. A., 332, 739. Furugori, N., 53. Fyfe, R. V., 86.

Gäbler, H., 196, 344, 664. Gahan, A. B., 227, 403. Gaines, J. C., 141. Gaines, R. C., 276. Gambrell, F. L., 392, 548. García, S. Planes, 701. Gardner, T. R., 552. Garman, P., 15, 246, 723, 724, 725. Garthwaite, P. F., 338. Gassner, L., **38.** Gautier, T. N., **439.** Geijskes, D. C., 464. Georgi, C. D. V., 120. Gerasimova, A. A., 305. Gersdorf, E., 520. Gersdorff, W. A., 439. Ghabn, Abdel Aziz Ali Elsayed, 458. Ghadimi, D., **685.** Ghilarov, M. S., 106, 200. Ghosh, C. C., 7, 294. Giavarini, I., 343. Gigante, R., 759. Gilliatt, F. C., **535.** Gilmore, J. U., 393. Gilyarov, M. S., 72, 356, 357, 368, 618. Gimingham, C. T., 101. Ginsburg, J. M., 21, 737. Girault, A. A., 502. Gleed, C. J., 383. Glover, L. C., 45. Glover, P. M., 311. Goetsch, W., 120. Goff, C. C., 64. Goffart, H., 461. Goidanich, A., 636. Goidànich, G., 119. Golding, F. D., 666. Golfari, R., 712. Goloviznin, D. D., 249. Gómez Clemente, F., 699, 702. Gomez-Menor, J., 10. González de Andrés, C., 700. Good, N. E., 64, 130. Gooden, E. L., 120, 282. Goodhue, L. D., 282, 321, 567, 695, 739. Goot, P. van der, 205. Goritzkaya, O. V., 471. Görnitz, K., 197, 522. Gösswald, K., 716. Götz, B., 154, 419. Götze, G., 197. Goux, L., 248. Gradojević, M., 685.

Graham, J. J. T., 331. Graham, S. A., 80. Grandi, G., 154. Granovsky, A. A., 385. Grassé, P. P., 413. Graves, H. W., 589. Gray, D. E., 538. Gray, R. A. H., 297. Green, E. E., 184. Green, W., 291. Greenslade, R. M., 613, 689. Greenwood, D. E., 562. Gregory, G. B., 448. Greimer, K., 347. Griot, M., 289. Grivanov, K. P., 73, 422, 423. Grosman, H. Francke-, 2, 252, 344, 762. Grünberg, A., 406. Gui, H. L., 399, 651. Guillaume, A., 182. Gunnell, F. H., 127. Gurney, W. B., 132. Gusev, V. I., 73, 432. Guseva, A., 427. Gwynn, A. M., 316, 376.

Hadden, F. C., 359. Hadersold, O., 420. Hadley, C. H., 727. Hadorn, C., 164. Hagmann, L. E., 565. Hahmann, K., 101. Haines, G. C., 202. Hale, H., **652.** Hall, D. G., 133. Haller, H. L., 120, 368, 567, 694. Ham, W. R., 772. Hama, T., 669, 670. Hambleton, E. J., 115, 502, 504. Hamilton, C. C., 542, 550. Hamilton, D. W., 489. Hammer, M., 155. Hammer, O. H., 561. Hampp, H., 522, 523. Hanf, M., 421, 522. Hanh, Nguyên-huu-, 188. Härdtl, H., 345. Hargreaves, E., 118, 239. Hargreaves, H., 372. Haring, M. M., 331. Harland, S. C., 82. Harman, S. W., 555, 574. Haroon Khan, M., 532. Harris, W. V., 112, 118, 642.

Harrison, C. J., 693. Hartman, G., 578. Hartzell, A., 173, 560. Hartzell, F. Z., 25, 130, 562. Harukawa, C., 53, 668, 769. Hase, A., 468, 519. Haseman, L., 79, 269. Hassan, A., 31. Hassan, A. A. G., 614. Haug, G. W., 390, 649. Hayhurst, H., 165. Headlee, T. J., 442. Heil, K. H., 234. Heinze, K., 68. Helm, A., 188. Hely, P. C., 754. Hem Raj, Bhalla, 180. Hempel, A., 502. Hengl, F., 518. Hennig, W., 716. Henrard, P., 315. Hepburn, G. A., 203. Hering, M., 247. Heriot, A. D., 171, 484. Hernandez Olmedo, I., 10. Herschler, A., 772. Herzig, J., 195. Hienton, T. E., 732. Higgins, J. C., 166. Hiley, J., 227. Hilgendorff, G., 188, 252, 522. Hill, C. C., 121. Hill, S. O., 587. Hille Ris Lambers, D., 440. Hinman, F. G., 26, 653. Hixon, E., 274. Ho, W. C., 387. Hodgkiss, W. S., 742. Hodson, A. C., 304. Hodson, W. E. H., 103, 349, 383. Hoerner, J. L., 281. Hoffmann, C. C., 248. Hoffmann, C. H., 394. Hofmann, C., 461, 660. Holaday, D. A., 330. Holbert, J. R., 391. Holst, E. C., 28. Hookom, D. W., 645. Hopf, H. S., 631. Hopkins, A. D., 1. Hori, M., 496, 770. Horn, W., 232. Hoskins, W. M., 131, 280. Hosny, M., 440. Hough, W. S., 271, 272, 554. Houser, J. S., 129, 399, 543.

Howard, Sir A., 666. Howard, L. O., 119. Howard, N. F., 739. Howes, F. N., 90. Howlett, F. M., 88. Hoyle, S. T., 367. Huber, L. L., 505. Huckett, H. C., 566. Humphreys, J. W., 645. Hunt, W. T., 275. Hus, P., 343. Husain, M. A., 180, 677, 686. Hussein, M., 147, 679. Husz, B., 254. Hutchins, R. E., 574. Hutchinson, H., 199, 364. Hutson, J. C., 358. Hutson, R., 733, 743. Hyre, R. A., 216.

Ignat'eva, Yu. V., 480. Ijima, K., 768. Il'ina, A. I., 306. Il'inskiĭ, A. M., 426. Illingworth, J. F., 214. Imai, S., 771. Imms, A. D., 46. Ingram, J. W., 28, 577. Inouye, M., 437, 452. Io (Chou), 641. Isaac, P. V., 243, 247. Isaakidès, C. A., 70, 71. Isely, D., 731. Ishii, G., **514.** Ishii, T., **296, 605, 671.** Ismailov, A., 472. Israel, P., 408, 602. Ito, K., 570. Ivanov, S. P., 157, 236, 237. Ivanova, N. A., 353, 482. Iwasa, T., 671.

Jacobs, S. E., 34, 118.
James, H. C., 105, 598.
Jancke, O., 2, 4, 52, 420, 761.
Janisch, E., 336, 419.
Jannone, G., 592.
Jarvis, H., 149.
Jary, S. G., 296.
Jaynes, H. A., 108, 390.
Jeannenot, J., 412.
Jehl, J., 522, 523.
Jenkins, C. F. H., 169.
Jepson, W. F., 317, 317.

Jerdon, T. C., 262. Jervis, L. A., 491. Jewet, H. H., 183. Johansson, E., 231. Johnpulle, A. L., 434. Johnson, H. W., 216. Johnson, J. P., 14, 723, 726. Johnston, C. J. R., 85. Johnston, H. B., 677, 679. Jones, E. T., 728. Jones, F. G. W., 1. Jones, H. A., 29, 120, 331, 368, 396, 739. Jones, L. S., 283. Jones, R. M., 571. Jones, S. C., 29, 60. Jones, S. E., 397. Jones, T. H., 589. Jourdan, M. L., 340, 500, 640. Judenko, E., 622, 719, 720.

Kadow, K. J., 587. Kalandra, A., 657. Kaloostian, G. H., 285, 292. Kalshoven, L. G. E., 262, 453, 706. Kamal, M., 456, 457. Kamel Bey, M., 682. Kamiński, E., 255, 256. Kamiya, K., 669. Kariya, S., 672. Karumidze, S., 481. Kaston, B. J., 14, 725. Katô, M., 603. Katsumata, K., 265. Katsura, S. K., 395. Kawada, T., 671. Kawecki, Z., 772. Kayashima, I., 52. Kayumov, S., 249. Kearns, H. G. H., 708. Keen, F. P., 400. Keenan, W. N., 23. Keifer, H. H., 650. Kelly, E. G., 184. Kelsall, A., 540. Kemp, H. J., 590. Kemper, H., 37, 48, 519. Kennedy, J. S., 6. Key, K. H. L., 583. Khan, M. Haroon, 532. King, C. B. R., 240, 337, 639. King, K. M., 170, 440, 590. Kintzel, W., 533. Kirchner, H. A., 660.

Kirichenko (Kiritshenko), A. N., 48, 305, 307, 560. Kirkpatrick, T. W., 177. Kiyama, R., 310. Klee, H., 619. Klein, H. Z., 405, 637. Klem, M., 248. Knechtel, W. K., 610. Knight, H. H., 247. Knowlton, G. F., 127, 128, 141, 247, 304, 369, 397, 496, 696, 772. Koblitsky, L., 441. Koehler, W., 523. Köhler, P., 146. Koidsumi, K., 207, 517. Kokina, E., 76. Koller, R., 38. Kondo, T., **496.** Kôno, H., **452, 516, 604, 673.** Konschak, M., 313. Korab, I. I., 159, 236. Koreishi, K., 206. Koreishi, T., 263. Kornfeld, A., 721. Kosmachevskii, A. S., 430. Kosobutzkii, M. I., 94. Kosswig, W., **419, 496.** Kotthoff, P., **101.** Kozhanchikov, I. V. (Kozhantshikov, I. W.), 351, 423, 440, 472, 475. Kozlova, E. N., 426. Kraĭter, A. D., 481. Kramer, 94. Krauss, N. H., 492. Kreibohm de la Vega, G. A., 362, 704. Kreiter, E. A., **424.** Krishnamurti, B., 409. Krüger, F., 251. Krukoff, B. A., 318. Kumashiro, S., 53, 54, 668, 672. Kunkel, L. O., 385. Kuntze, H. A., 560. Kuroda, S., 672. Kurosawa, M., **673.** Kurosawa, T., 515. Kuwayama, S., 309. Kylasam, M. S., 531.

Laan, P. A. van der, 603. Ladell, W. R. S., 630. La Due, J. P., 574. Laduizhenskaya, L. A., 476. Laffond, P., 371. La Forge, F. B., 120, 368, 651, 694. Laing, J., 150. Lal, K. B., 261. Lamas, C. J. M., 626. Lambers, D. Hille Ris, 440. Landaluze, P. Urquijo, 695. Landis, B. J., 13, 369. Lane, M. C., 569. Lang, A. L., 391. Lange, W. H., 117. Lange, jr., W. H., 226. Langford, G. S., **541.** Larchenko, K., 474. Larson, A. O., 648, 653. Larter, L. N. H., 490. Lathrop, F. H., 732. Laubert, R., 719. Laue, G., 101. Lavrekhin, F. A., 90. Lawson, F. R., 322. Leach, J. G., 43, 217, 386. Leefmans, S., 466. Legleu, R., 531. Lehmensick, R., 196, 460. Leib, E., 718. Lepage, H. S., 367. Lepesme, P., 184, 412, 501, 591, 624. Lepigre, A., **413.** Lesne, P., 416. Lesnikovskiĭ, A. Ya., 478. Lester-Smith, W. C., 360. Lever, R. J. A. W., 334, 497, 702. Levitt, E. C., 334. Lewin, C. J., 500. Lewis, H. F., 145. Li (Shou-sing), 244. Liebermann, J., 147, 655. Liebers, R., 196, 460. Light, S. F., 219. Lilly, J. H., 737. Lima, A. da Costa, 319, 704. Lindgren, D. L., 116, 575. Lindinger, L., 101. Linton, H. L., 137. Liro, J. I., 201. List, G. M., 28. Little, V. A., 749. Livingstone, E. M., 396. Liw (Tiou Fah), 244. López Cristóbal, U., 30, 289, 290. López Mansilla, E. E., 290. Lord, F. T., 536. Loughnane, J. B., 47.

Love, S., 368.
Lowman, M. S., 749.
Lozina-Lozinskiĭ, L. K., 257.
Luginbill, P., 696.
Luk'yanovich, F. K., 351, 475.
Luk'yanovich, L. M., 618.
Lund, H. O., 545.
Lupo, V., 496.
Lyle, C., 574.
Lyubischev, A. A., 94.

Mabry, J., **508.** McCauley, W. E., 748. McCormick, F. A., 749. MacCreary, D., 556. Macedo, W. de, **483**. Macfarlan, J., **462**. McGarr, R. L., **277**. McGough, J. M., 493. McGregor, E. A., 173. McHargue, J. S., **703.** Machatschke, J. W., **715.** McIndoo, N. E., **332.** McKenzie, H. L., 116, 226, 469. Mackie, D. B., 18, 116, 388, 747. Mackinnon, D. L., 598. McKinstry, A. H., 366. McLean, H. C., **140, 750.** MacLeod, G. F., **284, 542, 624.** McLeod, J. H., **537, 623.** McPhail, M., 142. Madden, A. H., **40, 132, 144.** Madel, W., **164.** Maercks, H., 421. Magerstein, V., 718. Mail, G. A., 26. Makeeva, E. A., 428. Maki, T., **437.** Makino, M., **560.** Malenotti, E., 713, 714, 757, 758. Mallis, A., 129, 588. Malysheva, E. W. Puzanova-, 304. Mamet, R., 66. Mani, M. S., **89, 440.** Manoff, I., **705.** Manolache, C. I., 193. Mansilla, E. E. López, **290.** Mansour, K., **248.** Manter, J. A., **726.** Marais, S. J. S., **684.** Marchal, P., 119. Marchionatto, J. B., 581. Marcovitch, S., 216, 509. Marcus, A., 235. Marcus, B. A., 69.

Marelli, C. A., 772. Markwood, L. N., 24, 368, 651. Marlowe, R. H., 213. Márquez, V. Moreno, **698.** Marsh, F. L., **330.** Marsh, R. W., 708. Marshall, Sir G. A. K., 106, 405, 612. Marshall, G. E., 732. Marshall, James, 174. Marshall, Joseph, 199, 364. Martelli, G. M., 5, 201, 260. Martelli, M., 712. Martin, H., 370, 708. Martin, J. P., 585. Martin, J. T., 616, **631, 772.** Martinez, M. Benlloch, **697.** Martorell, L. F., **288.** Mason, A. C., 84. Mason, H. C., 332. Mason, T. G., 65. Massee, A. M., 45, 103, 612, 613, 689. Matkovskiĭ, S. T., 75. Matsumoto, K., 310, 671. Maulik, S., **499.** Maxwell, C. W. B., **536.** Maxwell, J. M., **136.** Maxwell-Darling, R. C., 198. Mayo, J. K., 667. Mayton, E. L., 691. Medler, J. T., 137. Medvedeva, V. I., 473. Mégnin, J. P., 94. Meikle, A. A., 462. Melampy, R. M., **624.** Melis, A., 5. Mel'nichenko, A. N., 429. Melville, A. R., 592. Mendes, D., 642. Mendes, L. O. T., 82, 145. Mendizábal Villalba, M., 699, 700. Menor, J. Gomez-, 10. Menozzi, C., 161, 184, 527. Menusan, jr., H., **140, 284, 564.** Merino, G., **675.** Merkel, L., 101. Merkul'eva, A., 425. Merritt, J. M., 79, 733. Merwe, C. P. van der, 411. Metalnikov, S., 619. Metcalf, Z. P., 184. Meyer, T., 147. Meyers, M. T., 367, 505. Meyrick, E., 68, 151. Michael, A. D., 94.

Michael, H., 347. Michel, M. l'Abbé, 316. Michelbacher, A. E., 117, 280, 625, 650. Michelmore, A. P. G., 683. Miestinger, K., 234. Miles, H. W., 383. Miller, D., **58.** Miller, L. W., **168.** Miller, N. C. E., **198.** Miller, W. L., 367. Millot, J., 184. Milnthorpe, W., 111. Milum, V. G., 746. Mimeur, J. M., 48, 696. Minakawa, S., 515. Minkiewicz, S., 99, 256. Misaka, K., 263. Mischling, 255. Mishima, R., 669. Mistikawy, A. M., **687.** Mitchener, A. V., 540. Mitsuhashi, H., 53. Mo, Tjoa Tjien, 110, 358. Moniez, 94. Monte, O., 115. Monzen, K., 672. Moore, J. B., 25, 555, 562. Moore, R. H., 224. Moore, William, 567, 741. Mordvilko, A. K., 106. Moreno Márquez, V., 698. Morgan, W. L., 210, 334, 335, 496. Morgenthaler, 343. Mori, N., 771. Morison, G. D., 418. Morley, A. M., 298. Moroshkina, O. S., 617. Morris, H. M., 107. Morris, K. R. S., 103, 197, 341. Morris, V. H., 402. Morrison, H., 138. Morstatt, H., 248. Mossop, M. C., 203, 608. Motooka, K., 311, 438. Moutia, L. A., 176, 204. Muesebeck, C. F., 330. Mühlow, J., 154. Muller, H. R. A., 765. Müller, K. R., 520. Mundinger, F. G., 130. Munger, F., **575.**

Mungomery, R. W., 56. Munro, H. A. U., 541. Munro, H. K., 602. Munro, J. W., 177. Murat, M., 755. Murayama, J., 207, 436, 767. Murillo, L. M., 403. Murphy, P. A., 47, 495. Murray, G. H., 435. Musgrave, A. J., 598. Muzichenko, Yu. O., 184.

Nagel, C. M., 387. Nakamura, S., 604. Nakayama, S., 243, 767. Nanta, 205, 206. Narayanan, B. T., 360. Narayanan, E. S., 498, 532. Nardo, A. de, 469. Nawa, U., 53. Neiswander, C. R., 398, 505, 729. Neiswander, R. B., 399, 735. Nejedlý, J., 658. Nekrasova, Z. N., 481. Nelson, F. C., 109. Nelson, O. A., 331. Nelson, R. H., 397, 552, 553. Neu, W., 530. Neumann, H., **194.** Newcomer, E. J., **544.** Newman, L. J., 458, 754. Neves, M., **119.** Nevkrita, O. M., **184.** Nevskiĭ, V. P., **479.** Nguyên-huu-Hanh, 188. Nicholas, J. E., **488.** Nickels, C. B., **138.** Nikiforov, A. M., **6.** Nikol'skiĭ, V. L., **352, 431, 432.** Nikol'skiĭ, V. V., 470, **530.** Nishikawa, Y., 695. Nitsche, G., 419, 496. Nixon, G. E. J., 89, 688. Noble, L. W., **275.** Noble, N. S., **440, 450, 582.** Noda, I., **768.** Nonell Comas, J., 696. Nosova, M. I., **474.** Novikova, N. G., **482.** Nowicki, S., 72. Nunberg, M., 4, 154. Nutman, F. J., **183.**

Obarski, J., **72.**O'Connor, B. A., **495.**Oestlund, O. W., 484.
Ohta, Y., **771.**

Oishi, T., 311. Okada, I., 54, 310, 438, 606. O'Kane, W. C., 20, 45. Okazaki, K., 368, 770. Olmedo Hernandez, I., 11. Oman, P. W., 288. O'Neill, W. J., **555, 750.** Ong, E. R. de, **282.** Oosthuizen, M. J., 202. Oozi, Y., 438. Orchard, O. B., 711. Orelli, O. Schneider-, 459. Örösi-Pál, Z., 344, 664. Orr, L. W., 43. Osborn, H. T., 12, 586. Osburn, M. R., 285. Oshanin, V. F., 78. Ossiannilsson, F., 232. Ostrovskii, N. I., 422. Ota, Y., 265. Otero, J. I., 116. Oudemans, 94, 95. Owen, jr., W. L., 276. Ozol, A. A., 474. Ozols, E., 256.

Packard, C. M., 78. Paillot, A., 164, 192, 464. Painter, R. H., 121. Pál, Z. Örösi-, 344, 664. Palma, P., 757. Panasyuk, M. P., 235. Pankova, I. A., 601. Panshin, A. J., 318. Pantsios, A. A., 695. Paoli, G., 162, 757. Pape, H., 660. Parfent'ev, V. Ya., 74, 432. Park, T., 439. Parker, J. R., 676, 680, 686. Parker, L. B., 108. Parkhomenko, V. Yu., 158. Parkin, E. A., 635. Parmelee, F., 733. Parsons, F. S., 197, 199, 364. Pasquier, R., 685. Patay, R., 368. Patch, E. M., 400. Patch, L. H., 367, 729. Patterson, N. A., 534. Paul, L. C., 440. Pavlov, I. F., 355. Payne, N., 12. Pearce, G. W., 489.

Pearson, E. O., 365.

Peat, J. E., 366. Pelton, J. Z., 624. Pemberton, C. E., 584, 585. Pepper, B. B., 565. Pereira, C., 504. Perret, J., 417. Pescott, R. T. M., 84, 168. Petch, C. E., 440. Petch, T., 152, 153. Peterka, 255. Peters, G., 695. Peterson, A., 390. Petherbridge, F. R., 191, 462, 463. Petrik, A. I., **428.** Petty, B. K., 203. Peus, F., 48. Peyerimhoff, P. de, 410. Pfeffer, A., **657.** Phillips, E. W. J., **635.** Phillips, W. J., 304. Phillis, E., 65. Phipers, R. F., 616. Pichler, F., 715. Picket, A. D., 441. Pickles, A., 10, 369, 404, 533, 752. Piemeisel, R. L., **322.** Pieri, A., **758.** Pierpont, R. L., 551. Pilyugina, S. A., 76. Pinckney, J. S., 23. Piper, jr., W. R., 746. Planes García, S., 701. Plassche, A. W. van de, 465. Plessis, C. Du, 676. Plumb, G. H., 18, 547. Plummer, C. C., **505.** Poeteren, N. van, **299.** Poisson, R., 343. Polak, R. A., 529. Polivka, J. B., 399. Polizu, S., 71, 119, 192. Pollard, A. G., 166. Pollister, P. F., 184. Poluzzi, C., 715. Polzik, A. I., 480. Ponce, A., 181. Ponomarenko, D. A., 76. Ponomarev, A. V., 477. Poor, M. E., 65. Poos, F. W., 386, 609. Popov, S. L., 428. Popova, A. I., 480. Popp, H. W., 772. Portères, R., 531. Portman, R., 291. Pospelova, V., 425.

Potter, C., 496.
Powell, P., 369.
Powell, jr., T. E., 363.
Prendergast, D. T., 283.
Priesner, H., 440, 457.
Principi, M. M., 189.
Printz, Ya. I., 429.
Proeschel, Mlle., 182.
Pruthi, H. S., 8, 88, 181, 241, 340, 498, 765.
Pulley, G. N., 695.
Pushin, F. E., 482, 483.
Pussard, R., 301.
Pustovoit, A. F., 75.
Putman, W. L., 509.
Puzanova-Malysheva (Puzanowa-Malyschewa), E. W., 304.
Pyatnitzkii, G. K., 472, 473.

Quayle, H. J., **722.** Queiroz, J. B. R., **687.**

Rahman, K. A., 87. Raichoudhury, D. P., **34, 118.** Rainwater, C. F., 644. Raj, Bhalla Hem, 148. Ramakrishna Ayyar, T. V., 89, 240, 360. Ramchandra Rao, Y., 628, 678, 683. Rau, G. J., 226. Rau, S. Ananda, 148. Raucourt, M., 524, 525. Read, W. H., 711. Reed, M. P., 652. Reed, W. D., 396. Reeks, W. A., 77, 247. Regnard, P., 317. Regnier, R., 229. Regueral, F. G., 702. Rehn, J. A. G., 588. Rehn, J. W. H., 588. Reichmuth, W., 37. Reinhard, H. J., 279, 560. Reinmuth, E., 660. Rekk, G. F., 305. Reppert, R. R., 136. Reyne, A., 499. Rice, E. M., 772. Rice, P. L., 444. Richardson, H. H., Richey, F. D., 505. Richmond, H. A., 486. Riechen, F., 194.

Rieth, G., 346. Ripley, L. B., 202, 203. Ritcher, P. O., 304, 368, 736. Rivnay, E., 379. Roark, R. C., 48, 304, 624, 651. Roba, R. P., 533. Robbie, W. A., **304.** Roberti, D., 696. Roberts, R., 608. Robinson, J. M., 691. Rodendorf, B., 303. Rodionov, Z. S., 96. Roebuck, A., **341.** Romm, H. J., 217. Ronna, A., 115, 655. Roos, K., **459.** Rose, W. G., **694.** Rosenthal, H., 528. Ross, E. S., 650. Ross, H. H., 293, 573. Rossi, L., **593.** Roubaud, E., **410.** Roy, B. K. Datta, 772. Roy, D. N., 184. Rozsypal, J., 620. Rubtzov, I. A., 426. Rude, C. S., 274. Ruffo, S., 712. Ruhmann, M. H., 59. Ruisakov, S. V., 479. Ruiz Castro, A., 700, 701. Rumyantzev, M. D., 476. Rungs, C., 46, 500, 641. Rusanova, V. N., **617.** Rusk, H. W., **320.** Russell, G. A., 749. Russo, G., 119, 350. Rzhechitzkaya, Yu., 423.

Saalas, U., 518.
Sachtleben, H., 233.
Saito, K., 264, 310.
Saito, S., 247.
Sakai, K., 604.
Sakai, S., 208.
Sakimura, K., 142, 213, 692.
Salman, K. A., 394.
Salt, G., 102.
Sameshima, T., 439.
Samuel, C. K., 88.
Sandhouse, G. A., 369.
Santis, L. de, 290.
Saraiva, A. Coutinho, 315
Sattler, F., 421.
Sauer, H. F. G., 333, 689.

Sauter, H., 119. Savchenko, E. N., 236, 237. Sawamoto, T., 673. Sazama, R. F., 732, 732. Scales, A. L., 279. Scaramuzza, L. C., 171. Schedl, K. E., 528, 696. Scheibe, A., 251. Schilder, F. A., 601. Schimitschek, E., 154, 312, 663, 664. Schipper, A. L., 304. Schmidt, C. T., 143. Schmidt, H., 717. Schmitt, H., 521. Schneider-Orelli, O., 459. Schoene, W. J., 271, 556, 744. Scholl, E. E., 139. Schonwiese, F., 659. Schread, J. C., 14. Schuch, K., 91, 195, 717. Schulze, H., 94. Schüssler, H., 68. Schwerdtfeger, F., 194, 196, 253, 312, 368. Scudder, S. H., 744. Seamans, H. L., 751. Šeda, A., 656, 658. Seiferle, E. J., 387. Seín, jr., F., 223. Seitner, M., 663, 664. Sekiya, I., 265. Selivanova, S., 428. Semenov, A. E., 6, 475, 601. Sensburg, E., 519. Serdyukov, P., 74. Severin, H. C., **133.** Shabanova, M. P., **431.** Shabetai, C. R., **456.** Shafik, M., 591. Shands, W. A., 393. Shapiro, D. S., 257. Shaposhnikova, E., 479. Shaw, H., **613.** Shchegolev, V. N., 477. Shek, G. Kh., 471. Shepard, H. H., 389. Sherman, F., 575, 644. Shibuya, M., 670. Shinji, O., 496. Shiraki, T., 263, 309. Shirck, F. H., 747. Shuey, G. A., 216. Shumakov, E. M., 478.

Siegler, E. H., **321, 575.** Sievers, A. F., **749.**

Silver, E. A., 402. Silvestri, F., 161, 248, 260, 496. Simm, K., 518. Simmonds, H. W., 58, 59, 211. Simmons, P., 118, 573. Sindersberger, M., 235. Skoblo, I. S., 424. Skorkin, L. A., 306. Sleesman, J. P., 399. Smee, C., **104**, **113**, **203**. Smirnov, B. A., **306**. Smit, B., 151, 202. Smith, A. C., 318. Smith, A. J., 176, 177. Smith, C. F., 119, 247, 369, 696, 772. Smith, C. M., 120, 368. Smith, C. R., 183. Smith, F. F., 273, 396. Smith, G. L., 279. Smith, G. R., 725. Smith, J. B., 11. Smith, J. H., 246, 267, 494. Smith, K. M., 68. Smith, L. E., 575, 695. Smith, L. M., 62. Smith, R. C., 184. Smith, R. H., 387, 745. Smith, S. G., 215, 487. Smith, W. C. Lester-, 360. Snapp, O. I., 82. Snedecor, G. W., 688. Snelling, R. O., 63. Snitko, N. F., 306. Sobol', P. F., 235. Sokanowsky, B., 665. Solomon, M. E., 150. Sonan, J., 207, 263, 517. Sooter, C. A., 274. Soraci, F. A., 118. Sorauer, P., 703. Sorenson, C. J., 48, 126, 127. Soriano, P. S., 675. Sornay, P. de, 317. Spasskii, A. F., 357. Speicher, B. R., 119. Speyer, E. R., 349, 711. Speyer, E. R., 100, 346, 420, 716. Spoon, W., 766. Squire, F. A., 83, 175, 445. Staniland, L. N., 527. Stanley, G. W., 495. Stanley, W. W., 216. Stansfield, F. W., 152. Stark, V. N., 354, 428. Stearns, L. A., 443, 545, 556.

Steenburgh, W. E. van, 535. Steer, W., 297, 613, 614. Steiner, H. M., 287, 557. Steiner, L. F., 732. Steiner, P., 662. Steinitz, H., 636. Stellwaag, F., 154, 419, 661. Stepantzev, I. N., 94. Stevens, N. E., 386. Stevens, W. L., 166. Stiles, C. F., 139. Stocks, J. B., **625**. Stoddard, E. M., **15**, **723**. Stofberg, F. J., **377, 455.** Stone, W. M., **569.** Storey, H. H., **756.** Stracener, C. L., **218.** Straňák, F., **656.** Strand, A. L., **43.** Strawinski, K., 98. Stringfield, G. H., 505. Stritt, W., 560. Strong, L. A., 23, 323. Stultz, H. T., 540. Subklew, W., 496. Subrahmaniam, T. V., 204, 606. Suehiro, A., 213. Suenaga, H., 669. Sugimoto, M., 436. Şuire, J., **415.** Šulc, K., 560. Sullivan, W. N., 29, 567, 739. Summers, E. M., 577. Sung (Tsu-lien), 244. Sviridenko, P. A., 474. Svoboda, J., 255. Swain, A. F., 117. Swain, R. B., **133, 291.** Swan, D. C., **584, 703.** Swank, G. R., 217. Swezey, O. H., **213, 692.** Swingle, H. S., 82, **690, 741.** Swingle, M. C., **363.** Szelényi, G. von, **526.** Szymański, W., 97, 98, 300.

Taguti, K., 768.
Takagi, G., 264.
Takahashi, H., 769.
Takahashi, R., 48, 119, 262, 367, 496, 624, 674.
Takahashi, S., 673.
Takano, S., 263, 294, 768.
Takio, M., 515.
Takizawa, M., 304.

Tammes, P. M. L., 685. Tams, W. H. T., 405. Tamura, I., 516, 605. Tanabe, C., 265. Taranukha, M. D., 156. Tate, H. D., 65, 225. Tattersfield, F., 536, 616, 631. Taylor, G. G., 45, 46. Taylor, T. H. C., 83, 372, 376, 497. Taylor, W. L., **633.** Tesch, B., **38.** Thalenhorst, W., 461. Théry, A., 118. Thiem, H., 233, 345, 346. Thomas, C. A., 488. Thomas, F. L., 276. Thomas, H. R., **577.** Thomas, I., 191, 382. Thompson, B. G., 398. Thompson, R. W., 536. Thompson, W. L., 727. Thorpe, W. H., 1. Tiggelovend, L. M. J., 262. Timofeeva, T. V., 305. Timokhina, N. I., 482. Tischler, W., 51. Tissot, A. N., 696. Tjoa Tjien Mo, 110, 358. Todd, J. N., **644.** Tokunaga, M., 769. Tomaszewski, W., 522. Tomlinson, W. E., 576. Tompkins, C. M., 577. Torre-Bueno, J. R. de la, 11. Torres, D. Delgado de, 699. Townsend, J. F., 17, 725. Toyoshima, A., 771. Trägårdh, I., 155, 440, 611, 624, 634. Trappmann, W., 636. Trehan, K. N., 772. Treiman, F. S., 158. Tsai, P. H., 674. Tucker, R. W. E., 221, 446, 448. Turnbull, J., 440. Turner, N., 14, 15, 17, 247, 722, 723, 724. Twinn, C. R., 560. Tyumeneva, V. A., 477.

Uchida, H., 263, 436. Uchida, T., 54, 119. Ugarov, A. A., 470, 530. Uhlmann, E., 300. Underhill, G. W., 271. Urquijo Landaluze, P., 695. Uspenskaya, N. V., 479. Utter, L. G., 273. Uvarov, B. P., 72, 111, 677.

Vance, A. M., 730. Van den Brande, J., **69, 193.** Van den Bruel, W. E., 70, **313.** van de Plassche, A. W., 465. van der Goot, P., 205. van der Laan, P. A., 603. van der Merwe, C. P., 411. Van der Vecht, J., 674. van der Westhuysen, J. P., 203. van Poeteren, N., 299. van Steenburgh, W. E., 535. van Zwaluwenberg, R. H., 492. Vappula, N. A., 120, 201, 695. Vasil'ev, K. A., 307. Vasser, R. E., 470. Vayssière, P., 501. Vecht, J. Van der, 674. Vega, G. A. Kreibohm de la, 362, 704. Veitch, R., 703. Vejabhongse, N. P., **149**. Venables, E. P., **144**, **171**, **485**. Venkitasubban, C. S., 452. Vesey-Fitzgerald, D., 433, 611. Vidal, J., 340. Vielwerth, V., 656. Viennot-Bourgin, G., 259, 525. Vigor, S. H., **590.** Villalba, M. Mendizábal, **699.** Villiers, F. J. De, 681. Vinokurov, G. M., 353, 357. Vinson, J., **454.** Vitzthum, H., 94. Vivien, J. H., 314. Voelkel, H., 197, 248. Voitenko, M. P., 354. Volgin, V. I., 424. Volkonsky, M., **180, 239, 410, 590.** Volkov, V. F., **353.** Volodina, G., 423. von Szelényi, G., 526. von Winning, E., 368. Vore, Ll. De, 772. Vorontzov, A., 429. Voûte, A. D., 311, 336, 337, 639. Vuirzhikovskaya, A. V., 473.

Wadley, F. M., **39, 696.** Wagner, G. B., **118, 321, 362.**

Vukasović, P., 720, 759, 760.

Wakeland, C., 65, 747. Walden, B. H., 18, 247. Walkden, H. H., 135, 292. Walker, G. L., **285.** Walker, H. G., 13. Walker, M. G., 102. Wallace, C. R., **404.** Wallace, H. E., 445. Wallace, R., 210. Wallengren, H., 230. Walters, E. A., **121.** Walton, C. L., 462, **709.** Walton, R. R., 138. Wang (Fei-peng), **244.** Warburton, C., 296. Ward, K. M., **85, 494.** Watanabe, C., **247, 296, 452.** Watanabe, F., 208. Waterston, J. M., 220, 626. Watson, M. A., 32, 342, 615. Watson, J. R., 131, 727. Watts, J. G., 278, 644. Watzl, O., 662. Webb, jr., J. E., 30. Weber, A. L., 140, 750. Webber, R. T., 589. Webster, R. L., **488.** Weidner, H., **120, 193, 196, 439,** 469. Weigel, C. A., 553. Welch, H., 609. Welsch, J., 251. Went, J. C., 656. Westgate, W. A., 45. Westhuysen, J. P. van der, 203. Whitehead, F. E., 138, 398. Whitehead, T., 379. Whitney, R. J., 763. Wickens, G. M., 377. Wiesmann, R., **156**, **300**, **459**. Wilcoxon, F., 330. Wilford, B. H., 80. Wilkinson, D. S., **106, 342.** Willard, H. F., **84.** Willaume, F., 525. Wille, J. E., 81, 362, 533, 626. Williams, J. L., 440, 608. Wilmot, R. J., 439. Wilson, G. F., 349, 709. Wilson, J. W., **64.** Winning, E. von, **368.** Wirth, H., **665.** Woglum, R. S., 702. Wolcott, G. N., 222, 288. Wolff, N. L., **518.** Wood, A. A., 540.

Wood, M., 287.
Woodside, A. M., 507.
Woodworth, C. W., 78.
Worsley, R. R. LeG., 34, 37, 183, 258.
Worthley, H. N., 287, 488, 742.
Wright, D. W., 298, 463.
Wu (Chen-chung), 245, 304.

Yagi, N., 770. Yago, M., 53, 368, 604. Yakhontov, V. V., 238, 247, 358. Yamada, S., 671. Yamada, Y., 770. Yamashita, S., 671. Yamazaki, T., 264, 516, 669. Yang (Hung-ju), 245. Yarmolenko, I. M., 160, 236. Yashiro, H., 605. Yatomi, K., 671. Yeager, J. F., 439. Yokoo, T., **768.** Young, H. D., **321, 362.** Yuasa, H., **264, 497, 670.** Yusope, M., **31.**

Zacher, F., 94, 194, 248, 252, 301, 468, 706.

Zakhvatkin, A. A., 94, 95.

Zappe, M. P., 14, 15, 17, 723, 724.

Zehntner, L., 89.

Zhitkevich, E. (O.) N., 157, 236.

Zhukovskiĭ, A. V., 427.

Zhuravskaya, S. A., 472.

Zimin, G. S., 73.

Zimin, L. S., 246.

Zimmerman, E. C., 576.

Zolk, K., 461, 462.

Zolotarevsky, B. N., 679, 755.

Zotzenko, L. N., 478.

Zwaluwenberg, R. H. van, 492.

Zweigelt, F., 521.

GENERAL INDEX.

In the case of scientific names, the page reference is cited only under the heading of the generic name.

When a generic name is printed in brackets, it signifies that the name is not the one adopted.

abbreviatus, Diaprepes.

abdominalis, Macrocentrus; Thrips. aberrans, Doryphorophaga; Hetero-

tylenchus; Pyrilla.

Abies spp., pests of, in Britain, 191, 527; Chermes nordmannianae on, in Czechoslovakia, 657; Aphids on, in Germany, 3, 252, 344, 660; pests of, in Japan and Sakhalin, 452, 516; pests of, in U.S.A., 369, 581; susceptibility of species of, to C. piceae, 344; Dendrolimus spectabilis not maturing on, 672; Sirex cyaneus associated fungus in timber of, 191.

abietana, Argyroploce (Olethreutes). abietella, Dioryctria. abietina, Pristiphora (Nematus). abietinum, Rhopalosiphum (Neomy-

zaphis).

abietinus, Mindarus.

abietis, Chermes (Adelges, Sacchiphantes); Cryphalus; Hylobius. abietum, Nematus (see Pristiphora abietina).

abnormis, Tanaomastix (Leptomastidea).

abruptorius, Exenterus.

Abutilon, Dysdercus superstitiosus

on, in Sudan, 512.

Abutilon indicum, not infested with Platyedra gossypiella in Antigua, 446; P. gossypiella on, in Java, 766; cotton stainers on, in Uganda, 374.

abutilonea, Trialeurodes.

Abyssinia (see Africa, Italian East). Acacia, pests of, in Australia, 8, 449; Icerya purchasi on, Formosa, 263.

Acacia confusa, Bruchid on, in Formosa, 310; Anomala xantho-pleura on, in Loochoo Islands, 516.

decurrens, Acanthopsyche Acacia junodi on, in S. Africa, 203.

Acacia mollissima, Acanthopsyche junodi on, in S. Africa, 203.

Acacia, Yellow (see Caragana).

Acacia, White (see Robinia). acaciaria, Cleora (Boarmia).

Acalypha indica, insecticidal properties of, 360.

speisevi, Acanthoplus attacking Nomadacris septemfasciata Nyasaland, 114.

Acanthopsyche griseoalba, on Nipa fruticans in Indo-China, 184.

Acanthopsyche junodi, measures against, on Acacia spp. in S. Africa, 203.

Acanthopsyche nipae, on Nipa fruticans in Indo-China, 184.

Acanthoscelides (see Bruchus).

Acanthostichus, predacious on Gasterocercodes brasiliensis Brazil, 504.

Acanthostome insidiator, bionomics of, in Japan, 310.

Acarapis dorsalis, on honey-bees in Italy, 343.

Acarapis externus, possibly on honey-bees in Italy, 343.

Acarapis woodi (attacking honeybees), in Brazil, 655; in Italy, 343.

Acer (see Maple).

Acer pseudoplatanus, pests of, in France, 316.

Aceratagallia californica, acquiring virus of beet curly-top, 445.

Aceratagallia sanguinolenta, on narcissus in U.S.A., 288.

Acetic Acid, attraction of, for Cydia spp., 135, 137.

Acetone, as a solvent, 29, 50, 183, 574, 739, 740, 749; in baits for Cydia molesta, 135.

Achaea janata, on castor in Ceylon, 359.

Achroia grisella, parasitised by Nemeritis canescens, 1, 2.

Acicnemis vehemens, on jelutong in Malaya, 109.

Acidia lucida, on Lonicera in Switzerland, 156. acidusa, Anastrepha.

Acompsia pseudospretella (see Borkhausenia).

Acontia graellsi (on cotton), in Madagascar, 67; parasite of, in Nyasaland, 114; in Sudan, 513; in Transvaal, 67.

Acontia transversa, bionomics and control of, on Hibiscus sabdariffa in Netherlands Indies, 358.

Acrobasis caryae, on pecan in U.S.A., 138, 587; measures against, 138.

Acrobasis caryivorella, on pecan in Florida, 587.

Acrobasis juglandis, bionomics of, on pecan in Florida, 587.

acrobates, Telenomus.

Acrocercops, bionomics of, on chestnut and oak in Japan, 669.

Acrocercops bifasciata, on cotton in Belgian Congo, 315.

Acrocercops chalybophanes, sp. n., on coffee in Tanganyika, 68, 113.

Acrocercops tenera, on Schleichera trijuga in Ceylon, 359.

Acrolepia assectella, on onion in Germany, 39.

acroleuca, Conradtina.

Acromyrmex spp., tests of fumigants against, in Brazil, 533.

Acronicta psi, on apple and bush

fruits in Britain, 613.

Acronicta rumicis, on apple and bush fruits in Britain, 613; on poplar and willow in Germany, 461.

Acropolitis rudisana (rudis), on apple in Tasmania, 10.

Acrostalagmus, infesting Sipha flava in Porto Rico, 39.

Acrosternum hilare, in orchards in

U.S.A., 19, 127; bionomics of, 127.

Acrosternum spicatum, on Crotalaria anagyroides in Seychelles, 433.

Actia aegyptia, parasite of Earias spp. in India, 243.

Actia monticola, parasite of Laphygma exigua in India, 55.

Actinomyces scabies, Pnyxia scabiei associated with, on potato in Ontario, 540.

Acucephalus albifrons, taken on high buildings in U.S.A., 215.

aculealis, Tabidia.

aculeatus, Haplothrips; Hylesinus (Leperisinus).

acuminata, Aelia.

acupunctatus, Scyphophorus.

acuta, Leptocorisa. acutus, Platymetopius. Acyphas leucomelas, on vines in New South Wales, 334.

Adalia bipunctata, predacious on Aphids in Germany, 197; rearing of, on frozen Aphids in Ohio, 650. adamsi, Melasoma.

Adansonia (Baobab), cotton stainers on, in Uganda and Sudan, 374,

512.

Adelges (see Chermes). adherbal, Aporia crataegi.

Adhesives, banding with, **52**, **88**, **93**, **202**, **256**, **348**, **480**, **495**, **539**, **626**, **665**, **716**, **755**; comparison of, for banding against *Sexava* on coconut, **495**; methods of testing, **665**; other methods of catching insects with, **643**, **719**.

Adonia variegata, predacious on Aphids and Lepidoptera in

Ukraine, 353.

adonidis, Entomoscelis. adonidum, Pseudococcus.

Adoretus sinicus, measures against, on beans in Hawaii, 752.

Adoretus tenuimaculatus, food-plants of adults of, in Japan, 54.

Adrastus limbatus, soil conditions in relation to, in Russia, 425.

adspersa, Epicauta. adspersus, Exenterus.

advena, Aenasius; Ahasverus (Ca-thartus).

advens, Phyllocoptes.

Aegeria bibionipennis, measures against, on strawberry in Oregon, 291.

Aegeria exitiosa, on peach in U.S.A., 19, 271, 323, 726; measures against, 271, 323.

Aegeria opalescens, on peach in California, 20.

aegeus, Papilio.

Aegilops spp., relation of Mayetiola destructor to, in U.S.A., 728.

aegyptia, Actia.

aegyptiaca, Thermobia.

aegyptiacus, Retithrips; Scaphoideus.

aegyptium, Anacridium.

Aelia acuminata (on cereals), in Czechoslavakia, 659; bionomics of, in Germany, 51, 52; in Switzerland, 91; type of injury by, 51; characters of, 51.

Aelia rostrata, bionomics and control of, on wheat in Spain, 697,

698; in Portugal, 697.

Aenasius, revision of, 212; records of distribution and use of species of, including new ones, 212.

aenea, Anomala dubia; Dicerca.

aeneoviridis, Catolaccus.

aenescens, Naranga.

aeneus, Chalcodermus; Corymbites (Selatosomus); Meligethes.

Aenoplex carpocapsae, utilisation of, against Cydia pomonella in Idaho, 323, 747.

aequalis, Euaresta; Hemiliopsis.

aequatus, Rhynchites.

aereus, Monodontomerus.

Aerodromes, termite mounds

danger in, 449.

Aeroplanes, for broadcasting baits for grasshoppers, 398; for distributing insecticidal dusts, 19, 235, 398, 426, 433, 508, 654, 682, 748; for applying oil sprays, 134, 748; for distributing dusts against flying locusts, 681.

Aeropus sibiricus, distribution of, in Russian Union, 353, 426; bionomics of, 426; value of ploughing to destroy egg-pods of, 353.

aeruginosus, Limonius.

Aeschynomene sensitiva, cidal properties and constituents of, 224.

Aesiotes notabilis, on Araucaria cunninghami in Queensland, 494.

aestivum, Apion. aestivus, Rhizotrogus.

aethiopica, Tachina (Exorista).

aethiops, Orphinus.

affinis, Dasyneura; Neoaplectana; Pempheres.

Afghanistan, Cydia pomonella in apples from, 340.

afra, Spalangia.

Africa, locust problems in, 111,

679, 683, 684, 687.

Africa, French West, locusts in, 112, 684, 755; Trypetids in, 492 Aphis laburni and disease of ground-nuts in, 531; beneficial insects in, 112, 531.

Africa, Italian East, Aphis laburni and rosette disease of groundnuts in, 592; Coccids in, 496, 757; parasite of Aspidiotus destructor on banana in, 641; locusts in, 111, 684; survey of entomological problems in, 260.

Africa, North, Coccids of, 559, 696. Africa, Portuguese East, Coccid on Citrus in, 377; Diparopsis castanea in, 199; locusts in, 676,

678, 682, 683, 687.

Africa, South, Citrus pests in, 106, 176, 202, 377, 378, 455; Aonidiella aurantii on other plants in, 455; cotton pests in, 67, 177, 199, 364, 365, 366; forest pests

in, 68, 203; miscellaneous pests in, 364, 497; pests of leguminous crops in, 151, 292, 364; maize pests in, 106, 364, 411, 511; pests and disease of sugar-cane in, 411; Myzus persicae on tobacco in, 177; pests of stored grain in, 202; locusts in, 152, 411, 676, 681, 682, 683, 684, 687; termites in, 48, 411; new weevils in, 106, 612; beneficial insects in, 378, 411, 456, 610; Nosema causing disease in Cactoblastis cactorum in, 202; plant pest legislation in, 66, 691; beneficial insects introduced into Mauritius from, against Lachnosterna, 317; Halotydeus destructor probably introduced into W. Australia from, 582. Africa, South-west, locusts in, 676.

africana, Cassidocida; Gryllotalpa.

africanus, Lyctus.

Agallia constricta, on narcissus in U.S.A., 288. Agalliaphagus, parasite of Eutettix

tenellus in Utah, 128.

Agamermis decaudata, parasite of grasshoppers in U.S.A., 124.

Agar Agar, in mopping mixture against cotton boll weevil, 645. agavis, Chrysomphalus.

Agelastica coerulea, not attacked by Aiolocaria mirabilis in Korea, 310.

Ageneotettix deorum, destroying bluegrass in Iowa, 23.

Ageratum conyzoides, Bemisia gossypiperda transmitting disease to tobacco from, in India, 242.

agilis, Pristomeridia (Pristomerus). agilis, Can., Eberhardia (Cosmoglyphus).

agilis, Mich., Eberhardia (see E. michaeli).

Aglaope infausta, bionomics of, on fruit trees, etc., in Spain, 697; barium fluosilicate ineffective against, 697.

Agonoxena argaula, on coconut in Fiji, 211.

Agria mamillata, bionomics of, in Ukraine, 302.

Agria monachae, parasite of Lymantria dispar in Jugoslavia, 760. Agrilus ghesquierei, sp. n., on Albizzia in Belgian Congo, 118.

Agrilus hastulifer, on cork oak in Morocco, 46.

Agrilus mediocris, on Citrus Ceylon, 359.

Agrilus spinipennis, bionomics of, on elm and Zelkova in Japan, 265.

Agrilus viridis, on birch in Russia,

Agriotes gurgistanus, relation of, to type of soil in Russia, 72, 356; on rubber-producing plants, 72.

Agriotes lineatus, in Russian Union, 76, 424, 425, 426; on wheat and grasses, 76; soil conditions in relation to, 424, 425, 426.

Agriotes mencus, bacteria isolated

from, 624.

Agriotes obscurus, soil conditions in relation to, in Russian Union,

424, 425, 426, 429, 430.

Agriotes sputator, soil conditions in relation to, in Russian Union, 72, 356, 424, 425, 426; on rubber producing plants, 72.

Agromyza obtusa, in Burma, 451; bionomics of, on Cajanus cajan in India, 242, 451; measures

against, 451.

Agromyza phaseoli, bionomics and control of, on beans in New South Wales, 335.

pinguis, measures Agromyza against, on chicory in Belgium, 313.

Agromyza simplex (Asparagus Miner), bionomics of, in Britain,

Agromyza solani (Tomato Leafminer), bionomics and control of, in British Isles, 710.

Agropyrum ramosum, relation of wireworms to, in Central Asia, 76.

Agropyrum repens, pests of, in Australia, 494; relation of pests to, in Russian Union, 6, 257, 425, 426, 427, 428.

Agrotis longidentifera (see Euxoa). occulta, Tachina civilis ovipositing on, in Russian Union, 303.

Agrotis orthogonia (see Porosagrotis). Agrotis segetum (see Euxoa).

Agrotis vestigialis (see Euxoa).

Agrotis ypsilon, on potato in Hawaii, utilisation of parasites against, on maize in Netherlands Indies, 337.

Ahasverus advena, in stored cassava tubers in Madagascar, 455; in copra and stored rice in Malaya, 31; tests of insecticides on, 405, 610, 616.

Aholcus (see Telenomus).

aino, Tipula.

Aiolocaria mirabilis, bionomics of, in Korea, 310.

Aira flexuosa, Pentatomid on, in Germany, 51.

Alabama, pests of Citrus in, 690; cotton pests in, 327, 691; Chalcodermus aeneus on cowpeas in, 572, 691; miscellaneous pests in, 327, 690, 691.

Alabama argillacea (Cotton Leaf Worm), in U.S.A., 279, 329, 744; in W. Indies, 222, 752; Trichoparasitic on, 744: gramma 222, 329; measures against, tests of dusts against, 279.

Alaska, new Anobiid damaging

wood in. 214.

alaskensis, Pikonema (Pachynematus).

albella, Timocratica.

alberti, Eberhardia (Cosmoglyphus). albicincta, Pimpla (Exeristes).

albifrons, Acucephalus. albipalpus, Hemiteles. albipes, Technomyrmex. albisignata, Idiatella.

Albizzia spp., Indarbela quadrinotata on, in Assam, 601; Agrilus ghesquierei on, in Belgian Congo, 118: Xystrocera festiva on, in Java, 261.

albobasilaris, Chelonus. albocallosus, Bruchus. albohirtum, Lepidoderma. albolineatus, Dendrolimus.

albopictus, Elasmus; Lamachus; Xorides.

Alcides, in Indo-China, 185.

Alcides obesus, bionomics and control of, on kapok in Indo-China, 185.

Alcohol, for extracting insecticidal principles of plants, 34, 204, 206, 258, 360, 361, 412, 550, 740; spray formulae containing, 701.

Alcohols, Sulphated, uses of, in sprays, 286, 395, 396, 443, 704, 735, 736; contact performance of, 45.

Alder, Anomala chamaeleon on, in Korea, **767**.

Aleurites, Coccids on, in Nyasaland. 114.

Aleurites fordi, pests of, in Brazil. 502; pests of, in Burma, 340; Aspidiotus perniciosus experimentally developing on, 305.

Aleurites montana, pests Burma, **340**.

Aleurobius farinae (see Tyroglyphus).

Aleurocanthus woglumi, Eretmocerus serius controlling, on Citrus in Bahamas, 491.

Aleurodes azaleae, on azaleas in California, 20, 748; parasite of. 748.

Aleurodicus destructor, utilisation of Javanese parasite against, Celebes, 337.

Aleuroglyphus, gen. n., 95.

Aleuroglyphus ovatus, in stored cereals in Russian Union, 95, 96, 97; bionomics of, 96, 97.

aleurolobi, Eretmocerus.

Aleurolobus barodensis, on sugarcane in Bombay, 261.

Aleurolobus marlatti, new parasite of.

in Japan, 296.

Aleuroplatus kewensis, sp. n., on greenhouse ferns in Britain, 772.

Alfalfa (see Lucerne).

Alfalfa Snout Beetle (see Otiorrhynchus ligustici).

Alfalfa Weevil (see Hypera varia-

Algeria, Lindorus lophanthae predacious on Coccids in, 410; locusts in, 180, 239, 685; Tribolium castaneum in stored grain in, 413; vine and melon pests in, 371; summary of plant quarantine restrictions in, 134.

algirica, Cassida.

alienus, Lasius niger.

Alissonotum impressicolle, bionomics and control of, on sugar-cane in Burma, 294; Formosa, 294. in Assam and

Alkylphenylbenzenesulphonic Acid, reducing toxicity of derris, 553.

alliorum, Taeniothrips.

Allium (see Onion and Rocamboles). Allodontermes morogorensis, measures against, in Tanganyika, 113.

Allograpta exotica (predacious on Toxoptera graminum), parasite of, in Argentina, 289.

Allograpta obliqua, predacious on Aphis gossypii in Bermuda, 626.

Allotropa, parasite of mealybugs in Mexico, 586; failure to establish, in Hawaii, 586.

allyni, Eupelmus.

almanzori, Telenomus.

Almond, pests of, in N. Africa, 260, 340; Aonidiella aurantii on, in S. Africa, 455; Recurvaria nanella on, in Cyprus, 108; Ceresa bubalus on, in France, 343; pests of, in New Zealand, 266; Diloba coeruleocephala on, Palestine, 514; pests of, U.S.A., 18, 20, 722.

Almonds (Stored), pests of, 461,

573.

alope, Erinnyis. Alorco, a form of cryolite (q.v.), 16, 65, 736.

Alpha - naphthylamine, treated with, against Cydia pomonella, 507.

Alphitobius, measures against, in stored ragi in Mysore, 204.

Alphitobius laevigatus (piceus), in stored cassava tubers in Madagascar, 455.

alpium, Moma (Diphtera).

Alsike Clover (see Trifolium hybridum)

Alsophila pometaria, measures against, on shade trees in New Brunswick and U.S.A., 539, 547. alternans, Cleonus (Mecaspis); Pim-

alternata, Mycodiplosis.

alternatus, Nabis.

alternus, Stauropus.

Althaea rosea, Platyedra gossypiella on, in Brazil, 690; pests of, in British Isles, 33.

althaeae, Tetranychus (see T. tela-

Aluminium Stearate, in banding mixtures against Cydia pomonella, 507.

Aluminium Sulphate, as an adhesive for sprays, 443, 552.

Alysia manducator, factors cerned in host-finding by, 150.

amandus, Plochionus. amanii, Coptotermes.

Amara, attractiveness of cereals to, in Crimea, 356.

Amarantus, Loxostege bifidalis on, in Brazil, 333.

Amarantus retroflexus. Pyrausta nubilalis on, in Poland, 622.

Amaryllis, Brithys crini on, in Formosa, 52.

Amauromorpha metathoracica, parasite of Schoenobius bipunctifer in China, **109**.

ambigua, Myrmelachista.

ambiguella, Clysia. ambiguus, Psallus.

Amblymerus, parasite of Mayetiola

destructor in Morocco, 340.

Amblypelta lutescens, on papaya in Queensland, 246.

Amelanchier, Eriosoma americanum on, in U.S.A., 484.

America, North, Jassids of, 367. America, South, classification of rotenone-yielding plants in, 318, 319.

Malacosoma; Larra; americana, Meromyza; Periplaneta; Schistocerca.

americanum, Eriosoma.

americanus, Hister; Stigmus; Apan-

Ametastegia glabrata, on apple in Britain, 297.

amicroploides, Macrocentrus.

Amicroplus collaris, parasite Euxoa segetum in Italy, 528.

Amitermes, damaging cotton in Tanganyika, 113.

Amitermes (Globitermes) annamensis, fumigation tests against, Indo-China, 188.

amitinus, Ips.

Ammonia, in baits for Trypetids, 58, 697, 700; tests of, against cheese mites, 539.

Ammonium Carbonate, tests of, against wireworms, 631.

Ammonium Caseinate, uses of. in sprays, 174, 488, 552, 553, 739: reducing toxicity derris, 553.

Ammonium Fluoride, in baits and bait-spray for Trypetids,

Ammonium Oleate, uses of, in inverted spray mixtures, 174.

Ammonium Phosphate, increasing solubility of lead arsenate, 324.

Ammonium Sulphate, in bait and bait-spray for Trypetids, 108. Ammonium Thiocyanate, 703.

Amorphoidea pectoralis, sp. n., on cotton etc. in Java, 106, 766; bionomics and control of, 766.

ampelophaga, Haltica.

Amphimallon (see Amphimallus). Amphimallus majalis, bionomics of, damaging lawns in France, 229.

Amphimallus solstitialis, on mangel in Britain, 296; damaging lawns in France, 229; food-plants of, in Germany, 717; food-plants of, and soil conditions in relation to, in Russia, 72, 158, 356, 429, 430, 431, 432; measures against, 431, 432.

Amphimermis zuimushi, parasite of Chilo simplex in Japan, 671.

Amyl Acetate, in baits: for Euxoa, 236; for Orthoptera, 28, 139, 574. Amyl Alcohol, in bait for Dacus, 672.

Amyna octo, on cowpeas in Nyasaland, 114.

Anabadust, 236.

Anabasine, dusting tests with, against Aphids, 355.

Anabasine Sulphate, dusting with, 236; spraying with, 237, 355, 479, 739; and arsenicals, 479; and oil emulsions, 739;

soap, 237, 238, 355; other activators for, 237, 238; formulae containing, 236, 355; fixed compounds of, 479.

Anabrus simplex, in U.S.A., 44, 128, 580, 609, 686; loss due to, 44; poultry destroying, 128.

Anachaetopsis, parasite of Platyedra

gossypiella in Mexico, 275. Anachaetopsis tortricis, bionomics of,

in U.S.A., 286, 563. Anacridium aegyptium, bionomics

and control of, in Egypt, 685. Anacridium moestum, in Nyasaland, 115.

anactus, Papilio.

anaglypticus, Conotrachelus.

Anagrus, parasite of Aspidiotus perniciosus in Russian Union, 305.

Anagrus armatus var. nigriventris (parasite of Typhlocyba pomaria), bionomics of, in New York, 557, 558, 559; in New Zealand, 57, 208; introduced into Tasmania, 208; relation of insecticides to, 558, 559.

Anagyrus bohemani, parasite of Pseudococcus citri in Spain, 701.

Anagyrus coccidivorus, imported into Porto Rico against Pseudodococcus brevipes, 329.

Anaitis plagiata, introduction of, into Australia against Hypericum perforatum, 450.

analcidis, Microbracon.

Analeptes reticulata, on cotton in Tanganyika, 642.

Analeurops cuthbertsoni, gen. et sp. n., food-plants of, in S. Rhodesia, 106.

analis, Bruchus.

Anaphes anomocerus, parasite of Psallus seriatus in U.S.A., 328.

Anaphoidea nitens (parasite of Gonipterus scutellatus), establishment of, in New Zealand, 753; in Nyasaland, 204.

Anaphothrips orchidii, on Cyclamen europaeum in Japan, 673.

Anarsia lineatella (on stone fruits), in California, 19, 748; bionomics of, in Germany, 718; measures against, 718, 748.

anartoides, Orgyia.

Anasa armigera, bionomics of, in Connecticut, 17; characters of,

Anasa repetita, on cucurbits in Connecticut, 17.

Anasa scorbutica, on cucurbits in Jamaica, 490.

Anasa tristis (on cucurbits) in. U.S.A., 17, 269, 644, 725, 726, 739; bionomics of, 17, 726; 739; bionomics of, 17, measures against, 644, 725, 726, 739; characters of, 17.

Anastatus, parasite of Holochlora nawae in China, 244.

Anastatus bifasciatus, parasite of Lymantria dispar in Jugoslavia, 760.

Anastoechus nitidulus, egg-parasite of Dociostaurus maroccanus in

Italy, 163.

Anastrepha, not experimentally parasitised by Tetrastichus giffardi in Brazil, 628.

Anastrepha acidusa, treatment of fruit against, in Porto Rico, 325. Anastrepha fraterculus, on Citrus in Peru, 81.

Anastrepha longimacula, on mango

in Jamaica, 490.

Anastrepha ludens, in Mexico, 142, 325, 505; in Texas, 325; on mango, 142; legislation respecting, in U.S.A., 751; bionomics of, 325; measures against, 142, 325, 505, 751.

Anastrepha striata, extracts Haplophyton in bait-sprays for,

in Mexico, 505.

Anastrepha suspensa, treatment of fruit against, in Porto Rico, 325. Anatis ocellata, predacious

Aphids in Germany, 197.

ancylae, Sympiesis.

Ancylis comptana (Strawberry Leaf-Roller), in U.S.A., 45, 399, 735; bionomics of, 735; parasites and biological control of, 399, 735: other measures against, 45, 399, 735, 736.

Ancylis selenara, bionomics of, on apple and pear in Caucasus, 477. ancylivorus, Macrocentrus.

ancylus, Aspidiotus.

Andropogon, relation of Schistocerca paranensis to, in Argentina, 147.

Andropogon furcatus, Blissus spp.

on, in Iowa, 225.

Anethol, in baits for Cydia, 135.

Anethum graveolens, Listronotus latiusculus on, in New Jersey, 565. Angitia, parasite of Gracilaria theivora in India, 149; associated

with cotton pests in Mexico, 275. Angitia armillata, parasite of Cydia molesta in Italy, 154; parasite of Hyponomeuta in Ukraine, 302.

Angitia fenestralis (parasite Plutella maculipennis), attempted introduction of, into Australia, 449; introduced into Zealand, 449.

Angitia hellulae (see A. polynesialis). Angitia molestae, liberation of, against Cydia molesta in Connecticut, 16, 724.

Angitia paenerivalis, parasite of Sparganothis sulfureana in New Jersey, 563.

Angitia polynesialis, bionomics of,

in Virginia, 13.

Angitia punctoria (parasite Pyrausta nubilalis), in Poland, 623; establishment of, in U.S.A.,

Angitia rufipes, parasite of Platyedra malvella in Britain, 33.

Angola, locusts in, 684, 687, 688. Angoumois Grain Moth (see Sitotroga cerealella).

(Fergusobia, Anguillulina

gen. n.), association of, with Fergusonina in Australia, 86. angulata, Perisierola.

angusticollis, Zootermopsis. angustivostris, Eremnus.

Anilastus, parasite of Lixus junci in Morocco, 417.

Animula herrichi, on Eucalyptus in New South Wales, 334.

Aning, 236.

Anise Oil, in baits for Cydia, 506. Anisic Aldehyde, in baits for Cydia, 135.

Anisomeles ovata, Bemisia gossy-

piperda on, in India, 242.

Anisoplia austriaca, food-plants of, in Russia, 72, 356, 429, 430, 617; bionomics of, 617; soil conditions in relation to, 72, 356, 429, 430: measures against, 618.

Anisoplia segetum, on cereals in Germany, 717; on rubber-producing plants in Ukraine, 72.

annamensis, Amitermes (Globitermes). annonae, Ceratitis.

annulata, Campsomeris (Elis). annulatiorus, Microcryptus (Pezoporus).

annulicornis, Heterospilus; Hieroglyphus.

annulipes, Chelonus.

Anobium punctatum, damaging building timber in Connecticut, 17, 726.

Anomala chamaeleon, bionomics of, in Korea, 767.

Anomala cuprea, food-plants of adults of, in Japan, 54.

Anomala dubia aenea, damaging pine in Ukraine, 158.

Anomala errans, on cranberry in Massachusetts, 645.

Anomala horticola, damaging turf in Britain, 462; on apple in Finland, 201; on cereals in Germany, 717; not attacking bracken roots, 462.

Anomala orientalis, biological control of, on sugar-cane in Hawaii,

493.

Anomala pallidipennis, bionomics of, in Korea, 436, 767; measures against, 436.

Anomala rufocuprea, feeding habits of adults of, in Japan, 54, 516.

Anomala testaceipes (see Mimela). Anomala vitis, on vines in Czechoslovakia, 657.

Anomala xanthopleura, bionomics of, in Loochoo Islands, 516.

Anomis flava (see Cosmophila).

Anomis luridula (texana), bionomics and control of, on cotton in Peru, 533, 626, 627.

anomocerus, Anaphes.

Anona, weevils on, in Brazil, 115; Dacus cucurbitae on, in Philippines, 181.

Anona muricata, new Coccid on, in

Java, 607.

Anona reticulata, tests of insecticidal

properties of, 361.

Anona squamosa, Heterographis bengalella on, in Indo-China, 185. Anoplocnemis curvipes, on cotton

in Belgian Congo, 315.

Anoplolepis longipes, in Java, 95, 606; new mite in nests of, 95; associated with Pseudococcus citri, 606.

Anoxia pilosa, in forests in Ukraine, 158, 429; soil conditions in relation to, 429.

Antarchaea chionosticta, on cotton in Queensland, 494.

antennatus, Dasynus (Pendulinus).
Antestia, bionomics, synonymy and distribution of species of, on coftee in E. Africa, 177–180; associated with Nematospora, 178; measures against, 179.

Antestia faceta, 178. Antestia falsa, 178.

Antestra lineaticollis, on coffee in E. Africa, 178; parasites of, 179. Antestia trivialis, 178.

Antestia usambarica, 178.

antestiae, Corioxenos.

Antheraea pernyi, development of, in Japan, 247.

Antheraea yamamai, parasite of, in Japan, 310.

Anthidium maculosum, constructing nests in figs in California, 292, 293.

Anthocoris nemorum, predacious on mites in Holland, 465; killed by tar distillate sprays, 465.

anthonomi, Microdontomerus.

Anthonomus bisignifer (Strawberry Weevil), temperature limit of activity of, in Japan, 603.

Anthonomus campinas, sp. n., on

cotton in Brazil, 405.

Anthonomus cinctus (see A. piri).

Anthonomus eugenii, measures against, on Capsicum, etc., in U.S.A., 64, 328; bionomics of, 64.

Anthonomus grandis (Cotton Boll Weevil), parasites of, in Mexico, 275; in U.S.A., 142, 274, 276, 278, 279, 328, 329, 644, 691; cold-hardiness of, 274; measures against, 328, 644, 691; technique and results of insecticide tests against, 142, 276, 279.

Anthonomus grandis thurberiae (Thurberia Weevil), in U.S.A., 134, 329, 384; value of destroying Thurberia against, 329; legislation dealing with cotton-

seed against, 134, 384.

Anthonomus musculus (Cranberry Weevil), measures against, in Massachusetts, 645.

Anthonomus piri (Pear Blossom Weevil), bionomics and control of, on pear in Germany, 761.

Anthonomus pomorum (Apple Blossom Weevil), in Austria, 234; in Britain, 613, 689; in Cyprus, 260; in Czechoslovakia, 659; in Germany, 421, 762; in Italy, 351; on pear, 762; measures against, 234, 351, 421, 613, 659, 689; betanaphthol repellent to, 613.

Anthonomus rubi, measures against, on strawberry in Britain and

Estonia, 383, 462.

Anthonomus signatus, on strawberry in Delaware, 443, 545; measures against, 443.

Anthonomus spilotus, on pear in

Germany, 762.

Anthophila pariana, on apple in Britain, 527, 612; taken on high buildings in U.S.A., 215.

Anthores leuconotus, bionomics of, on coffee in N. Rhodesia, 500.

Anthracene, and chloropicrin, soil fumigation with, against beet pests, 236.

Anthracene Oil Emulsions, 301, 482. (See Tar Distillates.)

Anthrax hottentottus (see Villa).

Anthrenus scrophulariae, measures against, in clothing in U.S.A., 400.

Anthrenus verbasci, measures against, in clothing in U.S.A., 400.

Anthrenus vorax, in clothing and upholstered furniture in U.S.A., 227, 400; measures against, 400; used for tests of mothproofing solutions, 227.

Anticarsia gemmatalis, tests of insecticides on, 388.

Anticarsia irrorata, on cowpeas in Nyasaland, 114.

Antigastra catalaunalis, on Sesamum orientale in Cyprus, 108.

Antigua, Platyedra gossypiella in, 455; summary of plant pest legislation in, 491.

antillicola, Leptomastidea.

antilope, Xylotrechus.

Antimony, determination of, tartar emetic spray residues, 695. Antimony Potassium Tartrate (see

Tartar Emetic).

antiqua, Hylemyia; Orgyia (Bombyx, Notolophus); Promecotheca (see P. papuana).

Antirrhinum, Heliothis phloxiphaga on, in Vancouver, 293.

antonii, Helopeltis.

Antonina indica, on Cynodon and sugar-cane in Hawaii, 585.

Antrocephalus renalis, parasite of Tirathaba trichogramma in Fiji,

Ants, key to, in California, 129; damaging telephone poles and electric cables, 218, 262; associated with Aphids, 106, 107, 195, 196, 289, 588, 626; associated with Coccids, 202, 204, 411, 435, 588, 592, 606, 626, 642, 758; new mite associated with, 95; Labidostomis taxicornis in nests of, 70; destroying other insects, 22, 61, 208, 235, 289, 305, 339, 403, 435, 504; natural enemies of, 95, 292, 397; general accounts of, 120, 129, 228; measures against, 11, 93, 129, 202, 218, 222, 228, 286, 533, 588, 592; differences in sensitivity of species of, to arsenical baits, 716.

Ants, California Fire (see Solenopsis geminata var. maniosa).

Ants, Carpenter (see Camponotus herculeanus pennsylvanicus)

Ants, Leaf-cutting (see Atta and A cromyrmex).

Anuvaphis maidiradicis, measures against, on maize and cotton in U.S.A., 125, 645.

Anuraphis padi, on peach in India,

Anuraphis persicae (see A. schwartzi) Anuraphis roseus (on apple), in Britain, 527; measures against, in U.S.A., 16, 738.

Anuraphis schwartzi, experimentally parasitised by Aphidius platensis in Argentina, on almond in Libya, 260.

Anuraphis tulipae, on tulips in Britain, 350; measures against, on Gladiolus in Iowa, 65.

Anychus orientalis, ecology of, on Citrus in Palestine, 405.

Aonidiella, status and scope of, 226, 469.

Aonidiella aurantii (Citrus Red Scale), in S. Africa, 202, 377, 455; in Brazil, 502; in Cyprus, 706; in Indo-China, 185; in Japan, 604; in Transcaucasia, 481; in U.S.A., 20, 117, 324, 574, 575; in Victoria, 85; review of distribution of, 455; miscellaneous food-plants of, 455, 502, 604; bionomics of, 85, 324, 455, 456; measures against, 85, 117, 185, 202, 456, 481, 574, 575, 706, 707; susceptibility of strains of, to HCN, 324; fumigation tests with methyl iodide on, 575. Aonidiella eremocitri, sp. n., on

Eremocitrus glauca in Queensland,

Aonidiella orientalis, on Solanum arundo in Tanganyika, 757.

Aonidiella perniciosa (see Aspidiotus).

aonidum, auct., Chrysomphalus (see C. ficus).

Apanteles, parasite of Argyroploce illepida in Fiji, 497; parasite of Lymantria dispar in Germany and Jugoslavia, 234, 760; parasite of Lasiocampid in Korea, 264; parasite of Artona catoxantha in Philippines, 675; key to species of, 106.

Apanteles americanus, parasite of Sphingids in West Indies, 171, **490**; parasites of, **171**.

Apanteles circumscriptus (bicolor, Nees), characters, synonymy and host records of, 342.

Apanteles congregatus, parasite of Protoparce in Jamaica, 490.

Apanteles flaviventris, A. americanus recorded as, 171.

Apanteles glomeratus, parasite of Pieris rapae in Japan, 605; parasite of Naranga aenescens in Korea, 768; giant cells in larvae of P. brassicae parasitised by, 464.

Apanteles kariyai, sp. n., parasite of Cirphis unipuncta in Man-

churia, 296.

Apanteles militaris (parasite of armyworms), in Connecticut, 723; in Mexico, 586; introduced into Hawaii, 586.

Apanteles myeloenta, sp. n., parasite of Myelois ceratoniae in

Cyprus, 106.

Apanteles ruficrus, parasite of Spodoptera mauritia in India, 241; parasite of Cirphis unipuncta in Manchuria, 296.

Apanteles solitarius, establishment of, in U.S.A. against Stilpnotia

salicis, 590.

Apanteles sorghiellae, parasite of Celama sorghiella in Texas, 280.

Apanteles taragamae, parasite of Nephantis serinopa in Cochin, 452.

Apanteles tasmanica, 10.

Apanteles tirathabae, parasite of Tirathaba trichogramma in Fiji, 334.

apantelivorus, Horismenus.

Apate monacha, food-plants of, in Sierra Leone, 240.

Apate terebrans, food-plants of, in Jamaica, 490, 491.

aperta, Mylabris.

Aphalara calthae, on narcissus in U.S.A., 288.

Aphanogmus steinitzi, parasite of Coniopterygid in Palestine, 637. Aphelinus jucundus, parasite of

Aphids in Ontario, 537.

Aphelinus mali, utilisation of, against Eriosoma lanigerum, 41, 65, 74, 108, 192, 233, 347, 469, 485, 696, 771; bionomics of, 771.

Aphelinus marlatti, bionomics of, parasitising Aphids in Ontario,

537, 538.

Aphelinus mytilaspidis (see Aphytis). Aphelopus typhlocybae (parasite of Typhlocyba pomaria), bionomics of, in U.S.A., 557, 558, 559; attempted introduction of, into New Zealand against T. froggatti,

57; effect of sprays on, **558**, **559**.

Aphidecta obliterata, predacious on Chermes nordmannianae in Germany, 661.

Aphidencyrtus aphidivorus, parasite of Carolinaia setariae in Porto

Rico, **39.** aphidis, Pachyneuron.

Aphidius, parasite of Aphids in Ontario, 538.

Aphidius brassicae, parasite of Brevicoryne brassicae in Ukraine, 303

Aphidius phorodontis, parasite of Myzus persicae in Ontario, 537.

Aphidius platensis (parasite of Toxoptera graminum), new Cynipid parasite of, in Argentina and Uruguay, 31, 289, 290; bionomics of, 289, 290.

Aphidius ribis, giant cells in Aphid parasitised by, 464.

aphidivorus Aphidencyrtus.

Aphids, and plant diseases, 12, 32, 39, 40, 47, 68, 104, 239, 265, 377, 379, 383, 411, 484, 485, 495, 527, 531, 564, 577, 592, 615, 626, 759; ants associated with, 106, 107, 195, 196, 289, 588, 626; natural enemies of, 39, 41, 65, 74, 98, 108, 128, 189, 192, 195, 196, 197, 214, 233, 238, 289, 290, 296, 302, 303, 309, 329, 335, 347, 353, 354, 370, 459, 469, 469, 469, 469, 502 379, 450, 464, 469, 485, 523, 531, 537, 538, 611, 626, 661, 673, 696, 727, 763, 768, 771; technique for rearing Coccinellids on frozen, 649, 650; factors affecting wing development in, 104, 107, 632; methods of penetration of setae of, 225; tests of insecticides on, 17, 20, 21, 34, 35, 36, 37, 66, 616, 631, 632, 704; tests of contact agents on, 45; classification and new species of, 106, 119, 247, 265, 304, 367, 369, 440, 452, 496, 624, 696, 772; records of, in China, 367; key to, producing honey-dew in Central Europe, 459; of Formosa 119; Italy, 696; of Japan, 496; of Manchuria, 496; of Morocco, 48, 696; of Poona, 247.

Aphis chloris, imported into Australia against Hypericum perforatum, 450.

Aphis fabae*, ants associated with, in Germany, 195.

^{*} In view of the divergent opinions on the identity of the Aphids of this group, no attempt is here made to show the synonymy.

Aphis gossypii, in Argentina, 290; in Bermuda, 626; in China, 245, 304; in Belgian Congo, 315; in Formosa, 309; in Hawaii, 214; in Japan and Korea, 309; in Russian Union, 249, 353, 354, 472; in Sudan, 513; in U.S.A., 269, 399; on cotton, 245, 304, 309, 315, 353, 354, 472, 513; susceptibility of varieties of cotton to, 309; on cucurbits, 249, 269, 309, 399; transmitting rosette disease of lilies, 626; other foodplants of, 214, 245, 309; bionomics of, 245, 309; natural enemies of, 309, 353, 354, 626; experimentally parasitised by Aphidius platensis, 290; measures against, 245, 304, 309, 399, 472; effect of essential oils on, 249, 250.

Aphis laburni (leguminosae), and virus diseases of ground-nuts in Africa, 114, 239, 531, 592; bionomics of, on Astragalus and Phaseolus in Japan, 53; on cotton in Ukraine, 353; Coccinellids predacious on, 353, 531;

measures against, 531.

Aphis maidis, recorded as Rhopalosiphum prunifolii on cereals in Argentina, 31, 289; in Natal, 411; not transmitting virus diseases of Sorghum in Madras, 532; food-plants and natural enemies of, in Porto Rico, 39, 40; in U.S.A., 577; relation of, to sugar-cane mosaic, 40, 411, 577.

Aphis papaveris*, on beet in Czechoslovakia, 656.

Aphis plantaginis, on rubberproducing plants in Ukraine, 106.

Aphis pomi (on apple), in Britain, 297, 527; ants associated with, in Germany, 195; measures against, in New Jersey, 738; tests of ovicides against, 297.

Aphis rhamni, on potato in Britain, 527.

Aphis roseus (see Anuraphis).

Aphis rumicis*, in Britain, 296, 527; in Libya, 260; in U.S.A., 12, 19, 21, 445, 649; on beet and mangel, 19, 296, 445, 527; on leguminous plants, 19, 260, 527; on nasturtium, 21; on potato, 527; not a vector of certain virus diseases, 12, 445;

penetration of setae of, 225; technique for rearing Coccinellids on frozen, 649; tests of insecticides on, 17, 21, 616, 631, 632, 704; tests of contact agents on, 45.

Aphis sacchari, not transmitting virus diseases of Sorghum in Madras, 532; on sugar-cane in

Natal, 411.

Aphis sambuci, ants associated with, in Germany, 195.

Aphis sorghi, on Sorghum in Sudan, 513.

Aphis spiraecola, Coccinellid predacious on, on Citrus in Florida, 727.

Aphis tavaresi, on Citrus in Mysore, 361; tests of insecticides on, 34, 35, 36, 37, 361.

Aphis tulipae (see Anuraphis).

Aphis, Cabbage (see Brevicoryne brassicae).

Aphis, Corn Root (see Anuraphis maidiradicis).

Aphis, Cotton (see *Aphis gossypii*). Aphis, Hop (see *Phorodon humuli*). Aphis, Large Willow (see *Lachnus*

salignus).

Aphis, Mealy Plum (see Hyalopterus arundinis).

Aphis, Ornate (see Myzus ornatus).
Aphis, Pea (see Macrosiphum onobrychis).

Aphis, Peach (see *Myzus persicae*). Aphis, Spruce Gall (see *Chermes abietis*).

Aphis, Strawberry (see Capito-phorus fragariae).

Aphis, Sumach Gall (see Melaphis chinensis).

Aphis, Woolly Apple (see *Eriosoma lanigerum*).

Aphodius howitti, bionomics and control of, in New South Wales, 209.

Aphodius tasmaniae, damaging pastures in Victoria, 85.

Aphomia gularis (in dried fruit, etc.), in California, 573; cold storage against, in Holland, 599; characters of eggs of, 196.

Aphrophora quadrinotata, on narcissus in U.S.A., 288.

Aphthona bimaculata, on Sesamum in Nyasaland, 114.

Aphthona euphorbiae, on flax in Poland, 255.

^{*} In view of the divergent opinions on the identity of the Aphids of this group, no attempt is here made to show the synonymy.

Aphycus lounsburyi, imported into Peru against Saissetia oleae, 81.

Aphytis chrysomphali (parasite of Coccids), in S. Africa, 456; in Italian Somaliland, 641; introduced into Seychelles, 43.

Aphytis mytilaspidis (parasite of Aspidiotus perniciosus), in Idaho, 65, 747; in Transcaucasia, 305.

apicalis, Nephotettix (see N. bipunctatus).

apicicornis, Coryna.

Apina callisto, damaging grasses in New South Wales, 111.

Apiomyia bergenstammi, on pear in Czechoslovakia, 657.

Apion, measures against, on clover in Britain, 527.

Apion aestivum, on clover in Russia, 307, 474; bionomics and control of, 307, 308.

Apion apricans, bionomics and control of, on clover in Russia, 75, 307, 308, 474, 475.

Apion flavipes, bionomics of, on clover in Russia, 474.

Apion radiolus, on hollyhock in Britain, 33.

Apion seniculum, bionomics of, on alsike clover in Latvia, 256.

Apion xanthostylum, new parasite of, on cotton in Tanganyika, 610. apionidis, Entedon.

Apoderus tranquebaricus, on Schleichera trijuga in Ceylon, **359.**

Aporia crataegi (on fruit trees), in Czechoslovakia, 657; in Italy, 712, 713; measures against, 713. Aporia crataegi adherbal, on apple

in Japan, 673.

Apotetranychus (see Tetranychus).

appendigaster, Eurytoma.

Apple, Cydia pomonella on, Afghanistan, 340; Aonidiella aurantii on, in S. Africa, 455; pests of, in Australia, 8, 10, 149, 168, 208, 335, 449, 494, 510; Anthonomus pomorum on, in Austria, 234; new weevil on, in Brazil, 612; pests of, in Britain, 297, 527, 612, 613; pests of, in Canada, 59, 171, 484, 485, 534, 535, 541; Coccids on, in Chile, 221; pests of, in Cyprus, 107, 108, 259, 260; pests of, in Czechoslovakia, 620, 656, 657, 658, 659; Anomala horticola on, in Finland, 201; pests of, in France, 192, 259, 343, 416, 525; pests of, in Germany, 2, 52, 101, 233, 234, 250, 347, 421, 660, 719;

Paratetranychus pilosus on, in Holland, 464, 465; C. pomonella on, in Hungary, 254; pests of, in India, 340, 361, 409, 765; pests of, in Italy, 351, 593, 713; pests of, in Japan, 208, 604, 671, 673, 771; Lymantria dispar on, in Jugoslavia, 759; pests of, in Korea, 243, 671; Eriosoma lanigerum on, in Libya, 260; Eriosoma pests of, in Malta, 469; pests of, in Morocco, 340, 341; pests of, in New Zealand, 46, 56, 58, 208, 247, 266; C. pomonella on, in Palestine, 500; pests of, in Poland, 98, 99; E. lanigerum on, in Rumania, 192; pests of, in Russian Union, 74, 75, 159, 302, 305, 306, 307, 353, 356, 477, 478, 479, 480, 481, 482, 530; Aglaope infausta on, in Spain, 697; pests of, in Switzerland, 299; pests of, in U.S.A., 15, 16, 25, 41, 44, 61, 62, 65, 79, 125, 126, 136, 174, 226, 269, 271, 272, 287, 288, 291, 320, 332, 333, 399, 401, 442, 444, 484, 488, 507, 544, 545, 550, 551, 554, 555, 556, 557-559, 561, 562, 574, 587, 588, 646, 647, 652, 653, 723, 724, 725, 731, 732, 733, 734, 737, 738, 745, 747; varietal susceptibility of, to pests, 2, 250, 299; Dacus cucurbitae 271. experimentally developing in, 668; insecticides causing injury to, 79, 97, 98, 150, 216, 254, 300, 351, 355, 442, 466, 480, 481, 482, 510, 544, 556, 588, 614, 647, 653, 658, 708, 738; method for measuring effects of dormant sprays on growth of, 737; wetting properties of Ultrawet on, 285.

Apple (Fruit), pests in stored, and their control, 269, 306, 307, 541, 612; problems of spray residues on, 25, 79, 175, 259, 318, 320, 401, 402, 442, 449, 488, 510, 525, 555, 556, 614, 647, 652, 653, 732, 733, 750; determination of spray deposits on, 320, 368; uses of, in baits, 45, 168, 172, 566; (dried), not attacked by Aphomia gularis in California, 573.

Apple Aphis, Woolly (see *Eriosoma lanigerum*).

Apple Blossom Weevil (see Anthonomus pomorum).

Apple Bud Moth, Eye-spotted (see Eucosma ocellana).

Apple Leaf-curling Midge (see Dasyneura mali).

Apple Leaf-hoppers (see *Typhlocyba froggatti* and *T. pomaria*).

Apple Maggot (see Rhagoletis pomonella).

Apple Moth (see Hyponomeuta padellus malinellus).

Apple Moth, Light Brown (see Tortrix postvittana).

Apple Root Borer (see Dorysthenes huegeli).

Apple Sucker (see *Psylla mali*). Apple Worm, Lesser (see *Enar*monia prunivora).

Apple-tree Borer, Flat-headed (see Chrysobothris femorata).

Apple-tree Borer, Round-headed (see Saperda candida).

approximata, Aristobia.

apricans, Apion.

Apricot, pests of, in Australia, 8, 111, 335; Macropsis trimaculata on, in Canada, 173; Aspidiotus perniciosus on, in N. Caucasus, 481; Anarsia lineatella on, in Germany, 718; pests of, in India, 242, 765; Lymantria dispar on, in Jugoslavia, 759; Monosteira unicostata on, in Morocco, 340; Tortricids on, in New Zealand, 57, 58; Ceratitis capitata on, in Switzerland, 715; pests of, in U.S.A., 18, 19, 127, 173.

Apricot (Dried Fruit), pests of, in California, 30, 228; not attacked by Aphomia gularis, 573; (kernels) cold storage of, against

pests in Holland, 599.

Apricot Scale, Brown (see Lecanium

armeniacum).

Aprostocetus aspidomorphae, sp. n., doubtful identity of host of, in Uganda, 610.

apterus, Chasmodon; Lethrus.

Arabia, locusts in, 112, 198, 683,

Arachis hypogaea (see Ground-nuts). Araecerus (Araeocerus) fasciculatus, bionomics and control of, in Chinese yeast in Korea, 768; in stored cassava tubers in Madagascar, 455.

Aralia (see Fatsia).

Araucaria cunninghami, Aesiotes notabilis on, in Queensland, 494. Araucaria excelsa, Coccid on, in

New Zealand, 753. Arbela (see Indarbela). Arbor-vitae (see Thuja). arbustorum, Plagiognathus. archippus, Basilarchia. Archips (see Tortrix).

Archytas cirphis (parasite of armyworms), introduced into Hawaii from Mexico, 586.

Arctium lappa, Nymphalid on, in Japan, 673.

arcuatus, Chortoglyphus; Cirrospilus (Entedon) pictus.

areator, Hemiteles.

Areca catechu, new Capsid on, in Malaya, 198.

arecae, Parasthenaridea. arenosella, Batrachedra.

Areskap, as a wetting agent for derris, 645.

argaula, Agonoxena.

Argentina, Aphids on cereals in, 30, 289, 290; cotton pests in 362, 503, 704; miscellaneous in, 625, 689, 772; mite on orange in, 581; pests of solanaceous plants in, 290, 581; list of Heteroptera of, 696; locusts and their natural enemies in, 146, 147, 655, 689; vectors of plant diseases in, 581; beneficial insects in, 31, 146, 147, 289, 290, 689, 772; parasites of Diatraea introduced into U.S.A. from, 390, 391; summaries of plant quarantines in, 134, 385, 692.

argillacea, Alabama.

argiolus, Lycaenopsis (Lycaena).

Argyresthia conjugella, bionomics and control of, in Switzerland, 299; distribution of, 299.

Argyresthia ephippella (pruniella, auct.), on cherry in Finland, 201. Argyria sticticraspis (see Diatraea). Argyroploce abietana, measures against, on spruce in New York, 270.

Argyroploce illepida, on Inocarpus edulis in Fiji, 497; parasites and distribution of, 497.

Argyroploce leucotreta, on cotton in E. Africa, 376, 642; boll rots associated with, 376.

Argyroploce variegana, on apple and pear in Caucasus, 477.

argyrospila, Tortrix (Archips, Cacoecia).

semipurpurana (Oak Argyrotoxa Leaf-roller), host of Glypta rufiscutellaris in Ontario, 509; measures against, in U.S.A., 547.

aridula, Chaetocnema.

aristella, Lonchaea.

Aristobia approximata, on Xylia dolabriformis in Burma, 339.

Aristolochia spp., tests of insecticidal properties of, 361.

Arizona, cotton pests in, 227, 328, 329, 580; Phthorimaea lycopersicella on tomato in, 580; Vanduzea segmentata in, 395; beneficial insects in, 227, 328.

Arkansas, Cydia pomonella in, 731; new parasite of Psallus seriatus in, 227.

armata, Cylindromyia; Hypoga-

strura.
armatus, Anagrus; Myochrous.

armeniacum, Lecanium.

armicollis, Magdalis.

armigera, Anasa; Heliothis (Chlor-idea); Hispa.

Armillaria matsutake, Scatopse fuscipes on, in Japan and Korea, 438.

armillata, Angitia. arquatus, Xyleborus.

Arrhenatherum, possibly little

attacked by Melolontha, 521. Arsenic, in baits, 28, 37, 45, 139, 182, 224, 398, 520, 609, 686, 698; types of, in baits for cutworms, in bait-spray for Lygus, 540: 376; dusts containing, 69, 383, 542; uses of, against ants and termites, 449, 533; differences in sensitivity of ants to, 716; problem of residues of, on sprayed fruit, 25, 79, 140, 169, 259, 318, 328, 383, 402, 490, 525, 556, 614, 646, 647, 750; danger to bees and live stock of insecticides containing, 69, 609, 748; ferric hydrate as treatment for bees poisoned by, 255; action of, on leaves, 600; determination of, 695, 772; amorphous precipitates of salts of, as emulsifiers for oils, 482; standardisation of insecticides containing, in Switzerland, 156.

Artemisia, relation of wireworms to, in Central Asia, 76; insects on, in U.S.A., 27, 496, 650, 696,

772.

Artemisia vulgaris, as a food-plant for Pyrausta nubilalis, 475, 476, 622, 623.

Arthemis vulgaris, Tetranychus on, in Ukraine, 257.

Artichoke, Cossus cossus on, in Sicily, 93; liable to attack by Baris interstitialis, 450.

articulatus, Selenaspidus (Pseudaonidia).

Artocarpus integer, Margaronia caesalis on, in Ceylon, 358.

Artona catoxantha (on coconut), in Java, 674; parasite of, in

Philippines, 675; measures against, 674, 675.

arundinis, Hyalopterus.

Asaphes vulgaris, hyperparasite of Brevicoryne brassicae in Ukraine, 303.

Ascia (see Pieris).

Ascogaster carpocapsae (parasite of Cydia pomonella), in Canada, 535; in U.S.A., 30, 65, 291, 555, 747; utilisation of, 65, 291, 535; simultaneous propagation of Macrocentrus and, on C. molesta, 535; parasite of, 30; spray treatments effecting, 555. asella. Heterogenea.

Aserica castanea, in U.S.A., 215, 324, 325; biological control of, 324, 325; taken on high build-

ings, 215.

Aserica japonica, food-plants of, in Japan, 605.

Aserica orientalis, bionomics of, in Korea, 436, 767; measures against, 436.

asericae, Tiphia.

Ash (Fraxinus), pests of in Germany, 251, 345; Anomala pallidipennis on, in Korea, 767; Hylesinus aculeatus on, in New Jersey, 394; pests of, in Russian Union, 75, 429, 430; (timber), pests in, and their control in Britain, 190, 483, 631, 707; not experimentally attacked by Codiosoma spadix in Russia, 354; pests intercepted in Vancouver in crates of, 483; starch content of, in relation to pests, 631, 707.

Ash Bark-beetle (see Hylesinus

fraxini).
ashmeadi, Telenomus.

Asia, Central, apple pests in, 479, 530; cotton pests in, 238, 249, 250, 470, 471, 472, 530; locusts in, 77, 426; wireworms on wheat in, 76; keys to Coccids of, 301; Tetranychid mites of, 470, 530; utilisation of Coccinellids against *Hypera* and Aphids in, 238.

Asiatic Garden Beetle (see Aserica castanea).

Asiatic Locust (see Locusta migratoria).

asparagi, Crioceris.

Asparagus, pests of, in Britain, 32, 33; Scutigerella on, in California, 280; Crioceris spp. on, in Japan, 497.

Asparagus falcatus, Brachytes bicolor on, in Ceylon, 359.

Asparagus plumosus, Coccid on in Germany, 101.

Asparagus Miner (see Agromyza simplex).

asperatus, Sciaphilus.

Aspergillus flavus, infesting locusts, 591.

Aspergillus versicolor, experiments with, infesting Lepidoptera, 420. Asphondylia, on egg-plant in Tanganyika, 642.

Asphondylia capparis, on caper in

Sicily, 92.

Asphondylia gennadii (see Eumar-

chalia).

Aspidiotiphagus citrinus (parasite of Coccids), in S. Africa, 378; utilisation of, in Netherlands Indies, 337; in Mauritius, 66.

Aspidiotiphagus pseudoaonidiae, sp. n., parasite of Pseudaonidia spp. in Japan, 296.

Aspidiotus, characters guishing Aonidiella and, 470.

Aspidiotus ancylus, food-plants of, in U.S.A., 20, 172; measures

against, 172.

Aspidiotus aurantii (see Aonidiella). Aspidiotus destructor, in Netherlands Indies, 337; in Mauritius, 176, 204; in Italian Somaliland, 641; in Tanganyika, 113; in West Indies, 10, 41, 43, 268; on coconut, 10, 43, 176, 204, 268, 337; on other plants, 41, 113, 641; natural enemies and biological control of, 10, 41, 43, 176, 204, 268, 269, 337, 641.

Aspidiotus forbesi (Cherry Scale), measures against, on apple in

Virginia, 272.

Aspid otus harti, measures against, on yams on Cayman Islands, 269.

Aspid otus hederae, on carob in Cyprus, 458; fungus infesting, on asparagus fern in Germany, 101; in Libya, 260; food-plants of, in Nyasaland, 114; on olive in Spain, 700.

fumigation Aspidiotus lataniae. against, on avocado in California, 116; food-plants of, in Nyasa-

land, 114.

Aspidiotus orientalis (see Aonidiella). Aspidiotus ostreaeformis, measures against, on fruit trees in Italy, 351.

Aspidiotus (Aonidiella) perniciosus, in Austria, 662, 663, 715; in Br. Columbia, 59; in India, 765; in Russian Union, 305-307, 480, 481, 483; in U.S.A., 19, 25, 65, 125, 172, 272, 307, 389, 545, 738, 747; on fruit trees, 25, 59, 65, 125, 272, 305, 306, 480, 481, 483, 545, 662, 663, 738, 747, 765; on other plants, 19, 172, 305; bionomics of, 125, 305, 306, 662, 663: natural enemies and biological control of, 65, 305, 715, 747; other measures against, 25, 172, 272, 306, 480, 481, 483, 545, 662, 663, 738; susceptibility of, to methyl bromide, 389; generic position of, 470.

Aspidiotus personatus (see Chrysom-

phalus).

Aspidiotus vapax, food-plants of, in Nyasaland, 114.

Aspidiotus sylvaticus, on tea in Nyasaland, 114.

aspidistrae, Pinnaspis.

Aspidomorpha tecta, new eggparasites of, in Uganda, 610.

aspidomorphae, Aprostocetus.

Asplenium, Coccids on, in Britain, 712.

Aspongopus janus, on cucurbits in Ceylon, 358.

Asprothrips raui, gen. et sp. n., in greenhouses in New York, 490.

Assam, miscellaneous pests in, 294. 409, 601; tea pests in, 601, 693, 694.

assectella, Acrolepia.

assimilis, Gryllulus (Gryllus).

assulta, Heliothis.

Aster, pests of, in Canada, 293, 538; Jassid transmitting yellows disease of, in U.S.A., 78, 386.

Asterolecanium conimbrigensis, n., on oaks in Portugal, 315.

Asterolecanium pustulans, restricted to Nerium in Formosa, 263.

Asterolecanium seabrai, sp. n., on Quercus faginea in Portugal, 315. astheniata, Urapteroides.

Astragalus sinicus, Aphid on, in Japan, 53.

Astycus lateralis, on Aleurites spp. in Burma, 340; on kapok in Indo-China, 186.

Astylus atromaculatus, bionomics and control of, in Argentina, 704, 705.

Asynonychus godmani, on potato in Hawaii, 214.

Atacta geminata, parasite of Protoparce in Jamaica, 490.

atalantae, Theronia.

Atanycolus ulmicola, bionomics of, in Connecticut, 14.

Ataxia hubbardi, introduced into Australia to destroy Xanthium pungens, 450; attacking economic plants, 450.

ater, Hylastes.

aterrima, Epicauta.

aterrimus, Orchidophilus; Scymnillus.

Athalia, on watercress in China, 497.

Athalia japonica, on crucifers in Japan, 605.

Athalia sjöstedti, on turnip in Tanganyika, 113.

Athous niger, soil conditions in relation to, in Russia, 425.

Athyrium, Heptamelus ochroleucus on, in Britain, 152.

atlanis, Sarcophaga. atlantis, Diplotaxis.

atlas, Attacus.

Atomaria linearis, on beet in Czechoslovakia, 657.

atomaria, Epicauta. atra, Phyllotreta. atriceps, Oncophanes. atricolor, Geocoris. atricornis, Phytomyza. atriloqua, Batrachedra. atripes, Eurytoma.

Atriplex, relation of Eutettix tenellus to species of, in California, 323, 445; effect on Euxoa segetum of

feeding on, 424.

atriplicella, Phthorimaea. atrocoxalis, Pimpla (Itoplectis).

atromaculatus, Astylus. atropunctata, Eupteryx.

Atta nigra (see Acromyrmex).

Atta sexdens, in Brazil, 248, 533; bionomics of, 248; tests of fumigants against, 533.

Atta striata (see Acromyrmex). attaci, Pimpla (Itoplectis); Telenomus.

Attacus atlas, new parasite of, in Malaya and Siam, 89.

Attagenus piceus, in U.S.A., 227, 400, 568; bionomics of, 568; measures against, 227, 400. attenuata, Psylliodes.

Aucuba japonica, Coccids on, 560. augur, Leptocoris (Serinetha).

Aulacaspis pentagona (White Peach Scale), Prospaltella berlesei imported into Porto Rico against, 329.

Aulacizes irrorata, on cotton in Texas, 278.

Aulacophora cincta (see Ceratia). Aulacophora foveicollis (see Rhaphidopalpa). Aulocara elliotti, measures against, damaging grasses in Montana, 44

aurantiae, Toxoptera (see T. aurantii).

aurantii, Aonidiella (Aspidiotus, Chrysomphalus); Toxoptera.

aurata, Cetonia.
auratus, Dirhinus; Rhynchites.

aurichalceus, Dermestes.

auricularia, Forficula.

aurifer, Lachnopus. auriflua, Scirpophaga

nivella). auritus, Conoderus (Monocrepidius). aurostriatus. Cratopus.

S.

(see

Austracris guttulosa, in Australia, 584.

Australia, association of Nematodes and Agromyzids on Eucalyptus in, 86; miscellaneous pests in, 103, 362, 448-450, 497; Phylloxera vitifoliae on vines in, 608; grasshoppers in, 449, 583, 584, 629, 676, 686; beneficial insects and biological control of insects and plants in, 86, 448, 449, 450, 582, 629; Ephestia cautella intercepted in cacao in, 435; beneficial insects introduced into other countries from, 450, 753; spread of Citrus pests into New Zealand from, 752, 753, 754. (See also under the various States.)

Australia, South, grasshoppers in, 111, 583, 584, 703; new thrips transmitting spotted wilt in, 318; regulations against introduction

of Phylloxera into, 608.

Australia, Western, Bruchus pisorum in, 754; grasshoppers in, 169, 583, 584, 629, 676, 686; noxious mites in, 150, 494, 582; beneficial Arthropods and biological control in, 84, 169, 212, 458, 629; regulations against introduction of Phylloxera into, 608.

australicum, Trichogramma.

australis, Eucelatoria; Helicobia; Monolepta; Oedaleus.

australis, Frogg., Typhlocyba (see

T. froggatti).

Austria, Aphids producing honeydew in, 460; Bruchus obtectus in the field in, 301; forest pests in, 215, 589, 659, 660, 663, 664, 715; orchard pests in, 234, 662, 715; fungus infesting Aspidiotus perniciosus in, 715; beneficial insects in, 215, 589, 659, 660, 715; birds destroying Tanymecus

palliatus in, 255; danger of introduction of Leptinotarsa decembineata into, 518; summary of plant quarantines in, 134. austriaca, Anisoplia; Eurygaster. austriacus, Hemiteles albipalpus. Austroicetes cruciata (jungi, pusilla, auct.), biology and outbreaks of, in Australia, 111, 169, 583, 584, 629, 676, 686, 703; natural enemies of, 169, 629; measures against, 169.

Autographa brassicae (see Plusia). Autoserica (see Aserica).

avellanae, Phylus coryli. avenae, Macrosiphum.

avenae, auct., Siphocoryne Rhopalosiphum prunifoliae).

Avocado, pests of, in Brazil, 502; Aspidiotus lataniae on, in California, 116; thrips on, Caucasus, 354; Xyleborus morstatti on, in Fiji, 497; Pseudococcus nipae on, in Mexico and Hawaii, 586; Oncideres spp. on, in Peru, 82; Apate monacha on, in Sierra Leone, 240; termite damaging, in Tanganyika, 113; book on pests of, 722.

Avocado Mealybug (see Pseudo-

coccus nipae).

Axionicus insignis, bionomics of, on Brachychiton in New South Wales, 404.

axiridis, Coccinella (Harmonia). Azalea, pests of, in U.S.A., 20, 286. 328, 748; insects associated with disease of, 328.

azaleae, Aleurodes; Eriococcus.

В.

baccarum, Dolycoris.

Bacillus spp., infecting Schistocerca

gregaria, 184, 591.

Bacteria (from fermenting syrup), tests of, in baits for Cydia pomonella, 137.

Bacteria, Beneficial, 123, 147, 152, 184, 235, 267, 591, 717; spraying with against vine moths, 619.

Bacteria, Injurious, 202, 588: relation of insects to, 376, 385.

Bagworm, Common (see Thyridopteryx ephemeraeformis).

Bagworm, Wattle (see Acanthopsyche junodi).

Bahamas, biological control of pests of Citrus in, 491.

Bait-sprays, for Antestia, 179; for Lygus simonyi, 376; tests of, for moths, **50**, **51**; for Trypetids, **61**, **71**, **108**, **267**, **325**, **407**, **505**,

526, 619, 700, 713.

Baits, for ants, 11, 202, 219, 222, 223, 228, 588, 592, 716; for Coleoptera, 28, 45, 75, 153, 172, 209, 211, 269, 270, 393, 566, 630, 643, 705, 757; for crickets, 40, 340, 457, 528; for Dysdercus, 186; for Lepismids, 37, 224, 387, for locusts and grass-520: hoppers, 23, 138, 139, 140, 170, 326, 398, 410, 540, 574, 590, 608, 611, 682, 684, 685, 686, 698, 699, 748; for moths, 50, 57, 58, 125, 135, 137, 154, 168, 236, 423, 442, 473, 488, 506, 544, 652, 723, 745; tests of dry, for moths, 50; for Lepidopterous larvae, 282, 441, 498, 509, 540, 639; for Phorid parasite of honey-bees, 115; for thrips, 333, 727; for Trypetids, 58, 108, 142, 182, 407, 526, 672, 697, 699, 700, 713; formulae for, 37, 45, 75, 108, 135, 138, 139, 153, 168, 170, 172, 182, 202, 209, 223, 224, 387, 393, 398, 410, 506, 526, 540, 566, 574, 609, 639, 672, 682, 685, 686, 697, 698, 699, 700, 705, 713; use of aeroplanes for distributing, 398; mechanical spreaders for applying, 590; types of containers for, 108, 115, 168, 223, 506, 526, 697, 699, 713. bajulus, Hylotrupes.

Balaninus (see Curculio).

Balanogastris kolae, bionomics and control of, in Sierra Leone, 239. Balioptera tripunctata (see Geomyza). balteatus, Syrphus.

Baluchistan, Cydia pomonella in, 340; locusts in, 199, 628, 683.

Bamboo, Coccids on, in E. Africa, 611; Ochrophara montana on, in Burma, 339; Mycalesis gotama on, in China, 674; Acridid damaging, in Formosa, 769.

Banana, 269; pests of, in Australia, 210, 211, 494; thrips on, in Bermuda, 626; Ligyrus bitu-berculatus on, in Brazil, 502; pests of, in Ceylon, 360; pests of, in Formosa, 263; Aspidiotus destructor on, in Italian Somaliland, 641; Kophene cuprea on, in Madras, 499; pests of, in West Indies, 10, 145, 175, 223, 491.

Banana Rust Thrips (see Scirtothrips signipennis).

Banana Weevil (see Cosmopolites sordidus).

Anthonomus against Banding. pomorum, 613, 659; against ants, 93, 202, 592; against Cydia spp., 108, 168, 169, 192, 259, 291, 401, 442, 479, 507, 525, 594, 613, 713; against other Lepidoptera, 419, 715; lists of insects found in, on fruit-trees, 689; materials for, 108, 168, 169, 291, 507, 525, 613, 689, 713, 715; treatment of materials for, 108, 168, 169, 291, 479, 507, **525, 613.** (See Adhesives.)

banian, Hieroglyphus. bancrofti, Tarsonemus Boabab (see Adansonia).

Barathra brassicae, in forest nur-series in Korea, 767; parasite of, on crucifers in Japan, 605; food-plants of, in Ukraine, 159; as laboratory host of Tricho-

gramma spp., 103, 235.

Barbados, cotton pests in, 222, 446; sugar-cane pests in, 221, 222: sweet potato pests in, 222, 446: beneficial insects and biological control in, 222, 448; summaries of plant pest legislation in, 134, 385, 491; Hyperaspis trilineata imported into Porto Rico from, 223.

Barbasco, identity of plant known as, in S. America, 319.

barbita, Magdalis.

Barbuda, risk of importation of Platyedra gossypiella into Antigua from, 446.

Baris interstitialis (callida), introduced into Australia to destroy Xanthium pungens, 450; liable to attack artichokes, 450.

Barium Chloride, spraying with, 432, 527.

Barium Fluosilicate, in baits, 387, 457, 528, 639, 757; dusting with, 129, 133, 185, 186, 206, 299, 388, 447, 478, 483, 526, 639, 645, 654, 713, 736, 757; carriers for, 185, 186, 206, 299, 526, 639, 645, 736; oil-impregnated dusts of, 478; spraying with, 186, 201, 205, 478, 697; lime as an adhesive for, 186; ineffective in paint against borers, 543; test of toxicity of, to insects, 348, 387, 388, 572; formulae containing, 185, 186, 201, 205, 206, 299, 387, 478, 526, 639, 645, 736, 757; and cryolite, 736; and magnesium arsenate, 478; and sulphur, 757; tobacco foliage scorched by, 133; Dutox form of, 447, 736.

Bark-beetles, ecology of, in pine forests in Estonia, 461; records of, in Russian Union, 73; problems of, in Sweden, 611, 634; correlation of outbreaks of, with sun spots, 486; natural enemies of, 394; associated with Ceratostomella ulmi on elm, 20, 325, 547, 551, 656, 657, 658, 749; associated with blue-stain fungi on 386, 634; conifers, aseptic rearing of, 28. (See Scolytids.)

Barley, Aphids on, in Argentina, 289, 290; pests of, in Britain, 296, 382, 631; Mayetiola destructor on, in Cyprus, 108; Lema melanopa on, in Czechoslovakia, 657; pests of, in Germany, 51, 520, 521; Spodoptera mauritia on, in India, 241; slightly attacked by Lema tristis in Manchuria, 767; pests of, in Morocco, 500, 501, 641; Cirphis unipuncta on, in Nova Scotia, 441; pests of, in Russian Union, 356, 427, 617; wheat replaced by, against Aelia rostrata in Spain, 698; Pentatomids on, in Switzerland, 91; pests of, in U.S.A., 122, 144; type of injury to, by Pentatomids, 51, 520, 521; effect of soil fumigants on, 631.

Barley (Stored), pests in, and their control, 389, 390, 414; effect of fumigated, on animals, 414.

barodensis, Aleurolobus.

Barriers, against Blissus leucopterus, 63, 78; against Cirphis unipuncta, 723; against Coleoptera, 737, 757; against locusts, 611, 698; against clover mites, 85.

Barylypa perspicillator, parasite of Lymantria dispar in Jugoslavia, 760.

Barypithes pellucidus, measures against, in conifer nurseries in Britain, 625.

basalis, Microphanurus.

basifulva, Erycia.

Basilarchia archippus, parasite of, on willow in U.S.A., 744.

basilare, Xylobiops.

Bassia latifolia, tests of insecticidal

properties of, 360, 361.

Bassus [Braconid] (see Microdus). Bassus laetatorius [Ichneumonid], parasite of Allograpta exotica in Argentina, 289; parasite Cydia molesta in Italy, 154.

batatae, Euscepes (see E. postfas-

ciatus).

Bathyplectes curculionis, parasite of Hypera variabilis in California.

bathyzona, Dacnusa.

Batocera rubus, on fig in India, 765; on tapped jelutong trees in Malaya, 109, 110; against, 110. measures

Batrachedra arenosella, on coconut

in Fiji, 497.

Batrachedra atrilogua, on coconut in Fiji, 497.

Bats, destroying Locusta migra-

toria, 629.

Bauhinia reticulata, Helopeltis bergrothi on, in Sudan, 513.

Baumspritzmittel, 188, 234, 250.

(See Tar Distillates.)

Beach Plum (see Prunus maritima). Bean Beetle, Mexican (see Epilachna varivestis).

Bean Bruchid (see Bruchus obtectus). Bean Fly (see Agromyza phaseoli).

Bean Midge, Seedling (see Campto-

cladius macleayi). Bean Thrips (see Hercothrips fas-

Beans, Heliothis armigera on, in S. Africa, 364; Maruca testulalis on, in Brazil, 643; Aphis rumicis on, in Britain, 527; immune from certain Bruchids in Burma, 7; Cosmolyce boetica on, in Cyprus, 108; Melolontha on, in Germany, 521; pests of, in Hawaii, 214, 752; Myzus persicae on, in Japan, 265; Pseudococcus kenyae on, in Kenya, 592; A. rumicis on, in Libya,260; pests of, in New SouthWales, 209, 334, 335, 755; Tetranychus on, in Russian Union, 257, 471; Bemisia gossypiperda on, in Sudan, 513; pests of, in U.S.A., 12, 19, 127, 129, 270, 285, 327, 333, 545, 566, 580, 644, 649, 651, 704; susceptibility of varieties of, to Hercothrips fasciatus, 129; insects and virus diseases in, 12, 265; as trapcrops for Melolontha, 521; stimulating effect of copper sulphate and lime on, 333; translocation of constituents of derris in, 332.

Beans (Stored), Bruchids and their control in, 284, 414, 415, 649, 746; methyl bromide not impairing germination of, 389, 390. Beans, Lima (see Lima Beans).

Beans, Soy (see Soy Beans).

Beans, Sword (see Canavalia ensiformis).

Beauveria bassiana, infesting Diprion pini in Holland, 348.

Beauveria globulifera, infesting Blissus leucopterus in U.S.A., 78.

beckii, Lepidosaphes.

Beech, pests of, in Britain, 296, 711; Pyrochroa coccinea on, in France, 316; pests of, in Germany, 69, 344, 345, 419; Coccid on, in Massachusetts, 547; Melolontha on, in Rumania, 721.

Bee-hole Borer (see Xyleutes cera-

micus).

Bees, Black Stingless (see Melipona). Bees Honey, Acarapis spp. attacking, 343, 655; insect enemies of, 115, 133, 184, 344, 655, 664; other pests of, 655, 664; Mermis subnigrescens in, 746; use of honey obtained from Aphid honey-dew by, 460; influence of food on dimorphism in, 103; risk of insecticides to, 19, 69, 368, 748; treatment for arsenic poisoning in, 255; quassia sprays harmless to, 5; use of, for testing insecticides, 109.

Bees, Leaf-cutting, constructing nests in figs in California, 292; damaging fruit trees etc. in Germany, 522. (See Melipona

testacea.)

Beeswax, in banding mixtures, 93, 202, 480.

Beet, Tanymecus palliatus on, in Austria, 255; pests of, in Britain, 296, 383, 462, 527, 631; pests of, in Czechoslovakia, 656, 657; Pegomyia hyoscyami var. betae on, in France, 314; pests of, in Germany, 419, 461, 520, 521, 717; flea-beetles on, in Hungary, 197; pests of, in Italy, 161, 184, 461, 527, 528; Disonycha laevigata on, in Jamaica, 490; pests of, in Morocco, 416, 417, 526; pests of, in Poland, 255, 256; pests of, in Russian Union, 94, 157, 159, 235-237, 308, 356, 423, 424, 470, 472, 473; Phthorimaea atriplicella on, in Switzerland, 689; pests of, in U.S.A., 19, 44, 127, 134, 280, 282, 283, 322, 445, 691, 743, 747; insects and virus diseases of, 419, 445, 496; insecticides causing injury to, 236, 631, 747; in baits for cutworms, 160; mites and their control in stored seed of, 159.

Beet Armyworm (see Laphygma exigua).

Beet Leaf Bug (see Piesma quadratum).

Beet Leafhopper (see Eutettix tenellus).

Beet Webworm (see Loxostege sticticalis).

Beet Weevil (see Cleonus punctiventris).

Beet Weevil, Black (see *Psalidium* maxillosum).

Beet Weevil, Grey (see Tanymecus palliatus).

Begonia, thrips on, in greenhouses in Vancouver, 483; insecticides

causing injury to, 484.

Belgium, Agromyzids on chicory in, 313; Leptinotarsa decemlineata in, 120, 368, 719; Melolontha hippocastani in, 193; Myzus ornatus in, 589; parasite of Stilpnotia salicis in, 576; summary of plant quarantines in, 134; Lampyris noctiluca introduced into Australia from, 450.

Bell Heather (see Erica).

bellamyi, Pyrophorus.
Bellicositermes (see Macrotermes).

bellicosus, Macrotermes (Bellicositermes).

Belvosia unifasciata, parasite of Cirphis unipuncta in Connecticut, 723.

Bemisia, new parasite of, in Porto Rico, 41.

Bemisia goldingi, transmitting leafcurl of cotton in Nigeria, 666, 667.

Bemisia gossypiperda, on cotton in Belgian Congo, 315; food-plants and relation to virus diseases of, in India, 89, 242, 409; food-plants of, in Sudan, 512, 513.

Bemisia manihotis, sp. n., possible vector of cassava mosaic in Madagascar, 413, 455, 706; possibly in Réunion, 706.

Bemisia myricae, new parasite of, in Japan, 296.

bemisiae, Prospaltella. benedettoi, Empoasca.

benefica, Eumicrosoma; Pleurotropis; Pseudoleucopis.

beneficiens, Telenomus (Phanurus). bengalella, Heterographis.

bennigseni, Ootheca.

Bentonite, in dusts, 226, 648; uses of, in sprays, 79, 272, 323, 368, 448, 554, 555, 562, 652, 732, 733, 750; formulae containing, 226, 554, 652, 732, 733.

Bentonite Sulphur, in dusts, 544; uses of, in sprays, 25, 79, 149,

150, 448, 510, 750; formulae containing, 25, 149, 510.

Benzene, injected into clove trees against Cerambycids, 295; as a solvent, 190, 412, 707.

Benzine, for extracting toxic principles of plants, 206.

Benzoic Acid, in baits for ants, 222; esters of, attracting Cydia pomonella, 137.

Berberis nepalensis, Xyleborus arquatus on, in Ceylon, **359.**

Berberis vulgaris, Rhagoletis cerasi not observed in fruits of, in Switzerland, 156.

bergenstammi, Apiomyia (Oligotrophus).

bergrothi, Helopeltis.

berlesei, Caloglyphus; Podapolipus; Prospaltella.

Bermuda, miscellaneous pests in, 220, 626; summaries of plant pest legislation in, 229, 491.

Bermuda Cedar (see Juniperus bermudiana).

Berseem (see Trifolium alexandrinum).

betae, Pegomyia hyoscyami; Pemphigus.

Beta-naphthol (in oil), bands treated with, against Cydia pomonella, 108, 168, 169, 507, 525, 613; repellent to Anthonomus pomorum, 613; formulae and dipping methods for, 108, 507; effect of, on apple bark, 479, 480.

Bethelium mundum, on Eucalyptus in New Zealand and Australia, **754.** bhurmitra, Boarmia (Ectropis).

Bibio spp., new Nematode parasites of, in Denmark, 49, 50.

bibionipennis, Aegeria. bibionis, Neoaplectana. bicinctus, Hercinothrips.

bicolor, Brachytes; Leefmansia; Pegomyia.

bicolor, Curt., Microgaster (see Macrocentrus abdominalis).

bicolor, Nees, Apanteles (Microgaster) (see A. circumscriptus).

bicristata, Eutinophoea.

Bidens pilosa, Ĥelopeltis bergrothi on, in Sudan, 513.

bidentata, Euetheola (Dyscinetus). bidentatus, Ips (Pityogenes).

bifasciata, Aciocercops; Comperiella. bifasciatus, Anastatus; Physcus. bifidalis, Loxostege (Phlyctaenodes).

bifoveolatus, Dinoderus. biguttula, Empoasca (Chlorita).

biguttulus, Cremastus (see C. flavoorbitalis).

Bihar, sugar-cane moth borers in,

bimaculata, Aphthona; Virachola (Deudorix): Sturmia (see inconspicua).

bimaculatus, Schistoceros; chus (see T. telarius). bioculatus, Perilloides (1 Tetrany-

(Perillus); Tetranychus.

Biological Control, general account of, 247.

Biosteres caudatus (parasite Trypetids), in W. Africa, 492; introduced into Hawaii, 492.

Biosteres longicaudatus, parasite of Dacus spp. in Ceylon, 359; prointroduction of, posed Hawaii, 359.

bipunctata, Adalia (Coccinella).

bipunctatus, Collops; Echinocnemus; Nephotettix.

bipunctifer, Schoenobius. bipustulatus, Chilocorus.

Birch, Phyllotoma nemorata on, in Austria, 215; weevil on, in Germany, 344; pests of, in Russian Union, 75, 429; P. nemorata on, in U.S.A., 215, 745; pests intercepted in Vancouver from Britain in crates of, 483; susceptibility of timber of, to Codiosoma spadix, 354.

Birch Leaf-mining Sawfly $Phyllotoma\ nemorata$).

Birds, destroying noxious insects, 40, 61, 81, 128, 145, 169, 180, 255, 282, 291, 294, 341, 348, 429, 468, 520, 590, 598, 625, 629, 655, 675, 682, 715; question of danger of poisoned locusts to, 180; protection and value of, 120, 625, 633, 634, 675; insects spread by, 378, 446, 455; nests of, harbouring pests of stored products, 69.

Bischofia javanica, Phassus signifer on, in Formosa, 517.

Biscirus lapidarius, utilisation of, against Smynthurus viridis Australia, 84.

biselliella, Tineola.

Bishop Fly (see Lygus pratensis).

bisignifer, Anthonomus.

bispinosa, Frankliniella cephalica. bituberculatum, Nodostoma.

bituberculatus, Dolichoderus; Ligyrus.

bivittatus, Melanoplus.

Black Leaf 50, 387.

Black Leaf 155, forms of, 544, 554, 555, 556, 653, 733.

Black-headed Fireworm (see Rhopobota naevana).

Black-tipped Locust (see Chortoicetes terminifera).

Blackberry, Lygus pabulinus on, in Britain, 297; Nodonota puncti-collis on, in New York, 288; Typhlocyba froggatti on, in Queensland, 495.

blackburni, Chelonus.

blanchardi, Parlatoria.

Blapstinus substriatus (gregalis), bionomics and control of, in Montana, 26-28; synonymy of,

Blastophagus (see Myelophilus). Blastothrix sericea, parasite Lecanium in California, 18.

Blattella germanica, absorption of hydrocyanic acid by, **567.**

Blechnum, Heptamelus ochroleucus on, in Britain, 152.

Blissus iowensis, sp. n., on Andropogon furcatus in Iowa, 225.

Blissus leucopterus, on sugar-cane in Porto Rico, 40; on cereals and grasses in U.S.A., 63, 78, 124, 126, 144, 225, 283; resistance of varieties of maize and Sorghum to, 63, 125; bionomics of, 78, 144, 283; measures against, 63, 78, 125.

Blissus occiduus, in Saskatchewan, 170.

blitealis, Noorda.

Blood Albumen, as an adhesive for * dusts, 388; in sprays, 280, 542; mopping mixture against cotton boll weevil, 645.

Blosyrus ipomoeae, sp. n., on sweet potato in S. Rhodesia, 106.

Blue-grey Fly (see Aleurocanthus woglumi).

Fungi, Blue-stain bark-beetles associated with, 28, 386, 634.

Blueberry, pests of, in U.S.A., 20, 333; calcium arsenate causing injury to, 333.

Blueberry Maggot (see Rhagoletis pomonella).

Bluegrass (see *Poa*).

Blunt-nosed Leafhopper (see Euscelis striatulus).

Boarmia acaciaria (see Cleora).

Boarmia bhurmitra, on tea in India, 149.

Boehmeria nivea, Cocytodes coerulea on, in Japan, 266.

Cosmolyce (Lampides, boetica, Lycaena).

Bogosiella fasciata, parasite of Dysdercus spp. in Uganda, 374,

bohemani, Anagyrus; Calidea.

bolivari, Prochiloneurus.

bolliana, Gretchena.

Bombay, sugar-cane pests in, 261; Aphids of, 247.

Bombus, azalea flower spot disease spread by, in S. Carolina, 328. Bombyx antiqua (see Orgyia).

Bombyx mori, Aspergillus versicolor experimentally pathogenic 420; Dermestids attacking, 153, 514. 515; used in tests of insecticides, 135, 332, 695.

bondari, Naupactus.

Bonin Islands, Dacus ferrugineus dorsalis in. 672.

boninsis, Pseudococcus.

Books, pests of, 223, 329, 413, 706; bibliography of pests of, 624.

Borax, for treating timber against Lyctus, 190.

borbonica, Parnara.

Bordeaux Mixture, 343; against Epitrix cucumeris, 564, 648; against Jassids, 15, 140, 399, 400, 443, 557, 564, 609, 610, 701; ineffective against Tetranychid mites, 466, 693; repellent to Lepidoptera, 51, 55; not repellent to Stictodiplosis moringae, promoting increase of certain pests, 557, 728; formulae for, 131, 140, 301, 399, 400, 443, 537, 564, 609, 648, 693, 728, 743 formulae containing, 131, 234, 335, 537, 563, 743; and arsenicals, 99, 210, 252, 265, 285, 518, 522, 557, 563, 643, 664, 743; and nicotine, 131, 210, 648; and oils, 210, 298, 537, 728; and soap, 563; and forms of sulphur, 335; and tar distillates, 234, 235; for emulsifying anthracene oil, 301; addition of other insecticides to, 270, 566, 548; causing injury to plants, 210, 648; resistance to rainfall of sprays containing, 522.

Borkhausenia, Hofmannophila considered distinct from, 412.

Borkhausenia pseudospretella, destroying Lepidopterous pupae in cages in Britain, 298; in dried mint in France, 412; in Holland, 183; bionomics and taxonomy of, 412, 413; susceptibility of, to derris, 183.

Borkhausenia stipella on cembra in Austria, 664.

Borneo, Cerambycids on clove trees in, 295; locusts in, 112. bosqi, Celatoria; Lepidaphycus. Bothynoderes (see Cleonus).

botrana, Polychrosis.

Botrytis, infesting insects, 264, 504. Botrytis tenella, infesting Melolontha hippocastani, 193.

boucheanus, Dibrachys (see D. cavus).

Box (see Buxus). boxi, Diaprepes.

Brachartona (see Artona).

brachialis, Bruchus.

Brachyacma palpigera, on Tephrosia spp. in Porto Rico, 224.

Brachycaudus pruni, auct. Anuraphis padi).

Brachycerus citriperda, sp. n., on Citrus in S. Africa, 106.

Brachychiton, weevils on, in New South Wales, 404.

Brachyderes incanus, on pines in Germany, 312.

Brachyexarna lobipennis, in Australia, 584.

Brachymeria, parasite of Homona coffearia in India, 149; reared from Dipterous parasites in Jugoslavia, 760.

Brachymeria femorata, bionomics of, in Egypt, 456.

Brachymeria fiskei, parasite of Clania minuscula in China, 245.

Brachymeria intermedia, parasite of Polia oleracea in Italy, 528; parasite of Lymantria dispar in Jugoslavia, 760.

Brachymeria mikado, parasite of Clania minuscula in China, 245. Brachymeria obscurata, hosts of, in

Japan, 605, 669.

Brachymeria ovata, parasite Desmia funeralis in California,

Brachymeria rubripes, parasite of Tortrix postvittana in Tasmania, 10.

Brachymeria victoria, parasite of

Tortricids in Australia, 10, 169. Brachymeria vitripennis, hosts and utilisation of, in Italy, 528.

brachyntera, Thecodiplosis domyia).

Brachypeplus pilosellus, on cotton in Belgian Congo, 315.

Brachyplatys testudonigra, Dolichos in Nyasaland, 114.

Brachyrrhinus (see Otiorrhynchus). Brachytes bicolor, on Asparagus falcatus in Ceylon, 359.

Brachytrypes, baits for, on Aleurites spp. in Burma, 340.

Brachytrypes membranaceus. cotton in Tanganyika, 642. Brachytrypes portentosus, on cotton

in Formosa, 309.

Brachyurothrips hargreavesi, food-

brachyurus, Stiamus.

Bracon, Ipobracon recorded as, in Brazil, 704.

Bracon intercessor (see Microbracon). Braconids, of Japanese Empire, 247.

Bract Scale (see Atriplex bracteosa). Bran (including middlings, pollard, etc.), insects infesting, 322; in baits, 28, 40, 45, 58, 138, 139, 170, 209, 270, 282, 326, 333, 393, 398, 410, 441, 540, 566, 574, 608, 609, 639, 685, 686, 698, 699; baits for Ceratitis prepared from, 699, 700, 713; formulae containing, 45, 138, 139, 170, 209, 393, 398, 410, 540, 566, 574, 609, 639, 685, 686, 698, 699.

brasiliensis, Aenasius (Chalcaspis); Gasterocercodes; Pseudiastata; Theresia (Paratheresia).

Brassica pe-tsai, Aphids transmitting mosaic disease of, in California, 577.

brassicae, Aphidius; Barathra; Brevicoryne; Dasyneura; Phorbia (Hylemyia); Pieris; Plusia (Autographa).

Brassolis sophorae, on coconut in Br. Guiana, 369, 491; legislation against, 491.

Brazil, cacao pests in, 106, 405, 643; pests of Casuarina in, 145; Citrus pests in, 82, 502, 607; coffee pests in, 319, 607: cotton pests in, 115, 268, 333, 405, 502-504, 505, 689, 704; miscellaneous pests in, 106, 115, 502, 642; pests of solanaceous plants, 145, 290, 320; sugar-cane pests in, 82, 369, 370, 502; leaf-cutting ants in, 248, 533; Coccids in, 82, 212, 319, 367, 502, 607; natural enemies of honey-bees in, 115, 655; locusts in, 290; new weevils in, 106, 115, 405, 612; new Nematode associated with Gasterocercodes brasiliensis in, 504; beneficial insects in, 82, 212, 319, 369, 370, 504, 704; introduction of *Tetrastichus gif-faudi* aggint Tetrastichus gif-faudi aggint Tetrastichus gif-faudi fardi against Trypetids in, 627; rotenone-yielding plants in, 319; summary of plant quarantines in, 751; beneficial inintroduced into other sects countries from, 212, 223.

Brazilian Cotton Borer (see Gastero-cercodes brasiliensis).

Bregmatothrips iridis (Iris Thrips), bionomics and control of, in U.S.A., 273; possibly in Canada, 273.

bremii, Ceratitis.

Brenthia dendronympha, sp. n., on Shorea robusta in India, 68.

brevicomis, Dendroctonus. brevicornis, Microbracon.

Brevicoryne brassicae (on crucifers), in Britain, 463, 632; transmitting mosaic of Chinese cabbage in California, 577; in Russia, 302, 355; bionomics of, 463, 632; natural enemies of, 302, 464; measures against, 355, 463.

brevipennis, Leptidea. brevipes, Pseudococcus.

brevirostris, Lixus.

brevis, Calotermes (Cryptotermes). brevistylus, Dacus (see D. ciliatus).

breviusculus, Desmidophorus. Bristlegrass (see Setaria).

Brithys crini, bionomics of, in Formosa, **52**.

Brithys pancratii, tests of insecticides on, 35.

British Columbia, forest pests in, 293, 484, 485, 486; pests of greenhouse and ornamental plants in, 293, 483; orchard pests in, 59, 144, 171, 484, 485, 535; Sitona lineata on peas and clover in, 293; Sitodiplosis mosellana on wheat in, 60; grasshoppers in, 486, 680; correlation of outbreaks of insects with sun spots in, 486; parasites and biological control in, 485, 535; pests intercepted in quarantine in, 483.

British Empire, summary of plant

pest legislation in, 120.

British Isles, asparagus pests in, 32, 33; pests of beet and mangel in, 296, 383, 462, 527; cereal pests in, 102, 296, 382, 527, 709; pests of crucifers in, 296, 342; 382, 463, 632; forest pests in, 296, 527, 625, 631; importance of protection of birds in forests in, 633; pests of grasses and turf in, 341, 382, 462; pests of heather in, 296, 417; pests of leguminous plants in, 341, 527; miscellaneous pests in, 95, 296, 612, 613, 763; mushroom pests in, 297, 712; Hylemyia antiqua on onion in, 298; orchard pests in, 191, 297, 527, 612, 613, 614, 708, 709; list of insects found in fruit-tree bands in, 689; pests of small fruits in, 103, 296, 297, 383,

384, 612, 613; pests of ornamental and greenhouse plants in, 33, 152, 297, 349, 350, 527, 589, 615, 709, 710, 711, 712, 772; potato pests in, 47, 379, 527, 615, 709, 763; list of pests of stored products in, 165; pests of timber in, 34, 190, 191, 378, 380, 463, 707; new Aleurodids in, pseudos-772; Borkhausenia pretella destroying Lepidopterous pupae in, 298; problem of wireworms in, 383, 462, 630; insects and virus diseases in, 32, 47, 104, **342**, **349**, **379**, **383**, **615**; fungi associated with Siricids in, 632; fungus infesting Syrphid in, 101: beneficial insects in, 33, 102, 342, 382, 384, 463; thrips imported into, against Tetranychus telarius, 712; summary of plant quarantines in, 134; beneficial insects introduced into other countries from, 317, 450, 623; Bufo bufo introduced into Australia from, 449; pests intercepted in Vancouver from, 483.

bromeliae, Diaspis.

Bromus secalinus, relation of wireworms to, in Central Asia, 76.

Brontispa, revision and key to

species of, 499.

Brontispa longissima (froggatti var. selebensis), on coconut in Netherlands Indies, 337, 499; bionomics and synonymy of, 499; utilisation of Tetrastichus brontispae against, 337.

Brontispa sumatrana (see Plesispa

reichei).

brontispae, Tetrastichus (Tetrastichodes).

Brotolomia meticulosa, on chrysanthemum in Britain, 349.

Brown-tail Moth (see Nygmia phaeorrhoea).

Bruchidius (see Bruchus).

Bruchobius mayri (see Sphaerakis).
Bruchophagus gibbus (on lucerne),
in S. Africa, 152; measures
against, in Russia, 76.

Bruchus, measures against, i Burma, 8.

Bruchus albocallosus, bionomics of, on pigeon peas in Burma, 7.

Bruchus analis, bionomics of, in Burma, 7; effect of cold storage on, 8.

Bruchus brachialis (Vetch Bruchid), in U.S.A., 23, 329, 580; intercepted in U.S.A., 24; bionomics of, 24; parasites and biological control of, 24, 329, 580; other measures against, 24.

Bruchus chinensis, bionomics of, in Burma, 7; experiments against, in Mysore, 361.

Bruchus lentis, tests of trichlorethylene against, in stored lentils in France, 414.

Bruchus maculatus, bionomics of, in California, 648, 649.

Bruchus notatus, on Acacia confusa in Formosa, 310.

Bruchus obtectus (Bean Bruchid), in Central Europe, 197, 301; in U.S.A., 389, 648, 649, 746; bionomics of, 301, 648, 649; effect of heat on eggs of, 284; methyl bromide as a fumigant against, 389, 746; used for testing pyrethrum, 197.

Bruchus phaseoli, bionomics of, in Burma, 7.

Bruchus pisorum (Pea Bruchid), in W. Australia, 754; in Hungary, 197; history of outbreaks of, in Japan, 669; in Ukraine, 75; in U.S.A., 26, 65, 291, 327, 329, 653; effect of infestation by, on germination of peas, 26; bionomics of, 75, 653; parasites and biological control of, 65, 329, 654; other measures against, 65, 75, 291, 327, 654, 754; tests of pyrethrum on, 197.

Bruchus quadrimaculatus (see B. maculatus).

macaiais

Bruchus rufimanus, tests of trichlorethylene against, in stored beans in France, 415.

Bruchus terrenus, B. notatus misidentified as, 310.

brui, Thripoctenus.

brumata, Operophtera (Cheimatobia). Brumus octosignatus, bionomics and utilisation of, in Central Asia, 238.

brunnea, Eucolaspis. brunnescens, Hylemvia. brunneus, Lasius; Lyctus. bruzzonis, Tetrastichus. bryanti, Cephalodonta.

Bryobia praetiosa, measures against, in orchards in Washington, 488.

bubalus, Ceresa.

Bucculatrix gossypii, on cotton in Queensland, 494.

Bucculatrix thurberiella, on cotton in California, 19.

bucephala, Phalera.

Buckwheat, Cirphis unipuncta on, in Quebec, 145.

Bud Moth, Eye-spotted Apple (see Eucosma ocellana).

Buettneria pilosa, Phassus on, in Burma, 339.

Bufo bufo, introduced into Australia, 449; doubtful value of, against Oncopera, 449.

Bufo bufo miyakoensis, introduced into Loochoo Islands against noxious insects, 605.

Bufo insularis, destroying Lachnosterna smithi in Mauritius, 317.

Bufo marinus (destroying insects), in W. Indies, 40, 222, 317, 490; establishment of, in Fiji, 59; feeding on Bufo melanostictus in 317; problematical Formosa, value of introduction of, into Mauritius, 317; establishment of, in Queensland, 56; Dytiscid larvae destroying tadpoles of, 222.

Bufo melanostictus, introduced toad destroying, in Formosa, 317.

Building Timbers, Coleoptera in, 17, 34, 91, 155, 190, 191, 195, 354, 379, 381, 451, 463, 530, 599, 622, 707, 726; termites in. 17, 193, 220, 223, 314, 724, 748; surveys of pests of, in Sweden, 439, 624.

Bulb Mite (see Rhizoglyphus hyacinthi).

Bulgaria, summary of plant quarantines in, 692.

buoliana, Rhyacionia (Evetria).

Bupalus piniarius (Pine Geometrid), tests of dusts against, in Germany, 312; Aspergillus versicolor experimentally genic to, 420.

Bur Clover (see Medicago denticu-

lata).

Burma, Alissonotum impressicolle on sugar-cane in, 294; Bruchids of, 7; forest pests in, 338-340, 688, 763, 765; miscellaneous pests in, 409, 451; beneficial insects and biological control in, 294, 339, 688, 765; introduction of parasites of teak pests into Madras from, 765.

Bush Clover (see Lespedeza capi-

tata).

Busseola fusca, on sugar-cane in Natal, 411.

Butter-nuts (see Caryocar nuciferum).

Butyl Acetate, in baits for moths, 236.

buxi, Neotetranychus; Pinnaspis.

Buxus (Box), Neotetranychus buxi on, in Connecticut, 17.

Byrsocrypta gallarum, on elm in Finland, 201.

Byturus unicolor, measures against, on raspberry in California, 328.

C.

Cabbage, Plutellamaculipennis on, in Australia and New Zealand, 449; pests of, in British Isles, 48, 342, 463, 632; Halticids on, in Czechoslovakia, 657; Pieris rapae on, in Egypt, 456; Contarinia nasturtii on, in Holland, 466; pests of, in India, 241, 409; Aphids on, in Italy, 189; Lepidoptera on, in Java, 675; P. brassicae on, in Poland, 720; pests of, in Russia, 159, 257, 302, 355; pests of, in Spain, 695; Plutella maculipennis on, in Tanganyika, 113; pests of, in U.S.A., 13, 28, 285, 399, 647, 651; insecticides causing injury to, 241, 647.

Cabbage, Chinese (see Brassica petsai).

Cabbage Aphis (see Brevicoryne brassicae).

Cabbage Butterfly (see Pieris rapae). Cabbage Fly (see Phorbia brassicae). Cables, electric (see Electric Cables).

Cacao, pests of, in Brazil, 106, 405, 643; Phosphorus gabonator on, in Br. Cameroons, 469; Desmidophorus breviusculus on, Indo-China, 185; Heliothrips haemorrhoidalis on, in greenhouses in Russia, 358; Selenothrips rubrocinctus on, in Trinidad, 10.

Cacao (Stored), Ephestia spp. in, 11, 301, 315, 435, 523, 599; cold storage of, against E. elutella,

Cacao Moth (see Ephestia elutella). Cacao Thrips (see Selenothrips rubrocinctus).

Cacoecia (see Tortrix).

Cacoecia fumiferana (see Harmologa).

cacti, Chilocorus.

Cactoblastis cactorum, breeding of, against Opuntia in S. Africa, 202; Nosema causing disease in, 202. cactorum, Cactoblastis; Eriophyes.

cadaverum, Glycyphagus.

caesalis, Margaronia.

Caesalpinia coriaria, Apate terebrans on, in Jamaica, 491.

caesar, Zenillia. caesarion, Cryptolucilia.

Cages, for rearing and studying insects, 4, 143, 466, 561, 569, 595, 596, 650, 659, 717, 731, 745; for testing insecticides on honey bees, 109; descriptions of, 109, 143, 466, 561, 650, 659, 717.

caillaudi, Julodis.

Cajanus cajan (indicus), pests of, in E. Africa, 114, 592, 642; pests of, in Burma, 7, 451; pests of, in India, 181, 242, 361, 451; Frankliniella insularis on, in St. Vincent, 175.

calamistis, Sesamia.

Calandra granaria (Grain Weevil), in Germany, 519; in stored ginger in Jamaica, 491; in U.S.A., 116, 389, 390, 746; egg-production of, in relation to age, 90; measures and fumigation tests against, 116, 389, 390, 519, 632, 741, 746; tests of insecticides on, 183, 197; monograph on, 519.

Calandra oryzae (in stored cereals), in Assam, 409; in France, 414, 415; in Hawaii, 584; in Indo-China, 206; in Java, 311, 336, 639; in Madagascar, 455; in S. Rhodesia, 203, 608; in U.S.A., 116, 218, 362, 389, 642; on field maize, 203, 608, 643; in stored cassava tubers, 455; bionomics of, 90, 158, 203, 311, 336, 584, 608, 639, 643; uses and tests of fumigants against, 116, 362, 389, 409, 414, 415, 567, 633, 741; other measures against, 116, 203, 218; toxicity of extracts of Stemona to, 206.

calcaratus, Syagrus. Calcid, **86**, **663**, **703**.

Calcium Arsenate, 651; against cereal pests, 285, 309, 310, 757; against cotton pests, 142, 276, 277, 279, 328, 627, 644, 645, 691, 766; against forest pests, 69, 80, 393; against orchard pests, 25, 79, 174, 179, 252, 254, 255, 291, 333, 401, 478, 479, 489, 544, 559, 713; against pests of solanaceous plants, 11, 64, 252, 328, 361, 418, 518, 564, 606, 643, 648; against pests of other plants, 55, 76, 301, 333, 443, 566, 664, 675, 691, 713, 743; repellent to Lachnosterna, 226; ineffective against pea Bruchid, 654; tests

of toxicity of, to insects, 16, 17, 182, 388, 390, 572, 741, 742; in baits, 172, 209, 270, 393, 441, 540, 705; formulae for, in dusts, 142, 226, 276, 277, 333, 443, 566, 643, 644, 766; carriers for, 142, 226, 276, 333, 443, 566, 572, 643, 644, 645, 705, 766; aeroplane dusting with, 627, 654: oil-impregnated dusts of, 478; spraying with, 11, 25, 55, 79, 80, 252, 254, 255, 291, 301, 309, 333, 393, 401, 418, 479, 518, 544, 559, 564, 566, 606, 648, 664, 676, 743; other methods of applying, 644, 645; adhesives and spreaders for, 285, 310, 489, 522, 566; question of suspension power of, in sprays, 252, 518; resistance of sprays of, to rainfall, 522; and Bordeaux mixture, 252, 285, 518, 522, 648, 664, 743; and lime, 401, 489, 559, 566; and lime-sulphur, 489; and nicotine laurate, 301; as an emulsifier for oils, 482; and Paris green, 142, 279; and pyrethrum, 559; inverted spray mixtures containing, 174, 175; and injury to plants, 252, 328, 333, 600, 648, 691, 705; problem of residues of, 328, 402; components and analysis of, 252, 331, 567; physical properties of samples of, 282; other insecticides compared with, 16, 17, 25, 79, 182, 279, 310, 328, 333, 361, 388, 401, 402, 479, 645, 741, 742.

Calcium Arsenite, in baits for cutworms, 160; (in dusts), against locusts, 77, 426, 686; substitute for, against locusts, 76, 77; against other pests, 7, 431, 433; oil-impregnated dusts of, 426; aeroplane dusting with, 426, 433; spraying with, 433.

Calcium Caseinate, as a spreader and adhesive for arsenical sprays, 168, 310, 393; contact performance of, 45,

Calcium Cyanamide, soil treatment with, against pests, 190, 281, 412, 718.

Calcium Cyanide, 755; dusting with, 241, 286; as a greenhouse fumigant, 274, 355, 711, 712; as a soil fumigant, 327, 403, 431, 432, 645; tent fumigation with, 86, 117, 176, 177, 703, 748; mixed with stored beet seed against mites, 160; uses of, against ants, 202, 219; for destroying hornets' nests, 108; methods of applying, against

locusts, 426, 427; and oils, 431; forms of, 86, 176, 177, 703, 711; plants injured by, 241, 431, 712.

Calcium Fluosilicate, spraying with, against *Lema oryzae*, **309**, **310**.

calcitrator, Collyria.

Calendra (see Sphenophorus).

Calendula officinalis, Halticus saltator on, in Germany, 52.

Calidea spp., 112.

California, Aspidiotus lataniae on avocado in, 116; Aleurodes azaleae on azalea in, 20, 748; beet pests in, 19, 134, 280, 282, 322, 445; Aphids on Chinese cabbage in, 577; Citrus pests in, 18, 20, 173, 219, 283, 508, 574, 575, 650, 747, 748; cotton pests in, 19, 748; eradication of date scale in, 18; fig pests in, 30, 392; forest pests in, 18, 226, 394, 581, 650; pests of leguminous crops in, 18, 19, 118, 128, 387, 648, 747; miscellaneous pests in, 19, 589, 747, 748; orchard pests in, 18, 19, 20, 62, 745, 747, 748; pests of solanaceous plants in, 19, 20, 64, 117, 217, 327, 328, 744, 747; vine pests in, 19, 285, 286, 324, 748; pests of walnut and other nut trees in, 18, 20, 650, 747; Mayetiola destructor on wheat in, 326; pests of dried fruit in, 29, 118, 228, 324, 389, 573; pests of other stored products in, 116, 573, 588, 649, 748; injurious ants in, 129, 588; Eriophyids of, 173, 650; grass-hoppers in, 18, 19, 748; Scutigerella immaculata in, 280, 625; flooding against wireworms in, 569; insects and virus diseases in, 445, 577; beneficial insects in, 18, 19, 29, 212, 226, 285, 286, 748; Aenasius paulistus imported into, from Brazil, 212; lists of pests intercepted in quarantine in, 116; committee on border quarantines in, 116; beneficial insects introduced into other countries from, 81, 586, 712.

California Fire Ant (see Solenopsis geminata var. maniosa).

California Mixture, composition and use of, against *Paratetranychus* pilosus, **466**.

californica, Aceratagallia; Melan-

ophila.

californicus, Hippiscus; Pheletes (Limonius).

caliginosus, Epipolaeus.

Caliroa limacina, on pear and cherry in California, 748; on pear in Italy, 758; measures against, 748, 758.

Caliroa oishii, bionomics of, on

oaks in Japan, 311.

Calliceras (see Ceraphron).

callida, Baris (see B. interstitialis). Callidium rufipenne, bionomics of, on conifers in Japan, 515, 516. calligraphus, Ips.

Calliphora erythrocephala, susceptibility of, to injections of

nicotine, 332.

Calliptamus italicus (Italian Locust), in Central Asia, 77; in Cyprus, 108; notice of list of parasites of, in Italy, 163; losses and expenditure due to, in Jugoslavia and Iran, 685; in Libya, 260; outbreaks of, in Spain, 698; measures against, 77, 685, 698.

callisto, Apina. callosa, Sitona.

Callosobruchus (see Bruchus). Calluna vulgaris (see Heather).

Calocampa exoleta, fungus infesting, in Germany, 152.

Calocoris norvegicus, ovipositing on tansy in Germany, 234.

Caloglyphus, classification of, 94. Caloglyphus berlesei, type of genus, 94.

Caloglyphus moniezi, sp. n., 94; measures against, on mushrooms in Holland, 656; on cockchafer in Russia, 95.

Caloglyphus (Tyroglyphus) mycophagus, Mégn., identity of mites recorded as, by various authors, 94; new subgenus for, 94.

Caloglyphus mycophagus (Mégn.) Oudm. (see C. moniezi).

Caloglyphus rodionovi, sp. n., 94; distribution of, in stored seeds etc., 95; in stored cereals in Russian Union, 95, 97; effect of temperature on, 97.

Caloglyphus (Isoglyphus) sphaerogaster, sp. n., 94; in decaying

onion in Russia, 95.

Calomel (see Mercurous Chloride). Calomycterus setarius, bionomics of, in Connecticut, 15.

Calophyllum inophyllum, Cratopus aurostriatus on, in Seychelles,

Calosoma inquisitor, in Poland, 256. Calosoma semilaeve (predacious on armyworms), in Mexico, 586; introduced into Hawaii, 586.

Calosoma sycophanta (predacious on Lepidoptera), in Germany, 234; doubt as to establishment

of, in Java, 337.

Calotermes, damaging tea in India, 149; in Netherlands Indies, in Queensland, 706; damaging timber and books, 449, 706; measures against, 149, 220, 223, 338, 449.

Calotermes brevis, damaging timber and books etc. in W. Indies, 220,

223.

Calotermes castaneus. attacking living trees and timber in Bermuda. 220.

Calotermes dispar, sp. n., on chestnut and plum in Canary Islands, 413.

Calotermes flavicollis, in timber in

France, 314. Calotermes havilandi, in timber in Tanganyika, 642.

Calotermes lamanianus, in timber in Sierra Leone, 240.

Calotermes militaris, attacking tea in Ceylon, 240, 337; factors affecting damage by, 338.

ulotermes simplicicornis, in Mexico and U.S.A., **219**; biono-Calotermes mics and taxonomy of, 219.

Calotropis, Dacus longistylus fruits of, in India, 242.

Calotropis procera, decoction against thrips, 88.

calthae, Aphalara.

Camelina sativa, pests of, Ukraine, 257, 258.

Camellia, pests of, in Japan, 311, 438.

camelliae, Curculio (Balaninus); Parlatoria pergandei.

cameratus, Schizotrachelus.

Cameroons, British, Phosphorus gabonator on cacao and Cola nitida in, 469.

Camnula pellucida, in Canada, 680: in U.S.A., 19, 681; bionomics of, 19; cultivation favouring increase of, 680, 681; measures against, 680.

Camphor, Aspidiotus perniciosus not developing on, 305.

campinas, Anthonomus.

Camponotus herculeanus pennsylvanicus, measures against, damaging telephone poles Connecticut, 218.

Campsomeris annulata, parasite of Alissonotum impressicolle Burma, 294.

Campsomeris marginella modesta, parasite of Anomala orientalis in Hawaii, 493; Pyrophorus unlikely to affect, 493.

Campsomeris phalerata, establishment of, against Lachnosterna smithi in Mauritius, 317.

Campsomeris pilosella, establishment of, against Lachnosterna smithi in Mauritius, 317.

Camptocladius macleayi (Bean Midge), factors affecting infesta-tion of beans and cucurbits by, in New South Wales, 209.

Campylomma nicolasi, on cotton in

Sudan, **513**.

Canada, forest pests in, 103, 215, 293, 330, 487, 534, 538; miscellaneous pests in, 121, 133, 173; grasshoppers in, 590, 676, 680, 686; Porosagrotis orthogonia in, 751; summary of insect conditions in, in 1937, 560; beneficial insects and biological control in, 103, 133, 487, 623; beneficial insects introduced into other countries from, 328, 594, 595; historical review of plant legislation in, 23; index to entomological publications of Department of Agriculture in, 440. (See also under the various Provinces.)

canadensis, Epochra; Spathius. Canary Islands, new termite in, 413.

Canavalia ensiformis, new weevil on, in Brazil, 106; Bruchid on, in Burma, 7; Frankliniella insularis on, in St. Vincent, 175. canavaliae, Promecopsis.

cancellata, Schistocerca.

cancroides, Chelifer.

Candezea gossypii (see Monolepta).

candida, Saperda. canella, Diatraea; Paria.

canescens, Nemeritis.

canicularis, Fannia.

Cankerworm, Fall (see Alsophila pometaria).

Cannabis sativa (see Hemp).

cantor, Glenea.

canus, Pheletes (Limonius).

Cape-weed (see Cryptostemma calendulaceum).

Caper, pests of, in Sicily, 92.

capitata, Ceratitis.

capito, Locusta migratoria.

Capitonius saperdae, parasite of Eutetrapha tridentata, in Connecticut, 14.

Capitophorus fragariae, bionomics of, transmitting virus disease of strawberry in Britain, 103, 104, 383, 613; measures against, 104, 383; distinct from C. fragae-folii, 103.

Capitophorus fragaefolii, transmitting virus diseases of straw-berries in Oregon, 291; other food-plants of, 291; C. fragariae distinct from, 103.

Capnodis tenebrionis, on fruit trees

in Malta, 469.

Capparimyia savastani, bionomics and control of, on caper in Sicily, 92, 93.

capparis, Asphondylia.

apsella bursa-pastoris, gossypii on, in China, **245.** Capsella

Capsicum, pests of, in Chile, 654; thrips on, in India, 55: Aphid transmitting mosaic of tobacco to, in Italy, 759; Epitrix parvula on, in Jamaica, 490; Ceratitis capitata on, in Libya, 261; pests of, in U.S.A., 17, 64, 328; problem of arsenical residues on, 328.

capsiformis, Nabis.

Caradrina exigua (see Laphygma). Caragana, Sitona crinita on, in Russia, 432.

Caraway, legislation against Depressaria nervosa in, in Germany, 719.

Caraway Moth (see Depressaria

nervosa).

Carbolic Acid, in spray formula against Coccids, 701; dust containing, against clover mites, 85; as a repellent for Melolontha, 346, 721.

Carbolineum (see Tar Distillates). Carbon Bisulphide, against ants, 219, 533, 588; against borers, 187, 269, 295, 591; against Leptinotarsa decemlineata, 418; as a soil fumigant, 53, 54, 166, 236, 281, 383, 393, 398, 418, 436, 488, 729, 747, 757; against pests of stored products, plant material, etc., 24, 26, 31, 115, 160, 239, 240, 326, 359, 362, 409, 437, 515, 654, 766; relative effectiveness of, against various stages of Dacus spp., 263, 436; unsatisfactory as a greenhouse fumigant against thrips, 274; emulsions of, 398, 436, 729, 747; and carbon tetrachloride, 326, 362, 589; naphthalene and paradichlorobenzene dissolved 219, 591, 747; causing injury to

plants, 54, 437, 747; for extracting toxic principles of pyrethrum, 412.

Carbon Dichloride, toxicity of, to pests of stored cereals, 326.

Carbon Dioxide, combined with ethylene oxide (see also Cartox and T-gas), 38, 396, 413, 540, 571, 572, 597; combined with other fumigants, 539, 540, 571, 572.

Carbon Tetrachloride, unsatisfactory against weevils in Cola nuts, 239; and carbon bisulphide, fumigation with, 326, 362, 363, 589; and ethylene dichloride, against termites in books, 223 for extracting toxic principles of plants, 120, 183, 206, 258, 261, 412, 739, 740.

carbonarium, Monomorium.

Corcelia susurrans (see Zenillia).

carcharias, Saperda.

Cardiac Activity, method of recording, in insects, 439.

cardinalis, Dysdercus; (Novius).

cardui, Vanessa (Pyrameis).

Carica papaya (see Papaya). caridei, Sarcophaga

carinatus, Tetrastichus. carinosa, Syprestia.

cariocus, Aenasius.

Carnation, new weevils on, in S. Africa, 106, 612; new thrips on, in S. Australia, 318; pests of, in Britain, 349, 711, 712; Nephrotoma maculata on, in Switzerland, 689; Tetranychus telarius on, in U.S.A., 648; ceptibility of varieties of, to Thrips tabaci, 489, 711; injured by Paris green and sugar spray, 489.

carnifex, Cheyletus.

Carob Honey, in bait for Trypetids, 108.

Carolina, North, Bruchus brachialis and its parasites in, 24, 580; tobacco pests in, 393, 744.

Carolina, South, cotton pests in, 278, 328, 644, 645; miscellaneous pests in, 327, 643, 644; tobacco pests in, 393, 645; Bombus spreading azalea flower spot disease in, 328.

carolina, Megacephala.

Carolinaia cyperi, food-plants of, in Porto Rico, 39, 40; transmitting sugar-cane mosaic, 40. Carolinaia setariae, in Louisiana,

577; parasites of, on sugar-cane

and grasses in Porto Rico, 39; transmitting sugar-cane mosaic, 40, 577; penetration of setae and stylet sheaths of, 225.

Caroline Islands (see Tapanese

Mandated Islands)

Carpenter Ant (see Camponotus herculeanus pennsylvanicus).

Carpet Beetles, measures against, 227, 400. (See Attagenus piceus and Anthrenus spp.)

Carpocapsa (see Cydia).

carpocapsae, Aenoplex; Ascogaster; Stantonia.

Carpocoris pudicus (including var. fuscispinus), bionomics type of injury to cereals by, in Germany and Switzerland, 51, 52, 92.

Carpoglyphus taiwanensis, sp. n., in sugar in Formosa, 436.

Carpolonchaea (see Lonchaea).

Carpomyia vesuviana, in India, 242, **765**: on Zizyphus jujuba, **765**.

Carpophilus dimidiatus, associated with decaying copra in Malaya, 31; Necrobia rufipes predacious on, 31.

Carpophilus rickseckeri, on tomato

in California, 20.

parasites Carpophthoromyia, in Fr. Cameroons, 492; parasite of, introduced into Hawaii, 492.

Carrot, Cavariella on, in Britain, 296; pests of, in Germany, 468, thrips on, in Hawaii, 692; Papilio machaon on, in Japan, 673; pests of, in U.S.A., 489, 565, 647; as trap-crop for Melolontha, 521; methyl bromide fumigation not impairing germination of seed of, 389.

Carrot Rust Fly (see Psila rosae). Weevil Carrot (see Listronotus

latiusculus).

Cartox, regulations dealing with, as a fumigant in Germany, 38.

caryae, Acrobasis; Curculio: Longistigma.

caryaefoliae, Melanocallis.

caryaefoliella, Coleophora.

caryana, Enarmonia (Laspeyresia). caryivorella, Acrobasis.

Caryocar nuciferum, pests imported into Holland from Dutch Guiana in nuts of, 529.

Case-bearer, Clover (see Coleophora spissicornis).

Case-bearer, Larch (see Coleophora laricella).

casei. Piophila.

Casein, Attagenus piceus feeding on, in U.S.A., 568; in mopping mixture against cotton weevil, 645; as a wetting agent and adhesive for sprays, 542, 724.

Casinaria, parasite of Lymantria monacha in Germany, 233.

Cassava, Erinnyis spp. on, in Cuba, 171; Tetranychus telarius on, in Formosa, 674: pests of, in Madagascar, 413, 454, 455, 706; pests of, in Réunion, 706; pests of, in Sierra Leone, 240: pests of, in Tanganyika, 112, 642; question of relation of Aleurodids to mosaic disease of, 240, 413, 455, 706; pests of stored roots of, 455.

Cassia spp., Xyleutes persona on, in

Burma, 340.

Cassida, species of, and their parasites, on beet in Italy, 527, 528.

Cassida vittata (on beet), in Germany, 461; in Italy, 461, 527, 528; in Morocco, 416; biological control of, 528.

Cassidocida africana, sp. n., parasite of Aspidomorpha tecta in Uganda,

610.

Castalius hamada jezoensis, predacious on Doraphis populi in Japan, 673.

castanea, Aserica; Brontispa (see longissima); Diparopsis; Hyposerica (Serica); Pentilia.

castaneipennis, Planispa. castaneum, Tribolium.

castaneus, Calotermes.

castellanii, Tyrophagus putrescentiae. Castnia licoides (licus, auct.), on sugar-cane in Trinidad, 10.

Castor, pests of, in Ceylon, 359; pests of, in Formosa, 207, 674; Tetranychus telarius on, in India, 55; pests of, in Indo-China, 186, 205; Euproctis producta on, in Sudan, 513; sodium fluosilicate causing injury to, 205.

Castor Oil (see Oil, Castor).

Casuarina, pests of, in Brazil, 145: Icerya purchasi on, in Formosa, 263; species of, immune from Pantophthalmus pictus, 146.

Catagmatus clematidis, sp. n., on Clematis in Sumatra, 106.

catalaunalis, Antigastra,

catenifer, Stenoma.

Cathartus advena (see Ahasverus).

Catolaccus aeneoviridis, hyperparasite of Tortrix rosaceana in Utah, 141.

Catolaccus hunteri, hosts of, in Mexico, 275.

Catopsilia crocale, toxicity of extracts of Stemona to, in Indo-China, 206.

catoxantha, Artona (Brachartona). Cattle, question of risk of insecticides to, 180, 609, 638, 682, 748. Cattleya (see Orchids).

caudata, Lepidiota.

caudatus, Biosteres.

Cauliflower, Aphid on, in Britain, 464; Plutella maculipennis on, in Colorado, 28; Contarinia nasturtii on, in Holland, 466.

cautella, Ephestia.

Cavariella, on carrot in Britain, 296. cavus, Dibrachys.

Cayman Islands, miscellaneous pests in, 268.

Cecidomyia brachyntera (see The-codiplosis).

Cecidomyia seminis, on Pennisetum typhoideum in Sudan, 513.

cecropia, Semia.

Cedrela odorata, new Jassid on, in Porto Rico, 288.

Cedrela toona, Hypsipyla robusta on, in Burma, 340.

cedrelae, Dikraneura.

Cedria paradoxa, utilisation of, against Hapalia machaeralis in India and Burma, 339, 765; alternative hosts of, 765.

Celama sorghiella, on sorghum, maize, etc., in Texas, 279, 560; bionomics and control of, 279, 280.

Celatoria bosqi, sp. n., parasite of Diabrotica speciosa in Argentina, 689.

celer, Microbracon.

Celerio euphorbiae, parasite of, in Czechoslovakia, 529.

Celerio lineata var. livornica, measures against, on vines in Libya, 201.

Celery, Halticus saltator on, in Germany, 52; weevils on, in U.S.A., 565, 747.

cellularis, Perisierola.

cellulosa, Pulvinaria.

Celtis sinensis, Sasakia charonda on, in Japan, 673.

cembrae, Chermes (Pineus).

Centeter cinerea, liberation of, against Popillia japonica in Connecticut, 723, 726.

Centipedes, destroying Scutigerella immaculata in California, 626.

centralis, Monochamus (Monohammus).

Centrobia, parasite of Cassida spp. in Italy, **528.**

centuncularis, Megachile.

Cephaëlis ipecacuanha, Tetranychus on, in Formosa, 674.

cephalica, Frankliniella.

Cephalodonta bryanti, sp. n., on avocado in Brazil, **502.**

cephalonica, Corcyra.

Cephalonomia, parasite of Minthea rugicollis in Malaya, 764.

Cephalonomia tarsalis, parasite of Oryzaephilus surinamensis in Illinois, 369.

Cephalosporium coccorum (see

Verticillium).

Cephalosporium lecanii (infesting Coccids), in Czechoslovakia, 659; in Florida, 587.

Cephonodes hylas, bionomics of, on coffee in Madagascar, 67.

Cephus cinctus, loss due to, on wheat in Saskatchewan, 170.

Cephus pygmaeus (on wheat), in Britain, 102, 527; in Czechoslovakia, 657; in Estonia, 462; on rye, 462; parasites of, 102.

Cerambyx miles, on fruit trees in Malta, 469.

Cerambyx velutinus, on carob in Cyprus, 458.

ceramicus, Xyleutes.

Ceraphron (Calliceras), parasite of Fornicia ceylonica in Ceylon, 639; parasite of Sitodiplosis mosellana in Japan, 670.

cerasi, Rhagoletis.

cerasiops, Picroscytoides. Ceratia cincta, on cuci

Ceratia cincta, on cucurbits in Ceylon, 358.

Ceratitis, African parasites of, introduced into Hawaii, 492.

Ceratitis annonae, parasites of, in W. Africa, 492.

Ceratitis bremii, parasite of, in Kenya, 492.

Ceratitis capitata, in E. Africa, 492; in Bermuda, 626; in Brazil, 628; in Cyprus, 108, 260, 706; in France, 416; in Hawaii, 84, 325; first record of, in India, 602; in Italy, 712, 713; in Libya, 5, 260; in Malta, 469; in Palestine, 77, 406, 514; in Spain, 697, 699; in Switzerland, 715; legislation dealing with imported grapes etc. against, in U.S.A., 229; on Citrus, 5, 77, 261, 406, 469, 514, 626, 706,

713; susceptibility of species of Citrus to, 406, 407; other host fruits of, 84, 229, 260, 416, 602, 715; bionomics of, 406; parasites and biological control of, 84, 492, 628; host-fruits affecting parasitism of, 84; cold storage against, 229, 325; other measures against, 5, 108, 407, 697, 699, 713.

Ceratitis coffeae, in Sierra Leone, 239, 494; parasite of, 492; measures against, on coffee, 239.

Ceratitis colae, parasites of, in E. Africa, 492.

Ceratitis cosyra, parasites of, in E. Africa, 492.

Ceratitis giffardi, parasites of, in Sierra Leone, 492.

Ceratitis nigra, parasites of, in Kenya, 492.

Ceratitis punctata, parasites of, in Africa, 492.

Ceratitis rubivora, parasite of, in Kenya, 492.

ceratomiae, Zenillia.

Ceratonia, Lepidosaphes beckii on, in Palestine, 636.

Ceratonia siliqua, Eumarchalia gennadii on, in Crete, 71; pests of, in Cyprus, 106, 458.

ceratoniae, Myelois.

Cevatostomella ulmi (causing Dutch Elm Disease), in Czechoslovakia, 657, 658; in Holland, 656; in Italy, 119; in U.S.A., 20, 325, 547, 551, 749; relation of insects to, 20, 325, 547, 551, 656, 657, 658, 749; susceptibility of species of elm to, 656; reviews of data on, 119, 749.

Ceratovacuna lanigera (see Oregma).

cerealella, Sitotroga.

Cereals, pests of, in Argentina, 31, 705; pests of, in Britain, 296, 382; pests of, in Czechoslovakia, Lamellicorns on, Germany, 717; Trachea secalis on, in Holland, 299; locusts damaging, in India, 628; Mayetiola destructor on, in Morocco, 640; pests of, in Nyasaland, 114; Chortoicetes terminifera damaging. in Queensland, 494; pests of, in Russia, 422, 425; pests of, in Saskatchewan, 170; Hybalus cornifrons on, in Sicily, 757; pests of, in U.S.A., 19, 44, 126, 140, 580, 696, 723; type of injury to, by Pentatomids, 51, 92. 520, 521; review of literature on pests of, 386; stored grain mites surviving on, under field conditions, 422. (See Maize,

Wheat, etc.)

Cereals (Stored), pests of, and their control, 11, 38, 64, 74, 95-97, 116, 125, 159, 160, 202, 326, 389, 409, 413, 414, 415, 422, 501, 519, 568, 643, 644, 746; apparatus for recording absorption of ethylene oxide by, 414.

Ceresa bubalus, bionomics and control of, in France, 343; in Hungary and U.S.A., 343.

Cerocephala, probably hyperparasite of Minthea rugicollis in Malaya, 764.

Ceromasia juvenilis, parasite of Pyrausta nubilalis in Poland, **623**.

Ceromasia senilis, establishment of, against Pyrausta nubilalis in U.S.A., 730; Lydella stabulans grisescens considered correct name for, 730.

Ceroplastes giganteus, parasite of, on Ficus in Haiti, 41.

Ceroplastes rubens, measures against, on shrubs in New South Wales, 335.

Cerura vinula (see Dicranura).

Cetonia aurata, on roses and fruit trees in Germany, 717.

Ceuthorrhynchus macula-alba, bionomics and control of, on poppy in Czechoslovakia, 659.

Ceuthorrhynchus sareptanus, bionomics and distribution of, on cultivated and wild flax in Siberia, 6, 351, 475, 601; in Russian Far East, 351; measures against, 6.

Ceuthorrynchus syrites, bionomics of, on Camelina sativa in Ukraine,

257.

Ceylon, Coccids of, 184; coconut pests in, 359, 360, 434; miscellaneous pests in, 8, 358-360; tea pests in, 240, 337, 360, 639; beneficial insects and biological control in, 337, 359, 434, 639; summary of plant quarantines in, 692; proposed introduction of beneficial insects into other countries from, 204, 359.

ceylonica, Fornicia.

Chaetochloa setosa, Sipha flava on, in Porto Rico, 39.

Chaetocnema, on Agropyrum and cereals in Russia, 428.

Chaetocnema aridula, on cereals in Russia, 427.

Chaetocnema concinna, on strawberry in Britain, 612; on beet and Fagopyrum tataricum in Poland, 255.

Chaetocnema hortensis, on cereals in Russia, 427.

Chaetocnema tibialis, on beet in Italy, 528.

Chaetodacus (see Dacus).

Chaetoptelius vestitus (see Hylesinus). Chaitophorus lyropictus (see Periphyllus).

Chalcaspis, characters of, 212.

Chalcaspis brasiliensis (see Aenasius).

Chalcidoidea, of India, 440.

Chalcis, auct. (see Brachymeria).

chalcites, Plusia.

Chalcodermus aeneus, experiments against, on cowpeas in Alabama, 572, 691.

chalcographus, Ips (Pityogenes). Chalcoides (see Crepidodera).

Chalcophora virginiensis, in pine logs in Minnesota, 43.

Chalepus dorsalis, bionomics of, in U.S.A., 549, 550; measures against, 550.

Chalioides vitrea, on tamarind in Madras, 499.

chalybea, Pirene.

chalybeipennis, Planispa.

chalybeus, Orcus.

chalybophanes, Acrocercops.

chamaeleon, Anomala.

Changa (see Scapteriscus vicinus). chapadae, Aenasius.

Charagia virescens, bionomics and control of, in New Zealand, 266, 267.

Chariessa pilosa (with var. onusta), predacious on Chrysobothris femorata in Oklahoma, 137.

Charips grioti, sp. n., 290; hyperparasite of Toxoptera graminum in Argentina, 31, 289, 290; in Uruguay, 290; recorded as Xystus sp., 31.

Charips minuta, hyperparasite of Brevicoryne brassicae in Ukraine,

302, 303.

charonda, Sasakia.

Charops dominans, parasite of Spodoptera mauritia in India, 241.

Chasmodon apterus, parasite of Geomyza tripunctata in Britain, 382.

Cheese, Tyroglyphus farinae infesting, in Ontario, **539.**

Cheese Fly (see Piophila casei).

Cheimatobia brumata (see Opero-phtera).

Cheirinia repanda, relation of, to Psylliodes punctulata on beet in Utah, 743.

Cheletomorpha venustissima, predacious on grain mites in Russia,

95.

Chelifer cancroides, in bee-hives, 664; predacious on Galleria mellonella in Germany, 664.

Chelonella nitobei (see Chelonus

pectinophorae).

Chelonus, parasite of Spodoptera mauritia in India, 241; parasite of Platyedra gossypiella in Tanganyika, 112.

Chelonus albobasilaris, parasite of bollworms and boll weevils in

Mexico, 275.

Chelonus annulipes, importation and establishment of against Pyrausta nubilalis in U.S.A., 329, 730.

Chelonus blackburni (parasite of Platyedra gossypiella), attempted establishment of, in Mexico, 275; introduced into Texas from Hawaii, 276; reared on Ephestia kuehniella, 276.

Chelonus pectinophorae, parasite of Cydia glycinivorella in Manchuria, 452.

Chelonus texanus, parasite of cotton pests in Mexico, 275.

Chenopodium, relation of Schistocerca paranensis to, in Argentina, 147; Pegomyia hyoscyami var. betae on, in France and Germany, 314; effect on Euxoa segetum of feeding on, 424.

Chermes, on conifers in Germany, 189, 252, 253, 344; type of injury to silver fir by, 252, 253.

Chermes abietis, on spruce in Czechoslovakia, 657; development cycles of, on spruce and larch in Switzerland, 459; in U.S.A., 80, 215; effect of attacks of, on spruce, 80; carried by wind, 215; C. viridis not considered distinct from, 459.

Chermes cembrae, bionomics and control of, in Germany, 762; C. sibiricus preferred name for, 762.

Chermes ishiharai, bionomics and control of, on spruces in Japan, 438.

Chermes japonicus, bionomics and control of, on spraces in Japan, 438.

Chermes nordmannianae, on silver fir in Czechoslovakia, 657; bionomics of, in Germany, 3, 253,

660: type of injury to silver firs by, 253; measures against, 661. Chermes nüsslini (see C. nord-

mannianae).

Chermes piceae, on silver fir in Britain, 527; susceptibility of introduced and native silver firs to, in Germany, 344; Leucopis obscura imported into U.S.A. against, 329.

Chermes pini, liberation of parasites against, on pines in Aus-

tralia, 449.

Chermes prelli, bionomics of, in Germany, 2-4.

Chermes strobi, on Pinus strobus in

Germany, 189.

Chermes sibiricus (see C. cembrae). Chermes strobilobius, on spruce and larch in Czechoslovakia, 657, 715; bionomics of, 715; measures against, 716.

Chermes viridis, not considered distinct from C. abietis, 459.

chermesina, Rodolia.

Cherry, pests of, in Britain, 612; Pseudococcus on, in Br. Columbia, 59; Rhagoletis cerasi on, in Czechoslovakia, 657; Argyresthia ephippella on, in Finland, 201; R. cerasi on, in France, 526; pests of, in Germany, 52, 156, 347, 522; pests of, in Holland, 299, 465, 466; R. cerasi on, in Italy, 713; Lymantria dispar on, in Jugoslavia, 759; Tortrix spp. on, in New Zealand, 57, 58; Monosteira unicostata on, in Morocco, 340; pests of, in Russian Union, 159, 305, 352, 432, 530; pests of, in Spain, 697; R. cerasi on, in Switzerland, 156, 300; pests of, in U.S.A., 44, 127, 489, 748; book on pests of, in U.S.A., 722; insecticides causing injury to, 466, 489.

Cherry (Fruit), problem of spray residues on, 140, 489, 490; HCN unsuitable as fumigant for, 307.

Cherry Fruit-fly (see Rhagoletis cerasi).

Cherry Scale (see Aspidiotus forbesi). Chestnut, Thysanoptera Britain, 711; new termite on, in Canary Islands, 413; Cydia Canary pomonella on, in Czechoslovakia, 658; Acrocercops on, in Japan, 669; pests found in nuts of, for export from Italy, 384; ants damaging timber of, in Connecticut, 218.

chevrolati, Thonalmus.

Chevletus carnifex, predacious on grain mites in Russia, 95.

Cheyletus eruditus, predacious on other mites in stored seed, etc., in Russian Union, 95, 97, 160; temperature preference of, 97. Chick Pea (see Cicer arietinum).

Chicory, pests of, in Belgium, 313; wireworms on, in Britain, 463.

Chile, Coccids on apple and pear in, 221; pests of Capsicum in, 654.

chilensis, Tetracha (see Megacephala carolina).

Chillies (see Cabsicum).

Chilo, on sugar-cane in China, 243: parasite of, in Madras, 408.

Chilo infuscatellus, on sugar-cane in Formosa, 673; effect of submersion on, 673.

Chilo plejadellus, types of damage to rice by, in Louisiana, 218.

Chilo simplex (on rice), in China, 498; in Japan, 54, 669, 670, 671, 770; effects of infestation by, 671; parasites of, 54, 669, 670, 671; measures against, 498, 770.

Chilo zonellus, on sugar-cane in India, 243, 261, 499; susceptibility of varieties of sugar-cane to, 499; parasite of, 243.

Chilocorus, predacious on Lepidosaphes pinnaeformis in S. Africa.

Chilocorus bipustulatus (predacious on Coccids), in Palestine, 637; in Russian Union, 305.

Chilocorus cacti, introduced into Santo Domingo against Aspidio-

tus destructor, 10.

Chilocorus distigma (predacious on Coccids), in Tanganyika, 112: establishment of, in Seychelles, 433.

Chilocorus politus, utilisation of, against Aspidiotus destructor in Netherlands Indies, 337; introduced into Mauritius against A. destructor, 204.

Chilocorus venipustulatus, predacious on Aspidiotus perniciosus

Russian Union, 305.

Chilocorus wahlbergi, establishment of, against Coccids in Sevchelles, 433.

China, cotton pests in, 245, 304, 498; pests of litchee in, 674, 693; miscellaneous pests in, 244, 497; rice pests in, 89, 109, 244, 245, 498, 674; sugar-cane pests in, 243; records of Aphids including new species in, 265, 367:

Aphids producing tannin galls on Rhus in, 264, 265; beneficial insects in, 89, 109, 119, 244, 245, 693; Bureau of Entomology in, 109; summary of plant quarantines in, 229; Rhizoglyphus echinopus imported into Formosa on Allium from, 674.

Chinch Bug (see Blissus leucop-

terus).

Chinch Bug, False (see Nysius ericae)

Chinch Bug, Western (see Blissus

chinensis, Bruchus: Melaphis (Schlechtendalia); Oxya.

Chinese Cabbage (see Brassica pe-

Chinese Juniper (see Juniperus chinensis).

Chionaspis furfura, measures against, on apple in U.S.A., 272, 561, 738; bionomics of, 561. Chionaspis pinifoliae, bionomics

and control of, on pines in U.S.A., 270, 548, 549, 723.

Chionaspis salicis, on Sorbus aucuparia in Finland, 201.

Chionaspis tegalensis, susceptibility of varieties of sugar-cane to, in Réunion, 501; predacious enemy and control of, 501.

Chionaspis vitis, on Hydnocarpus wightiana in Ceylon, 359.

chionosticta, Antarchaea.

chittendeni, Dialeurodes.

Chlaenius obesus, destroying locust eggs in Nyasaland, 114.

Chloridea (see Heliothis)

Chloris, food-plant of Schistocerca paranensis in Argentina, 147. chloris, Aphis; Nezara.

Chlorita (see Empoasca).

Chlorcresylic Acid, dusting with, against onion fly, 298.

Chlorochroa sayi, on wheat in Saskatchewan, 170; on cereals and lucerne in U.S.A., 44, 126, 170; bionomics of, 44.

Chlorodinitrobenzene, soil fumigation with, against wireworms, 630.

Chlorodryinus pallidus, parasite of Pyrilla in India, 243.

Chloroform, for extracting toxic principles of plants, 183, 206, 739.

chlorophanes, Opogona.

Chlorophorus herbsti, on Robinia and mulberry in Russian Union, 429.

Chlorophorus varius, on Robinia and mulberry in Russian Union, 429.

Chloropicrin, injected into trees against borers, 517; as a soil fumigant, 236, 630, 631; addition of other agents to, 236; causing injury to beet, 236; against pests of stored products, etc., 160, 326, 389, 514, 515, 654, 768: reducing germination of beet seed, 160; other fumigants compared with, 326, 389, 390.

Chlorops pumilionis (see C. tae-

niopus)

Chlorops taeniopus (on cereals), in Britain, 296; time of sowing and varieties of wheat in relation to, in Czechoslovakia, 658, 659; bionomics of, in Poland, 518; trap-strips against, 518.

Chlorosol, fumigation of stored

grain with, 125.

Chlorotettix unicolor, on narcissus in U.S.A., 288.

Chlorotettix viridius, on narcissus in U.S.A., 288.

Chocolate (see Confectionery).

Cholam (see Sorghum).

Chortoglyphus arcuatus, bionomics of, in stored cereals in Russian Union, 95, 96, 97.

Chortoicetes terminifera, bionomics and outbreaks of, in Australia, 111, 440, 449, 450, 494, 583, 584, 629, 676, 686, 703.

Chortophila flavopicta (see Hyle-

myia).

Chromoderus (see Cleonus).

Chrysanthemum, pests of, in Britain, 297, 439, 709; Aphis gossypii on, in Korea, 309; Tortrix postvittana on, in Tasmania, 8; pests of, in U.S.A., 20, 395, 396, 727; thrips as vectors of tomato spotted wilt in, 349; shading favouring pests of, 396.

Chrysanthemum cinerariaefolium, cultivation of, for pyrethrum

(q.v.), 412.

Chrysanthemum Leaf-miner Phytomyza atricornis).

Chrysobothris chrysostigma, conifers in Finland, 518.

Chrysobothris dentipes, in pine logs in Minnesota, 43.

Chrysobothris femorata, trees attacked by, in U.S.A., 136, 543, 587; bionomics of, 136, 137; measures against, 543.

Chrysocharis, parasite of Phytomyza ilicicola in Maryland, 541.

Chrysocharis laricinellae, parasite of Coleophora laricella in Austria, 660; establishment of, against C. laricella and Phyllotoma nemorata in U.S.A., 745.

Chrysomela decemlineata (see

Leptinotarsa).

Chrysomelids (sens. strict.), larval characters of, in Germany, 716. chrysomelina, Epilachna.

chrysomphali, Aphytis.

Chrysomphalus, characters distinguishing Aonidiella and, 226, 470.

Chrysomphalus agavis, on Juniperus spp. in Bermuda, **220, 626.** Chrysomphalus aonidum, auct. (see

C. ficus).
Chrysomphalus aurantii (see Aoni-

diella).

Chrysomphalus dictyospermi, on Citrus in California, 20; in Libya, 260; on cinnamon in Seychelles, 433; surface tension affecting toxicity of oil emulsions

to, 702.

Chrysomphalus ficus, measures against, in Florida, 728; in Palestine, 77; fungi infesting, on coconut and cinnamon in Seychelles, 433; parasite of, in West Indies, 41; on Citrus, 41, 77, 728.

Chrysomphalus obscurus, measures against, on fruit and nut trees in U.S.A., 18, 323, 587, 747.

Chrysomphalus personatus, systematic position of, 470.

Chrysopa, predacious on Aphids in Germany, 196.

Chrysopa flavostigma, predacious on Icerya seychellarum in Seychelles, 433.

Chrysopa perla, bionomics of, predacious on Aphids in Italy, 189; predacious on Tetranychus telarius in Bessarabia, 71.

Chrysopa phyllocroma, in Italy, 189. Chrysopa prasina, in Italy, 189.

Chrysopa septempunctata cognata, predacious on Eriosoma lanigerum in Japan, 771.

Chrysopa vulgaris, predacious on Tetranychus telarius in Bessarabia, 71; in Italy, 189.

chrysorrhoea, auct., Euproctis (see Nygmia phaeorrhoea).

chrysostigma, Chrysobothris.

Chrysothamnus, insects on, in U.S.A., 27, 247.

Cicada sexnotata (see Macrosteles).

Cicada, Seventeen-year (see Magicicada septendecim).

Cicadula, Zett. (see Thamnotettix). Cicadula, auct., generic attribution of species of, 78.

Cicadula divisa (see Macrosteles). Cicadula sexnotata (see Macrosteles). Cicadulina mbila, transmission of

streak disease of maize by, in

Tanganyika, 756.

Cicer arietinum, relation of Heliothis armigera to, in S. Africa, 364; little damaged by Bruchids in Burma, 7; not attacked by Pachytychius strumarius in Morocco, 641.

Cichorium intybus (see Chicory).

Cider, in bait for Cydia, 168.

Cienfuegosia hildebrandti, Diparopsis castanea on, in southern Africa, 199, 200, 365; distribution and ecology of, 199, 200.

Cigarette Beetle (see Lasioderma servicorne).

ciliatus, Dacus.

cilicrura, Phorbia (Hylemyia).

Cimex lectularius, toxicity of various fumigants to, 633.

Cinara, producing honey-dew in Central Europe, 459.

Cinara laricis, on Larix decidua in Czechoslovakia, 657.

Cinara piceicola, on Picea jezoensis, in Sakhalin, 452.

Cinara pichtae, type of injury to silver fir by, in Germany, 252, 253.

Cinara shinjii, n. n. (Lachnus fasciatus, Shinjii), on Pinus himekomatsu in Japan, 452.

Cinara thatcheri, sp. n., on Pinus ponderosa in Idaho, 369.

Cinara utahensis, sp. n., on Abies lasiocarpa in Utah, 369.

Cinara utahensis zoolathridi, subsp. n., on spruce in Oregon, 369.

Cinchona, Hypomeces squamosus on, in Indo-China, 186.

cincta, Ceratia (Aulacophora).

cincticeps, Nephotettix bipunctatus. cinctus, Anthonomus (see A. piri);

Cephus.

Cineraria, Phytomyza atricornis on, in Australia, 404.

Cineraria Leaf Miner (see Phytomyza atricornis).

cinerascens, Cnephalia (Gonia); Tachina (Larvaevora).

cinerea, Centeter; Zeuxia. cingulata, Herse; Rhagoletis.

cingulatorius, Exenterus (see E. abruptorius).

cingulatus, Dysdercus; Oncideres. Cinnamomum camphora (see Camphor).

Cinnamon, Coccids on, in Sev-

chelles, 433.

Cinquefoil (see Potentilla canadensis). Cionus fraxini (see Stereonychus). circumflexus, Myzus.

circumscriptus, Apanteles.

Cirphis, attempted introduction of Tachinid parasite of, into Hawaii, 586.

Cirphis leucosticha, on sugar-cane in Natal, 411.

Cirphis loreyi, parasite of, in S. Rhodesia, 89; on onion in

Tanganyika, 113.

Cirphis unipuncta, outbreaks of, in Canada and U.S.A., in 1937, 145, 441, 545, 580, 723; utilisation of ducks against, on rice in China, 498; on Juncus effusus decipiens in Japan, 672; in Manchuria, 296; bionomics of, 498, 723; parasites of, 296, 672, 723; measures against, 441. 498, 540, 723.

cirphis, Archytas.

Cirrospilus pictus var. arcuatus, parasite of Coleophora laricella in Austria, 660.

citrella, Phyllocnistis.

citri, Dialeurodes; Diaphorina; Paratetranychus; Prontaspis; Prospaltella; Pseudococcus.

Lepidosaphes citricola. (see beckii).

citrinus, Aspidiotiphagus.

citriperda, Brachycerus.

colocynthis, insecticidal Citrullus properties of, 360, 361.

Citrus, Lepidosaphes pinnaeformis on, in Portuguese E. Africa, 377; pests of, in S. Africa, 106, 176, 202, 377, 378, 455; mites on, in Argentina, 581; pests of, in Australia, 56, 85, 268, 335, 494, 502, 754, 755; Ceratitis capitata on, in Bermuda, 626; pests of, in Brazil, 82, 502, 607; pests of, in Ceylon, 359; Clania minuscula on, in China, 244; pests of, in Cyprus, 107, 706; pests of, in Formosa, 207, 263, 438, 674; pests of, in India, 55, 361, 765; pests of, in Indo-China, 185, 186 Coccids on, in Italy and Sicily, 70, 93, 713; pests of, in Japan, 53, 296, 438, 604; new Coccid on, in Java, 607; pests of, in Libya, 5, 260, 261; Phyllocnistis citrella on, in Loochoo Islands, 438: Papilio demodocus on, in Madagascar, 167; pests of, in Malta, 469; Anastrepha ludens on, in Mexico, 325; pests of, in New Zealand, 266; pests of, in Palestine, 77, 405, 406, 407, 514, 636, 637, 638; pests of, in Peru, 81, 82; pests of, in Russian Union, 305, 354, 481; Lepidosaphes beckii originally imported into Spain from Palestine on, 636; pests of, in Tanganyika, 35, 36; pests of, in U.S.A., 18, 20, 41, 131, 173, 219, 283, 286, 324, 325, 508, 574, 575, 650, 690, 722, 727, 728, 747, 748; pests of, in W. Indies, 10, 41, 268, 402, 405, 448, 490, 491, 751, 752; book on pests of, and their control, 722; susceptibility of species of, to C. capitata, 406, 407; Dacus cucurbitae experimentally developing in, 668; mites possibly transmitting disease of, 581; data on and dosage tables for HCN fumigation of, 70, 702, 703; not injured by methyl iodide fumigation, 575; oil emulsions causing injury to, 82, 355; absorption of selenium by, 131.

Citrus (Fruit), in bait for grasshoppers, 139.

Citrus poonensis, heat treatment of fruit of, against Dacus dorsalis in Formosa, 207.

Citrus trifoliata, Aspidiotus perniciosus experimentally developing on, 305.

Citrus Mussel Scale (see Lepidosaphes beckii).

Citrus Red Mite (see Paratetranychus citri).

Citrus Red Scale (see Aonidiella aurantii).

Citrus Whitefly (see Dialeurodes

citri). civilis, Tachina (Larvaevora).

Cladosporium, associated Meromyza americana in Kansas, 123.

Clania crameri, parasite of, in Madras, 499.

Clania minuscula, bionomics of, on Citrus in China, 244, 245.

Clania pryeri, on banana in Formosa, 263.

claripalpis, Theresia (Paratheresia). clavatus, Pilophorus.

Clavigerus smithiae (see Pterocomma).

clavipes, Otiorrhynchus.

Clay, as a carrier for dusts, 15, 17, 133, 550, 647, 648, 725; spray formula containing, 350.

Clear-winged Grasshopper (see Camnula pellucida).

clematidis, Catagmatus.

Clematis, new weevil on, in Sumatra,

Clemelis bullata (see Zenillia).

Clensel, in baits for Trypetids, 58, 108, 407, 697, 699, 700; ineffective in bait-spray, 108; onion seedlings immersed against Thrips tabaci, 626.

Cleonus alternans, not injurious to

beet in Italy, 161.

Cleonus fasciatus (on beet), not injurious in Italy, 161; in Poland, 256.

Cleonus foveicollis, on beet in

Poland, 256.

Cleonus luigionii, on beet in Italy, 161, 527.

Cleonus mendicus, on beet in Italy, 161, 527; parasites and control of, 527.

Cleonus morbillosus, not attacking beet in Italy, 161.

Cleonus pedestris, not injurious to

beet in Italy, 161.

Cleonus piger (on beet), not injurious in Italy, 161; in Poland, 256.

Cleonus punctiventris (on beet), in Poland, 256; in Russian Union, 236, 237, 356, 473; bionomics of, 237, 256; other crops not attractive to, 356, 473; measures against, 236, 256.

Cleonus tigrinus, not attacking beet in Italy or Poland, 161, 256.

Cleora acaciaria, on Schleichera trijuga in Ceylon, 359.

clerkella, Lyonetia.

Clidemia hirta, utilisation of Liothrips urichi against, in Fiji and Solomon Is., 59, 703.

Clinodiplosis, on hollyhock in Br. Isles, **33.**

Clinodiplosis mosellana (see Sitodiplosis).

Clinodiplosis oleisuga, on olive in

Spain, **700**.

Clivina impressifrons (on maize), measures against, in Australia, 352; probably in New Zealand, 361.

Closterocerus insignis, parasite of Oscinis theae in India, 149.

Closterocerus tricinctus, parasite of Phytomyza ilicicola in Maryland, 541.

Clothes Moths, leaflet on, in U.S.A., 367. (See Borkhausenia pseudospretella and Tineola biselliella.)

Clove Tree, Cerambycids on, in Netherlands Indies, 295; termite damaging, in Tanganyika, 642.

Clover, pests of, in Australia, 209, 582; Apion on, in Britain, 527; Coleophora spissicornis on, in Denmark, 155; Otiorrhynchus ligustici on, in Germany, 520; Myzus persicae on, in Japan, 265; Lymantria dispar on, in Jugoslavia, 759; Cirphis unipuncta on, in Nova Scotia, 441; pests of, in Russian Union, 75, 307, 308, 425, 470, 471, 473, 474, 475; pests of, in U.S.A., 12, 15, 126, 136, 287, 444, 723, 747; Sitona lineata on, in Vancouver, 293; review of literature on pests of, 386; insects and diseases of, 12, 265, 386; in crop rotation against pests, 78, 125; in bait for weevils, 75.

Clover, Alsike (see Trifolium

hybridum).

Clover Case-bearer (see Coleophora spissicornis).

Clover Weevils (see Apion aestivum

and A. flavipes).

Clysia ambiguella (Vine Moth), parasite of, in France, 415; in Germany, 50, 154, 419, 521, 661; in Switzerland, 299; notice of lists of food-plants of, 4: bionomics of, 415, 419, 661; tests of baits for, 50, 154; possible use of bands against, 419; value of spraying against, 521.

Cnaphalodes (see Chermes).

Cnephalia cinerascens, parasite of Prodenia litura in Formosa, 207. Cnephasia conspersana, on iris in Scilly Isles, 527.

coarctata, Scotinophara. coccidivorus, Anagyrus.

coccinea, Graphocephala; Pyrochroa. Coccinella axiridis, predacious on Eriosoma lanigerum in Japan, 771. Coccinella bipunctata (see Adalia).

Coccinella septempunctata, predacious on Aphids and Lepidoptera in Ukraine, 353, 354.

Coccinella trifasciata, fed on frozen Aphids in Ohio, 650.

Coccinellids, colour patterns indicating feeding habits of, in E. Africa, 611.

Coccobacillus, mite possibly connected with, infecting Schistocerca paranensis in Argentina, 147.

Coccophagus gurneyi, explanation of immunity of Pseudococcus citri from, 287.

Coccotrypes, characters of, 529.

Cocculus suberosus, tests of insecticidal properties of, 361.

Coccus hesperidum, measures against, on Citrus in Cyprus, 706, 707; in New Zealand, 753; in Peru, 82; on Citrus, 82, 706, 707; Coccinellids predacious on, 82, 753.

Coccus mangiferae, fungus infesting, on mango in Florida, 587; on cinnamon in Seychelles, 433.

Coccus pseudomagnoliarum, measures against, on Citrus in Cali-

fornia, 575.

Coccus viridis (on coffee), measures against, in Indo-China, 185; toxicity of various plant extracts to, in Mysore, 361; in Réunion, 706; on guava, 361.

Cochin, coconut pests in, 452.

Cochin China (see Indo-China).

cochleariae, Phaedon.

cockerelli, Horismenus; Paratrioza. Cockroaches, tests of insecticides

on, **35, 183, 225, 439.**

Coconut, scarcity of Coccids on, in E. Africa, 611; pests of, in Ceylon, 359, 360, 434; pests of, in Cochin, 452; pests of, in Fiji, 211, 212, 334, 497; Homaledra sabalella on, in Florida, 22; pests of, in Br. Guiana, 369, 491; Hypomeces squamosus on, in Indo-China, 186; pests of, in Netherlands Indies, 337, 674; Hispids on, in Japanese Mandated Islands, 336; Oryctes rhinoceros on, in Malaya, 434; Coccids on, in Mauritius, 66, 176, 204; pests of, in New Guinea Territory, 435, 495; Artona catoxantha on, in Philippines, 675; Aspidiotus destructor on, in Santo Domingo, 10; pests of, in Seychelles, 433, 611; Pyroderces paroditis on, in Solomon Islands, 497; pests of, in W. Indies, 43, 121, 268.

Coconut Spike Moth (see *Tirathaba trichogramma*).

Coconut Zygaenid (see Artona catoxantha).

Coconut-oil Soap (see Soap, Coconut-

Cocos australis, Homaledra sabalella on, in Florida, 22.

Cocos nucifera (see Coconut).

Cocytodes coerulea, in Japan, 266, 672; measures against, on ramie, 266; parasite of, 672.

Cod-Ban, 507.

Codiaeum, Orthezia praelonga on, in Grenada, 752.

Codling Moth (see Cydia pomonella). Coelinius (Coelinidea) spp., parasites of Meromyza americana in Kansas, 123.

Coeliodes fuliginosus, on poppy in Czechoslovakia, 657.

Coelognathus dimidiatus (see Tyrophagus putrescentiae).

Coeloides scolytivorus, parasite of Hylesinus aculeatus in New Jersey, 394.

Coelophora inaequalis, predacious on Aphis gossypii in Korea, 309. coerulea, Agelastica; Cocytodes.

coeruleiviridis, Lucilia. coeruleocephala, Diloba.

coeruleus, Aenasius; Curinus; Rhynchites.

Coffea (see Coffee).

coffeae, Ceratitis (Trivhithrum); Pseudococcus; Rhembastus; Saissetia; Xyleborus; Zeuzera.

coffearia, Homona.

Coffee, pests of, in Br. E. Africa, 68, 113, 151, 177-180, 372, 405, 592, 610; pests of, in Brazil, 319, Stephanoderes hampei on, in Ceylon, 360; records of insects on, in Colombia, 533; Ceratitis capitata on, in Hawaii, 84; pests of, in Indo-China, 185, 261; Coccids on, in Java, 182, 606, 607; Cephonodes hylas on, in Madagascar, 67; pests of, in Mysore, 55, 361; Ferrisiana virgata on, in New Guinea Territory, 495; pests of, in Réunion, 706; pests of, in N. Rhodesia, 500; pests of, in Sierra Leone, 239; pests of, in West Indies, 40, 222, 223, 490; Antestia spp. associated with Nematospora on, 178.

Coffee (Stored), Stephanoderes hampei and its control in, in Ceylon, 359.

Coffee Stem Borer (see Xylotrechus quadripes).

coffeella, Leucoptera.

cognata, Chrysopa septempunctata.

cognatus, Epicaerus.

Cola, pests of, in W. Africa, 239, 240, 469; treatment of nuts of, against Balanogastris kolae, 239. colae, Ceratitis.

Cold Storage, uses of, against insect pests, 229, 238, 269, 325, 363, 599; Aphids kept in, for rearing Coccinellids, 650; effects of, on eggs of Ephestia and Trichogramma, 670.

Cold-hardiness, studies on, in in-

sects, 257, 274, 304.

Coelophora caryaefoliella, on pecan in Florida, 587.

Coelophora hemerobiella, in orchards in Czechoslovakia, 657.

Coleophora laricella (Larch Casebearer), in Austria, 659, 660; in U.S.A., 543, 745; parasites and biological control of, 659, 660, 745; other measures against, 543.

Coleophora nigricella, on fruit-trees

in Britain, 612.

Coleophora spissicornis, bionomics of, on clover in Denmark, 155.

Coleoptera, list of, attacking stored products etc. in Germany, 194; key to, injurious to forests in Central Europe, 312.

Coleus, Coccids on, 377, 598.

Colias electo, bionomics and control of, on lucerne in S. Africa, 151, 152.

Colias hyale poliographus, on leguminous crops in Japan, 673.

collaris, Amicroplus.

Collops bipunctatus, predacious on Chlorochroa sayi in Montana, 44.

Collyria calcitrator, analysis of superparasitism of Cephus pygmaeus by, in Britain, 102.

Coloborhombus fasciatipennis, bionomics and control of, on clove trees etc. in Netherlands Indies, 295.

Coloborhombus hemipterus, on clove trees in Netherlands Indies,

Colocasia, Hepialid attacking roots of, in Formosa, 517; Megamelus proserpina on, in Hawaii and

Philippines, 212.

Colombia, insects on coffee in, 533; Sacadodes pyralis on cotton and its natural enemies in, 403; new weevil in, 405; new Encrytid introduced into Hawaii from, against Pseudococcus brevipes, 212.

colombiensis, Aenasius.

Colophony, emulsifier for pyrethrum containing, 412; in banding adhesives, 480.

Coloradia pandora, on Pinus contorta in Colorado, 580.

Colorado, forest pests in, 580; Stretchia plusiaeformis on gooseberry in, 281; Hypera variabilis in, 326; vegetable pests in, 28, 327, 643; parasite of Hyphantria cunea in, 291.

Colorado Potato Beetle (see Lepti-

notarsa decemlineata).

Colours, reactions of insects to, 506, 573, 732.

combinata, Geomyza.

Combretum, Helopeltis bergrothi on, in Sudan, 513.

comes, Erythroneura.

communis, Lygus.

Comocritis pievia, bionomics and control of, on tea in Assam, 601, 602.

Comperiella bifasciata, utilisation of, against Aspidiotus destructor in Netherlands Indies, 337.

Comperiella unifasciata, attempted introduction of, into Mauritius against Aspidiotus destructor, 176.

complanatus, Laemostenus.

complanella, Tischeria.

Compsilura concinnata, parasite of Pieris spp. in Britain, 342; parasite of Lymantria dispar in Jugoslavia, 760; hosts of, in U.S.A., 589, 590.

compta, Psilopa nitidula.

comptana, Ancylis.

concinna, Chaetocnema (Tlanoma).

concinnata, Compsilura.

concinnus, Stigmatrachelus; Wes maëlius.

Confectionery, insects infesting, in Germany, 460, 461.

conformis, Leis.

Confused Flour Beetle (see Tribolium confusum).

confusum, Tribolium.

confusus, Ips.

Congo, Belgian, Agrilus ghesquierei on Albizzia in, 118; cotton pests in, 315; locusts in, 679, 687; summary of plant quarantines in, 134.

congregatus, Apanteles.

coniferana, Cydia (Laspeyresia).

conigerum, Sinoxylon.

conimbrigensis, Asterolecanium.

Coniopteryx pusana, bionomics of, predacious on Pyrilla in India, 241.

conjugalis, Signiphora. conjugella, Argyresthia.

Connecticut, pests of forest and shade trees in, 14, 15, 17, 18, 546, 547, 723, 725; maize pests in,

14, 723, 726, 730, 745; miscellaneous pests, 14-18, 246, 247, 722-726; pests of timber in, 17, 218, 724, 726, tobacco pests in. 132, 726; vegetable pests in, 14, 17, 722, 725, 731; beneficial insects and biological control in, 14, 15, 16, 723, 724, 725; nematode parasite of grasshoppers in, 124; Conopid parasite of honey bees in, 133; new Coccid in, 226.

connexa, Diabrotica; Eriopis. Conoderus auritus, predacious on Tyloderma fragariae in Kentucky,

737.

Conoderus exsul, on potato Hawaii, 214; introduced into California, 589.

Conopia (see Aegeria).

Conorrhynchus (see Cleonus).

Conotrachelus anaglypticus, Ceratostomella ulmi cultured from, on elm in New Jersey, 325. Conotrachelus licaniae, sp. n., on

Licania rigida in Brazil, 106, 115;

measures against, 115.

Conotrachelus nenuphar, in Ontario, 473; measures against, in U.S.A., 15, 16, 323, 333, 443, 444, 489, 647, 653, 723, 724, 734; on apple, 15, 333, 647, 723; on cherry, 489; on peach, 16, 443, 444, 653, 724, 734.

Convadtina acroleuca, parasites of, in Nigeria, 492.

consocialis, Oechalia.

conspersana, Cnephasia.

constricta, Agallia.

Contarinia, natural enemies of, on wheat in Japan, 670.

Contarinia loti, bionomics and control of, on lucerne in Italy, 189, 190.

Contarinia medicaginis (on lucerne), in Hungary, 527; bionomics of, in Sweden, 232; parasites of, 232, 527.

Contarinia nasturtii, bionomics and control of, on crucifers in Holland,

466, 467.

Contarinia pyrivora (on pear), in Czechoslovakia, 657; in Italy, 636; in New York, 130, 131; measures against, 130, 131, 636.

Contarinia soya (see Profettiella). Contarinia torquens (see C. nastur-

tii).

tritici (Wheat Gall-Contarinia midge), in Germany, 619; in Sweden, 154, 230, 231, 620; bionomics of, 230 ; natural enemies of, 155, 231, 620.

contariniae, Inostemma.

Contheyla rotunda, on coconut in Cochin. 452.

contrahens, Drosicha.

contraria, Ochrogaster.

convergens, Hippodamia.

Convergent Ladybeetle (see Hippodamia convergens).

Convolvulus, Tetranychus on, in Russian Union, 257, 471.

Cook Islands (see Rarotonga).

cooki, Cremastus.

Copa occidentalis, on cotton in Belgian Congo, 315.

cophta, Gryllotalpa gryllotalpa.

Copidosoma geniculatum, imported into U.S.A., against Rhyacionia buoliana, 329.

copiosella, Ocnerostoma (see O. pini-

ariella).

Coposil (see Copper Ammonium Silicate).

Copper Acetoarsenite, toxicity of homologues of, to Tribolium confusum, 30. (See Paris Green.)

Copper Ammonium Silicate, spraying with, against Lepidoptera on tomato, 752; in mixed sprays, 554, 752.

Copper Carbonate, uses of, as an insecticide, 125, 248, 609, 713.

Copper Cyanide, mixed with stored wheat, against pests, 125.

Copper Oxide, as an insecticide, 248. Copper Phosphate, as an insecticide, 125, 609.

Copper Sulphate, toxicity of, to Dacus cucurbitae, 182; and lime, dusting with, 333, 443, 564, 565, 566; and calcium arsenate, 333, 443; and cubé, 566; and sulphur, 565; spray formula containing, 350. (See Bordeaux Mixture.)

Copra, insects associated with, in Malaya, 31; Ephestia cautella in, in New Guinea Territory, 435.

Coptocercus rubripes, on Eucalyptus in New Zealand and Australia, 754.

Coptotermes, measures against, damaging tea in India, 149.

Coptotermes amanii, damaging trees and timber in Tanganyika, 113, 642.

Coptotermes curvignathus, bionomics and control of, in Indo-China, 186, 187, 188.

Corchorus olitorius, Platyedra gossypiella on, in Sudan, 511.

Corcyra cephalonica, in copra and stored rice in Malaya, 31; rearing of Trichogramma spp. in eggs of, in Mauritius and India, 176, 409. Cordyceps fuliginosa (see Hirsutella

exoleta).

Cordyceps militaris, identity of, infesting Lepidoptera, 152.

Cordyceps pistillariaeformis, infesting Lecanium in Czechoslovakia, 659.

coriaceus, Eriococcus.

Coriandrum sativum, tests of oil of, against Aphids and mites, 249.

Corigetus dejeani, on kapok in Indo-China, 186.

Corioxenos antestiae (parasite of Antestia), bionomics of, in E. Africa, 179.

Cork Oak (see Quercus ilex var. suber).

Corks, Borkhausenia pseudospretella infesting, 413.

Corn Borer, European (see Py-rausta nubilalis).

Corn Earworm (see Heliothis armigera).

Corn Root Aphis (see Anuraphis maidiradicis).

Corn Rootworm, Southern (see Diabrotica duodecimpunctata).

Corn Stalk Borer, Southern (see Diatraea crambidoides).

Corn Worm, Pink (see Pyroderces rileyi).

Cornflower, Aphid on, in Britain, 350.

corni, Lecanium (Eulecanium).

cornifrons, Hybalus.

Cornus, Lecanium corni on, in Germany, 251; insects on, in U.S.A., 288, 546.

Cornus florida, Prionus laticollis on, in New York, 547.

Cornus mas, Rhagoletis cerasi not observed in fruits of, in Switzerland, 156.

cornutus, Gnathocerus.

Corrosive Sublimate (see Mercury Bichloride).

corrupta, Epilachna (see E. varivestis).

corticalis, Xenylla.

coryli, Emphytus; Phylus.

Corylus (Filbert, Hazel), pests of, in Germany, 69, 234, 560; pests of, in U.S.A., 398, 547.

Corymbites aeneus, ecology of, in Russian Union, 425.

Corymbites latus, on wheat and grasses in Central Asia, 76; ecology of, in Russian Union, 425, 429.

Corymbites sjaelandicus, ecology of, in Russia, 425.

Corymbites spretus, ecology of, in Siberia, 425.

Coryna apicicornis, on Cajanus cajan in Tanganyika, 642.

Corythuca cydoniae, on Cotoneaster and quince in Ohio, 399; tests of insecticides against, 399.

Cosmoglyphus, subgenus of Eberhardia (q.v.), 95.

Cosmolyce boetica (on pulse crops), in Cyprus, 108; measures against, in Hawaii, 752.

Cosmophila erosa, species recorded as, on cotton in Madagascar and Nyasaland, 67, 114.

Cosmophila flava, on cotton in Madagascar and Belgian Congo, 67, 315; measures against, on Hibiscus sabdariffa in Netherlands Indies, 358; bionomics of, 67, 358.

Cosmopolites sordidus (on banana), in Ceylon, 360; in Formosa, 263; introduction of Plaesius javanus into Jamaica against, 145; in New South Wales, 210; in Porto Rico, 223; measures against, 210, 223.

cossis, Hylecoetus.

Cossus cossus, measures against, on Citrus and Cynara in Sicily, 93.

Costa Rica, summary of plant quarantines in, 134.

Costalimaspis, gen. nov., new species of, in Brazil, 367.

costipunctata, Hemithea. cosyra, Ceratitis.

Cotoneaster, pests of, in U.S.A., 399, 547.

Cotton (Gossypium), Diparopsis castanea on, in Portuguese E. Africa, 199; pests of, in S. Africa, 177, 199, 200, 364, 365, 366; pests of, in Argentina, 362, 503, 704; pests of, in Brazil, 115, 268, 333, 405, 502-504, 505, 689, 704; Pempheres affinis on, in Burma, 409; pests of, in China, 245, 304, 498; Sacadodes pyralis on, in Colombia, 403; pests of, in Belgian Congo, 315; Platyedra gossypiella on, in Cyprus, 108: pests of, in Egypt, 457, 458, 685, 686; pests of, in Formosa, 309; pests of, in India, 41, 55, 89, 180, 204, 242, 261, 408, 409, 532, 666 pests of, in Indo-China, 186 pests of, in Java, 106, 765, 766;

Aphis gossypii on, in Korea, 309, 674; pests of, in Madagascar, 67; pests of, in Mexico, 274, 275; pests of, in Nigeria, 666, 667; pests of, in Nyasaland, 113, 114, 177, 367, 372; Platyedra vilella on, in Persia, 249; pests of, in Peru, 533, 626; pests of, in Queensland, 494; pests of, in N. Rhodesia, 177, 366, 500; pests of, in S. Rhodesia, 366: pests of, in Russian Union, 238, 249, 250, 353, 354, 470, 471, 472, 530; pests of, in Sudan, 511, 512, 513; pests of, in Tangan-yika, 112, 113, 610, 642; pests of, in Uganda, 83, 372–375, 376; pests of, in U.S.A., 19, 41, 129, 132, 133, 134, 136, 140, 141, 142, 229, 274, 275, 276, 277, 278, 279, 327, 328, 329, 384, 385, 398, 580, 644, 645, 686, 691, 748; textbook on, and its pests in U.S.A., 722; pests of, in W. Indies, 41, 42, 65, 83, 222, 445, 446, 752; distribution and ecology of wild, in relation to Diparopsis castanea in Africa, 199, 200; time of 471, 503, 504, 505, 644, 666, 667, 766; Bemisia goldingi transmitting leaf-curl of, 666, 667; insects associated with boll rots in, 249, 375, 376, 457, 513; factors affecting attacks of insects and fungi on, 316, 532, 666; analysis of nectar of, as food of Heliothis 197; insecticides armigera, causing injury to, 249, 472, 705. Cotton Aphis (see Aphis gossypii).

Cotton Boll Weevil (see Anthonomus grandis). Cotton Bollworm (see Heliothis

armigera).

Cotton Bollworm, Pink (see Platy-edra gossypiella).

Cotton Bollworm, Red (see Diparopsis castanea).

Cotton Borer, Brazilian (see Gasterocercodes brasiliensis).

terocercodes brasiliensis).
Cotton Flea-hopper (see Psallus

seriatus).
Cotton Leaf-roller (see Sylepta

derogata). Cotton Leaf Worm (see Alabama

argillacea).

Cotton Lint, legislation dealing with, against *Platyedra gossy-piella* in U.S.A., **133, 134.** Cotton Stainers (see *Dysdercus*).

Cottonseed, prescribed measures against pests in, in U.S.A., 134, 229, 384; other measures against *Platyedra gossypiella* in, 408, 446; products of, in baits for insects, 139, 509, 574, 705.

Cottonseed Oil (see Oil, Cottonseed). Cottony Cushion Scale (see *Icerya*

purchasi).

Couch Grass (see Agropyrum repens). Cowpea Weevil (see Chalcodermus

aeneus).

Cowpeas, Bruchids on, in Burma, 7; Cosmolyce boetica on, in Cyprus, 108; Noctuids on, in Nyasaland, 114; Dacus cucurbitae on, in Philippines, 181; new weevil on, in S. Rhodesia, 106; Coleoptera on, in U.S.A., 572, 649, 691; calcium arsenate injurious to, 328, 691.

Cowpeas (Stored), pests of, and their control 7, 11, 642, 649.

Crabgrass (see Digitaria sanguina-lis).

Crabro davidsoni, sp. n., predacious on leafhoppers in Ohio, 369.

crambidoides, Diatraea.
Crambus hortuellus, on cramberry in Washington, 172.

crameri, Clania.

Cranberry, Tortrix rosana on, in Holland, 299; pests of, in U.S.A., 171, 172, 508, 562, 563, 645, 646; problem of spray residues on fruit of, 646.

Cranberry Fruitworm (see Mineola

vaccinii).

Cranberry Girdler (see Crambus hortuellus).

Cranberry Tipworm (see Dasyneura vaccinii).

Cranberry Weevil (see Anthonomus musculus).

crassicornis, Gallobelicus (see Engytatus tenuis).

crassifemur, Eulimneria. crassitarsis, Phytodietus.

crataegi, Aporia; Eriosoma; Glossonotus.

Crataegus (Hawthorn), identity of woolly aphis on, in N. America, 484, 485; Aphids on, in Germany, 196; pests of, in Holland, 183, 465; Tortricid on, in Ontario, 509; Aglaope infausta on, in Spain, 697; experimentally attacked by Otiorrhynchus hirticornis, 345.

Cratopus aurostriatus, on Calophyllum inophyllum in Seychelles,

433.

Cremastus biguttulus (see C. flavo-orbitalis).

Cremastus cooki, parasite of Ancylis comptana in Ohio, 735.

Cremastus flavo-orbitalis, parasite of Argyroploce illepida in Hawaii, 497; parasite of Chilo simplex in Japan, 54; establishment of, against Pyrausta nubilalis in U.S.A., 730.

Cremastus forbesi, hosts of, in Ontario, 510.

Cremastus hymeniae (see C. flavoorbitalis).

Cremastus interruptor, establishment of, against Rhyacionia buoliana in U.S.A., 745, 746.

Cremastus minor, hosts of, in Ontario, 510.

Creontiades pallidus, on cotton, in Sudan, 513.

Creosote, as a barrier for *Blissus* leucopterus and clover mites, **63**, **85**; methods of treating timber with, against pests, **514**, **707**, **748**; other uses of, against insects, **219**, **604**, **645**.

Creosote Tar Oil, **270.** (See Tar Distillates.)

Crepidodera (Chalcoides) helxines, on peach in Connecticut, 17.

Crepidodera rufipes, on apple and mallow in Britain, 612.

Cresylic Acid, and oils, spraying with, against Aphids, Coccids and mites, 16, 738.

Crete, Eumarchalia gennadii on carob in, 71.

cretica, Sesamia.

Cricket, House (see Gryllulus domesticus).

Cricket, Mormon (see *Anabrus* simplex).

crini, Brithys. crinita, Sitona.

Crioceris asparagi, Pentatomids reared on, in Ohio, 13, 14.

Crioceris orientalis, on asparagus in Japan, 497.

Crioceris quatuordecimpunctata var. sibirica, on asparagus in Japan, 497.

cristata, Homalopoda.

crocale, Catopsilia.

Cronartium ribicola, 625.

Crossotarsus wallacei on poisone

Crossotarsus wallacei, on poisoned and girdled Styrax in Malaya, 453.

Crotalaria, Xyleborus fornicatus on, in India, 149; thrips on, in Sudan, 513.

Crotalaria anagyroides, Acrosternum spicatum on, in Seychelles, 433.

Crotalaria juncea (Sunn Hemp), Bemisia gossypiperda and disease of, in India, 89; new weevil on, in S. Rhodesia, 106; relation of Bemisia gossypiperda to virus diseases of, in India, 242.

Crotalaria striata, Coccinellid favoured by, in Florida, 727.

Croton, Lepidosaphes pinnaeformis on, in S. Africa, 377.

crotonis, Pseudococcus (see P. lila-cinus).

cruciata, Austroicetes.

crucifer, Spaniopterus.

cruciferae, Phyllotreta. cruciferarum. Plutella (se

cruciferarum, Plutella (see P. maculipennis).

cruentatus, Rhipiphorothrips.

Cryolite (Sodium Fluoaluminate), in baits, 387; dusting with, 65, 129, 145, 203, 327, 328, 388, 489, 526, 643, 845, 654, 736; adhesive for dusts of, 388; carriers for, 327, 328, 489, 736; spraying with, 16, 25, 62, 79, 100, 145, 327, 399, 401, 443, 488, 547, 724, 735, 736; adhesives and spreaders for, 16, 25, 547, 724, 735, 736; tests of toxicity of, to insects, 216, 387, 388, 572, 654; formulae containing, 16, 25, 327, 387, 443, 547, 724, 735, 736; and barium fluosilicate, 736; and bentonitesulphur, 25; and other forms of sulphur, 25, 129, 443, 724; and lime, 724; and oils, 62, 401, 488; inverted spray mixtures of oil and, 174, 488; factors affecting amount of deposit of, on foliage, 203; problems of residues of, on sprayed fruits and vegetables, 25, 62, 216, 327, 401; forms of, 16, 65, 100, 399, 735, 736; other insecticides compared with, 25, 62, 79, 327, 387, 401, 547, 645, 735, 736.

Cryphalus abietis, on pine in Estonia, 462; on spruce in Sweden, 634.

Cryphalus saltuarius, on spruce in Sweden, 634.

Cryptocephalus incertus, measures against, on cranberry, etc., in Massachusetts, **645**, **646**.

Cryptognatha nodiceps, introduced into Santo Domingo against Aspidiotus destructor, 10.

Cryptolaemus montrouzieri (used against Coccids), establishment

of, in Netherlands Indies, 337; bionomics of, in Massachusetts, 648; attempted establishment of, in New Zealand, 753; in Spain, 701.

Cryptolucilia caesarion, in drying

fruit in California, 228.

Cryptomeria japonica, Cerambycids

on, in Japan, 516.

Cryptorrhynchus lapathi, bionomics and control of, on willow in Czechoslovakia, 658.

Cryptostemma calendulaceum, mites on, in Australia, 85, 151.

Cryptotaenia japonica, Papilio machaon on, in Japan, 673. Cryptotermes (see Calotermes).

Cryptotympana japonensis, measures against, on pear in Japan, 604.

cryptus, Pseudococcus.

Ctenopseustis obliquana, bionomics of, in New Zealand, 57, 58.

ctenopterus, Systoechus.

Cuba, Hymenopterous parasites in, 171, 330; summary of plant quarantines in, 385; Chilocorus cacti introduced into Domingo from, against Aspidiotus

destructor, 10. Cubé, dusting with, 17, 129, 132, 279, 285, 327, 333, 371, 399, 443, 566, 645, 647, 648, 723, 735, 740, 741; carriers for, 133, 399, 566, 647, 648, 735, 740; conditioning agents for dusts of, 285, 327, 740; spraying with, 128, 285, 327, 443, 489, 542, 543, 547, 548, 553, 566, 646, 647, 708, 709, 723, 725, 755; spreaders, wetters and adhesives for, 285, 489, 543, 547, 548, 646, 647, 725, 755; tests of toxicity of, to insects, 17, 133, 279, 333, 371, 543, 566; comparison of water suspensions and extracts of, 648; formulae containing, 128, 489, 542, 547, 548, 646, 647, 755; and copper compounds, 566; and nicotine, 740, 741; and oil emulsions, 708, 709; and pyrethrum, 740; and sulphur, 489, 566; and thiocyanate, 740, 741; rotenone contents of, 17, 120, 128, 129, 132, 279, 319, 333, 371, 399, 489, 543, 548, 553, 566, 645, 646, 647, 651, 708, 723, 735, 739, 740, 741; chemistry and evaluation of, 120, 331, 574, 651, 739; effect of attack of Dinoderus bifoveolatus on rotenone content of stored roots of, 396; identity of plant known as, in S. America, 319; other insecticides compared

with, 132, 279, 547, 553, 645, 647, 651, 723, 735, 740.

cucullata, Rondania.

Cucumber, little attacked by Heliothis armigera in S. Africa, 364; pests of, in Britain, 615, 712 Dacus cucurbitae on, in Formosa, 206; D. cucurbitae on, in Hawaii, 325; Oxytelus sculptus on, in Holland, 299; Aphis gossypii on, in Korea, 309; Camptocladius macleavi on, in New South Wales. 209; pests of, in U.S.A., 17, 269. 333, 399, 553, 704; Aphids transmitting virus diseases of, 615; insecticides causing injury to, 712.

Cucumber Beetle, Striped (see Diabrotica melanocephala).

cucumeris, Epitrix.

cucurbitae, Dacus (Chaetodacus). Cumbu (see Pennisetum typhoideum). cunea, Hyphantria.

cunicularius, Hylastes.

Cunninghamella, infesting Schistocerca paranensis in Argentina, 147. Cuphocera hova, parasite of Cirphis loreyi in S. Rhodesia, 89.

Cuphocera varia, parasite of Spodoptera mauritia in India, 241.

cupira, Melipona (Trigona) testacea. cuprea, Amonala; Kophene.

cupreoviridis, Earias.

Cuprous Cyanide, against Phthorimaea lycopersicella, 327.

cupulatus, Platypus.

Curculio, in chestnuts for export from Italy, 384.

Curculio camelliae, bionomics of, on Camellia in Japan, 438.

Curculio caryae, on pecan in U.S.A., 587, 690; measures against, 690.

Curculio, Plum (see Conotrachelus nenuphar).

curculionis, Bathyplectes.

curculionum, Eurytoma.

Curinus coeruleus (predacious on Pseudococcus nipae), in Mexico, introduced into Hawaii, 586; 586.

Curly-top of Beet, experiments with insects and, in California, 445.

Currant, new mite on, in Central Asia, 530; Lygus pabulinus on, in Britain, 297; pests of, in Finland, 201; Capsids on, in Germany, 234; Trypetids on, in U.S.A., 60; HCN unsuitable fumigant for fruit of, 307.

Currant, Black, Dasyneura tetensi on, in Britain, 612; Tortrix rosaceana on, in Utah, 141; effect of oil emulsions on, 708.

Currant Fly (see Epochra canaden-

curvignathus, Coptotermes. curvipes, Anoplocnemis. cushmani, Microbracon. cuspidatae, Pseudococcus.

cuthbertsoni, Analeurops; Hvalomya.

Cutworm, Pale Western (see Porosagrotis orthogonia).

Cutworm, Red-backed (see Euxoa

ochrogaster).

Cutworms, baits for, 160, 236, 441, 498. 540: other measures against, 236, 498, 723, 751.

cyanea, Scutellista. cyaneus, Sirex.

Cyanogas, 86, 241, 703, 711.

Cyclamen, Thysanoptera on, 483, 673, 712.

Cyclohexylamine, derivative of. against Paratetranychus pilosus,

Cycloneda sanguinea, predacious on Sipha flava in Porto Rico, 39, 289; not attacking Diatraea saccharalis, 289.

Cydia coniferana, on Pinus cembra in Austria, 664.

Cydia dorsana, C. nigricana recorded as, in Russia, 428.

Cydia funebrana (on plum), in Czechoslovakia, 657; bionomics and control of, in Switzerland, 99, 100.

Cydia glycinivorella (Soy Bean Pod Borer), in Japan, 310; bionomics of, in Manchuria, 54, 310, 452, 606.

Cydia latiferreana, in acorns in Ontario, 510; on Corylus in Oregon, 398; parasites of, 398, 510.

Cydia molesta (Oriental Peach Moth), in Australia, 169, 448, 450; legislation against, in New South Wales, 608; in Canada, 509, 510, 535; in Italy, 154, 163, 594, 714, 758; in Japan, 604; in Switzerland, 163; in U.S.A., 16, 125, 135, 154, 270, 399, 443, 444, 448, 450, 506, 544, 644, 647, 653, 724, 725, 735; in Uruguay, 582; on pear, 169, 594, 604; on quince, 647; hibernation of, 594, 714; artificial interruption of diapause in, 758; parasites and biological control of, 16, 125, 154, 270, 399, 448, 450, 509, 510, 535, 544, 582, 647, 653, 724, 725, 735; other measures against, 125, 135, 443, 444, 448, 506, 544, 594, 608, 653, 714, 725, 758.

Cydia nigricana (Pea Moth), bionomics of, in Ouebec and Russia, 60, 428, 623; susceptibility of

varieties of peas to, 428.

Cydia pomonella (Codling Moth), in Australia, 149, 168, 449, 450, 510; legislation against, in New South Wales, 608; in Brazil, 320; in Britain, 612, 613, 689; in Canada, 59, 535, 541; in Cyprus, 108, 259, 260; in Czechoslovakia, 657, 658; in France, 192, 259, 464, 525; in Hungary, 254; in India, 340, 765; Italy, 593, 712, 713, 758; Morocco, 341; in New Zealand, 137, 174, 272, 291, 323, 330, 389, 401, 442, 444, 450, 488, 507, 544, 554, 555, 557, 588, 652, 653, 647, 723, 731, 732, 733, 745, 747; on chestnut, 658; on peach, 19; on pear, 19, 99, 168, 192, 259. 340, 389, 500, 525, 658, 758, 765; on quince, 500; on walnut, 192, 500, 658; bionomics of, 61, 99, 168, 192, 259, 341, 401, 478, 479, 500, 510, 593, 612; effect of temperature on oviposition of, 731, 732; birds destroying, 291; parasites and biological control of, 30, 65, 291, 323, 330, 356, 450, 464, 535, 555, 747; other measures against, 19, 25, 44, 46, 61, 74, 79, 99, 108, 125, 137, 149, 168, 174, 192, 254, 259, 260, 272, 291, 299, 320, 389, 401, 442, 449, 478, 479, 480, 488, 500, 507, 510, 525, 541, 544, 554, 555, 557, 588, 608, 613, 647, 652, 653, 658, 689, 713, 723, 732, 733, 745, 758; tests of organic compounds on, 575; susceptibility of, to injections of nicotine, 332. Cydia splendana (with var. reau-

murana), found in chestnuts for export from Italy, 384.

Cydia strobilella, bionomics of, on Picea jezoensis in Japan, 516.

Cydonia lunata, predacious Aphis laburni in French Sudan, 531.

Cydonia vicina, predacious on Aphis laburni in French Sudan, 531. cydoniae, Corythuca.

Cylas formicarius (Sweet Potato Weevil), in Ceylon, 358; in Loochoo Islands, 669; hibernation of, in Louisiana, 286; factors affecting infestation by, in Porto Rico, 223; on morning glory, 669.

Cylas formicarius var. elegantulus, measures against, on sweet potato

in Cayman Islands, 269.

cylindricum, Sinodendron, cylindrirostris, Orthorrhinus.

Cylindroiulus frisius (see C. oweni). londinensis Cylindroiulus teutonicus, bionomics and control of, in Germany, 461, 468.

vlindroiulus oweni, destroying carrots in Germany, 468. Cylindroiulus

Cylindromyia armata, parasite of Chlorochroa sayi in Montana, 44. Cylindronotus (see Helops).

cylindrus, Platypus.

Cymbopogon citratus, tests of oil of, against Aphids and mites, 249, 250.

cymoides, Nysius.

Cynara (see Artichoke).

cynica, Eriopyga.

Cynipids, as parasites of insects, 496.

Cynodon dactylon, relation of Schistocerca paranensis to, in Argentina, 147: Antonina indica on, in Hawaii, 585.

Cynoglossum, Empoasca libyca on, in Spain, 700.

Cynosurus cristatus, Geomyza tripunctata on, in Britain, 382.

cyperi, Carolinaia.

Cyperus, Pseudococcus brevipes on, in Formosa, 263.

Cyperus rotundu⁻, Carolinaia cyperi on, in Porto Rico, 40.

Cyphocleonus (see Cleonus).

Cyprus, pests of carob in, 106, 458; pests of fruit trees in, 107, 108, 259, 260, 706; miscellaneous pests in, 107, 108; new Braconid in, 106; failure to establish Aphelinus mali in, 108; summary of plant pest legislation in, 107.

Cyrtomorpha flaviscutellaris, attacking eggs of Austroicetes in W. Australia, 169, 629; prepupal and pupal stages of, 629.

Cyrtopeltis, status and scope of, 688. Cyriopeltis geniculata, Fieb., type of genus, 688, 689.

Cyrtopeltis geniculata, Reut. (see Engytatus).

Cyrtopeltis tenuis (see Engytatus).

Cyrtopeltis varians (see Engytatus geniculatus).

Cyrtorhinus, predacious on Megamelus proserpina in Hawaii, 213. cyrus, Telenomus.

Cytherea obscura, bionomics of, in

Italy, 163.

Czechoslovakia, pests of crucifers in, 656, 657, 718; new Coccid on Festuca in, 560; forest pests in, 215, 657, 658, 659, 715; miscellaneous pests in, 657, 658; orchard pests in, 620, 656, 657, 658, 659; Noctuid on Typha latifolia in, 345; parasites of Lepidoptera in, 529.

D.

dacicida, Tetrastichus.

Dacnusa bathyzona, parasite of Agromyza simplex in Britain, 33. Dactylis glomerata, Geomyza tripunc-

tata on, in Britain, 382.

Dactylopius vitis (see Pseudococcus). Dacus, parasites of, in Africa, 492; parasites and biological control of, in Fiji, 58, 59.

Dacus ciliatus (brevistylus), parasites of, in Nigeria and Zanzibar,

492.

Dacus cucurbitae, in Ceylon, 358, 359; in Formosa, 206, 436, 517, 667, 668; in Hawaii, 213, 325, 359, 752; in India, 242, 359, 765; danger of establishment of, in Japan, 667, 668; in Philippines, 181; on cucurbits, 181, 206, 325, 358, 359, 667, 765; other host fruits of, 181, 242, 325, 668, 752; susceptibility of varieties of tomato to, 213; bionomics of, 182, 206, 667; effects of low temperature on, 517; parasites and biological control of, 359; other measures against, 182, 325; susceptibility of stages of, to carbon bisulphide, 436.

Dacus dorsalis (see D. ferrugineus

dorsalis).

Dacus ferrugineus, in Ceylon, 359; in Hawaii, 359; in India, 242; in Malaya, 359; in Queensland, 246; host fruits of, 242, 246, 359; parasites and biological control of, 359.

Dacus ferrugineus dorsalis, apparently introduced into Bonin Islands from Saipan, 672; in Formosa, 207, 263, 517; in India, 765; host fruits of, 207,

263, 672, 765; effects of temperature on, **517;** measures perature on, 517; measures against, 207, 672; susceptibility of stages of, to carbon bisulphide, 263.

Dacus humeralis, parasites of, in

Africa, 492.

Dacus longistylus, in fruits of

Calotropis in India, 242.

Dacus oleae (Olive Fly), in Cyprus, 108; in Greece, 71; in India, 242; in Italy, 161, 261, 350; parasite of, in Libya, 261; in Malta, 469; in Spain, 700; bionomics of, 161; measures against, 71, 108, 161, 350, 700.

Dacus passiflorae, in Fiji, 58, 497, 702: Syntomosphyrum indicum imported from New South Wales against, 702; host fruits of, 497, 702; baits for, 58.

Dacus pectoralis, parasites of, in Zanzibar, 492.

Dacus punctatifrons, parasite of, in Sierra Leone, 492.

Dacus tryoni (see D. ferrugineus).

Dacus xanthodes, Syntomosphyrum indicum imported into against, 702; on guava, 702. Dacus zonatus, in India, 242.

Dadap (see Erythrina).

daghestanica, Podonta.

Dahlia, Aphids on, in Tanganyika, 36; pests of, in U.S.A., 15, 286, 723.

Dakota, North, Anabrus simplex in, 580 : Camnula pellucida in, 681.

Dakota, South, Anabrus simplex in, 580; Conopid parasites of bees in, 133.

Dalbergia sisso, guide to pests of, in India, 531.

Daldinia concentrica, Xiphydria prolongata associated with fungus resembling, in Britain, 632.

Damson, pests of, in Britain, 191, 297; pests of, in Germany, 251, 718.

Dasychira pudibunda, measures against, in forests in Germany, 69; infected with polyhedral disease, 70.

Dasyneura affinis, on violets in Britain, 350.

Dasyneura brassicae, on Camelina sativa in Ukraine, 258.

Dasyneura lini, bionomics of, on flax in India, 181, 242.

Dasyneura mali, measures against, on apple in Massachusetts, 647. Dasyneura tetensi, on black currant in Britain. 612.

INDEX.

Dasyneura vaccinii, on cranberry in U.S.A., 172, against, 172. 646: measures

Dasynus antennatus (on Citrus), in Ceylon, **359**; in India, **55**.

Dasyscapus parvipennis, proposed introduction of, into Santo Domingo against thrips, 10.

Datana integerrima, on pecan in Florida, 587.

datanae, Winthemia (see W. quadripustulata).

Date Palm, pests of, in Sudan, 513; book on pests of, in U.S.A.,

Date Scale (see Parlatoria blanchardi).

Datura fastuosa, insecticidal properties of, 360.

Datura stramonium, Ptinus tectus feeding on dried, in Germany, 164.

davidsoni, Crabro.

Davisonia, gen. n., key to N. American species of, 78.

dea, Indarbela (Arbela).
Death-watch Beetle (see bium rufovillosum).

Decahydronaphthalene, injected into timber against ants, 219.

decaudata, Agamermis.

decemlineata, Leptinotarsa (Chrysomela).

deceptor, Pseudococcus. decoratus, Geocoris.

Dectes spinosus, introduced into Australia against Xanthium pungens, 450.

deesae, Glyptomorpha (Stenobracon). deflorata, Cassida.

defluiteri, Pseudococcus.

defoliaria, Erannis (Hybernia).

Deguelin, plants containing, 29, 36. 616; forms and estimation of, 36, 616.

Deilephila (see Celerio).

dejeani, Corigetus; Oncideres.

Delaware, cereal pests in, 545, 730: miscellaneous pests in, 443, 545; orchard pests in, 41, 443, 444, 551; liberation of Tiphia spp. against Popillia japonica in, 545; introduction of Aphelinids into Haiti from, 41.

Delaware Bay, Jassids migrating across, 556.

delicatus, Macrocentrus.

Delitia Method, for fumigation with hydrogen phosphide, 519.

Delomerista japonica, parasite of Diprion nipponicum in Japan, 296.

Delphinium, Tarsonemus latus on,

in New Hampshire, 20.

Deltocephalus dorsalis, on rice in Japan, 167, 439; transmitting dwarf disease, 167; taken in light traps, 439.

demodocus, Papilio.

Dendroctonus, oil sprays against, on pines, 394.

Dendroctonus brevicomis, on Pinus ponderosa in California, 394.

Dendroctonus frontalis, aseptic rearing of, 28.

Dendroctonus jeffreyi, on Pinus jeffreyi in California, 394.

Dendroctonus monticolae, on Pinus spp. in California, 394.

dendrolimi, Trichogramma.

Dendrolimus albolineatus, on conifers in Japan, 516.

Dendrolimus jezonsis, on conifers in

Japan, 516.

Dendrolimus spectabilis, in China, 119; in Japan, 669, 671; in Manchuria, 672; on pines, 669, 672; suitability of various conifers for, 672; parasites of, 119, 669, 671.

dendrolimusi, Trichogramma (see T. dendrolimi).

dendrouympha, Brenthia.

Dendrotettix quercus, outbreak of, on oak in U.S.A., 588; parasite of, 588.

Dendrurgus (see Poecilips).

Denmark, Coleophora spissicornis on clovers in, 155; Melolontha spp. in, 468; Thaumetopoea pinivora in, 518; associations between Nematodes and insects in, 49.

dentata, Eurytoma.

dentipes, Chrysobothris; Mono-dontomerus.

deorum, Ageneotettix.

Depressaria heracleana, on parsnip in Britain, 297.

Depressaria nervosa (Caraway Moth), legislation and measures against, in Germany, 719.

depressella, Emmalocera.

Dermestes, tests of cold storage against, in cow hair in Holland, 599; bait for, 153.

Dermestes aurichalceus, 153.

Dermestes frischi, 153.

Dermestes lardarius, bionomics and control of, attacking silkworm pupae etc. in Italy, **153.**

Dermestes vorax, bionomics and control of, attacking silkworm pupae etc. in Japan, 515.

Dermestes vulpinus, 153; Clerid predactions on, in timber in New South Wales, 451; oviposition of, 164, 165; effect of temperature on, 528.

Derocrepis (see Crepidodera).

derogata, Sylepta.

Derris, against Aphids, 34, 35, 128, 309, 327, 443, 523, 543, 645, 651, 675, 740; against Coccids, 361, 574, 575; against other Rhynchota, 15, 35, 361, 397, 690, 739, 755; against Coleoptera, 17, 65, 132, 164, 183, 194, 209, 218, 265, 268, 285, 291, 297, 299, 312, 327, 371, 382, 383, 418, 421, 443, 522, 527, 543, 550, 551, 566, 572, 645, 651, 743, 762; against Diptera, 16, 172, 333, 466, 467, 713; against Lepidoptera, 35, 50, 172, 183, 207, 218, 327, 347, 348, 399, 409, 421, 422, 443, 510, 529, 543, 547, 548, 563, 646, 648, 651, 652, 674, 675, 723, 735, 752; against mites, 191, 192, 328, 543, 553, 583, 648, 709, 712; against saw-flies, 183, 191, 348, 543, 614, 615, 708, 748; against thrips, 191, 192, 274, 333, 553; tests of toxicity of, to insects, 17, 34, 35, 36, 37, 183, 347, 361, 371, 390, 397, 405, 421, 422, 572, 610, 616; in baits, 50, 713; dusting with, 15, 17, 35, 36, 65, 132, 164, 183, 194, 207, 209, 268, 285, 291, 297, 299, 312, 327, 333, 347, 348, 371, 382, 383, 397, 399, 409, 418, 443, 466, 467, 522, 527, 529, 550, 566, 572, 583, 652, 674, 675, 712, 735, 740, 766; carriers for, 15, 17, 65, 268, 371, 397, 399, 550, 572, 652, 735, 740; conditioning 675. agents for dusts of, 285, 327 aeroplane dusting with, 563, 748; spraying with, 16, 17, 34, 128, 172, 274, 297, 309, 327, 382, 421, 443, 484, 510, 523, 542, 543, 547, 548, 551, 553, 566, 614, 615, 645, 646, 652, 674, 675, 723, 739, 743, 752, 755, 762, 766; comparison of water suspensions and extracts of, in sprays, 648, 674; and soap, 309, 484, 755; other spreaders and adhesives for, 16, 128, 191, 274, 382, 443, 543, 547, 548, 551, 553, 615, 645, 646, 674, 690; other uses of, against insects, 218, 267, 328, 645; formulae containing, 16, 17, 128, 172, 183,

191, 192, 268, 274, 399, 443, 542, 547, 551, 563, 614, 615, 645, 646, 652, 713, 735, 743, 755; and copper ammonium silicate, 752; and nicotine, 65, 172, 740, 741 and oil emulsions, 191, 192, 510, 708: intermediary solvents for, when used with oils, 574; vegetable oils and acetone increasing toxicity of, 739; and pyrethrum, 268, 421, 422, 522, 523, 553, 740, 762; and sulphur, 65, 397, 443; and thiocyanate, 740, 741; rotenone content of, 15, 17, 34, 35, 128, 183, 274, 299, 333, 347, 348, 361, 371, 382, 383, 397, 399, 443, 543, 547, 550, 553, 563, 614, 645, 646, 651, 652, 675, 708, 735, 739, 740, 741, 743, 766; chemistry, evaluation and standardisation of, 120, 156, 183, 258, 331, 616, 631, 651, 674, 675, 731, 766; importance of fineness of particles of, 651, 674: duration of activity of dusts and sprays of, 421, 422; translocation of constituents of, in bean plants, 332; possible danger of, to man and fish, 383, 529; method of harvesting and drying, 120; other insecticides compared with, 17, 34, 35, 36, 90, 132, 191, 258, 274, 333, 347, 348, 443, 522, 523, 547, 553, 614, 651, 690, 723, 735, 740.

Derris spp., pests of, in Porto Rico, 224; Megachile seychellensis defoliating, in Seychelles, 433; insecticidal properties and constituents of, 37, 183, 224, 318,

361, 631, 675, 766.

Derrisol, spraying with, against Aphids, 241.

Desert Locust (see Schistocerca gregaria).

desertus, Gryllulus (Gryllus).

designata, Epicauta.

Desmia funeralis, parasites of, on vines in California, 19, 285, 286.

Desmidophorus breviusculus, bionomics and control of, on cacao and kapok in Indo-China, 185, 186.

Desmodium canadense, Calomycterus setarius on, in Connecticut,

destructor, Aleurodicus; Aspidiotus; Hadrobregmus; Halotydeus; Lepidoglyphus (Glycyphagus); Mayetiola (Phytophaga); Merisus; Monomorium.

Detal (dust containing dinitrocresol), as an insecticide, 69, 345, 661.

Deudorix bimaculata (see Virachola). Deudorix epijarbas, bionomics of, on litchee in China, 693.

Deudorix lorisona (see Virachola).

devastans, Empoasca.

devastator, Melanoplus.

devoniensis, Eriococcus.

Dewberry, Tortrix rosaceana on, in Utah, 141.

Dexia rustica, parasite of white grubs in Britain, 317; introduced into Mauritius, 317.

Dextrin, and blood albumen, as an adhesive for dusts, 388.

Diabrotica connexa, in Mexico, 28; transported in vegetables U.S.A., 28.

Diabrotica duodecimpunctata, maize and cucurbits in U.S.A., 124, 269, 391; resistance of maize varieties to, 124, 391.

Diabrotica longicornis, on maize in Connecticut, 726.

Diabrotica melanocephala, measures against, on cucurbits in U.S.A., 17, 269, 285, 333, 647.

Diabrotica pallipes, on cucurbits in Jamaica, 490.

Diabrotica soror, in U.S.A., 19, 291: on apricot and peach, 19; derris against, 291.

Diabrotica speciosa, new parasite of, in Argentina, 689.

Diabrotica themei (see D. pallipes). Diabrotica vittata (see D. melanocephala).

Diachasma fullawayi (parasite of Trypetids), in Hawaii, 84; in Kenya and Uganda, 492.

Diachasma tryoni, parasite of Ceratitis capitata in Hawaii, 84.

diadema, Stomatodexia (Leskiopalpus).

Dialeurodes chittendeni, on rhododendrons in Britain, 350.

Dialeurodes citri (Citrus Whitefly), in India, 765; new parasite of, in Japan, 296; measures against, in U.S.A., 286, 690; attempted eradication of, in California, 18. 747.

Diamond Back Moth (see Plutella maculipennis).

dianthi, Ocladius; Protostrophus.

Diaphania hyalinata (on cucurbits). in Jamaica, 490; in Missouri, 269.

Diaphania nitidalis, on cucurbits in Missouri, 269.

Diapheromera femorata, bionomics and control of, in Michigan, 80.

Diaphnidia pellucida (predacious on Typhlocyba pomaria), bionomics of, in New York, 557, 558. 559; insecticides destroying. 558, 559.

Diaphorina citri, on Citrus in India,

765.

Diaprepes, bionomics and control of, on Citrus in West Indies, 402. 405, 448, 751.

Diaprepes abbreviatus, in W. Indies, 222, 402, 751; on Citrus, 402, 751; hand-collection against, on sugar-cane, 222.

Diaprepes boxi, sp. n., food-plants of, in St. Lucia, 405.

Diaprepes esuriens, on Citrus in W. Indies, 402, 751.

Diarthronomyia hypogaea, on chrysanthemums in Britain, 349.

Diaspis, species of, in Italy, 496. Diaspis bromeliae, restricted to pineapple in Formosa, 263.

Diaspis leperii (see Epidiaspis). Diastocera reticulata (see Analeptes).

Diatomaceous Earth, as a carrier for dusts, 65, 129, 133, 327, 328. 389, 652, 725. (See Kieselguhr.)

Diatraea (on sugar-cane), in Brazil, 82: hand-collection against, in China, 243; in Br. Guiana, 369; in India, 243; in W. Indies, 10, 82, 121; parasites and biological control of, 10, 82, 121, 243.

Diatraea canella, on sugar-cane in Br. Guiana, 221, 369, 491; parasites and biological control of,

221, 491.

Diatraea crambidoides, on maize in S. Carolina, 644.

Diatraea saccharalis (Sugar-cane Moth Borer), in Argentina, 390, 391; in Br. Guiana, 221, 369, 491; in Peru, 390, 391; in U.S.A., 218, 390; in W. Indies, 40, 43, 221, 269, 288, 448, 490; susceptibility of varieties sugar-cane to, 221, 490; damage to rice by, 218; method of determining larval dispersion of, 448; parasites and biological control of, 43, 221, 222, 289, 369, 390, 391, 448, 491; predacious enemies of, 269, 288.

Diatraea (Argyria) sticticraspis, on sugar-cane in India, 261, 408, 499; varietal susceptibility of sugar-cane to, 499; parasite of, 408.

Diatraea venosata (striatalis) (on sugar-cane), in Formosa,

673; in India, 408, 499; bionomics of, in Madagascar, 67: in Mauritius, 176, 317; varietal susceptibility of sugar-cane to, 263, 317, 499; natural enemies of, 67, 317, 408; effect of submersion on, 673.

Diaxenes phalaenopsidis, sp. n., bionomics of, on orchids in Netherlands Indies, 262.

Dibenzothiaphene, less effective than calcium arsenate against Anthonomus eugenii, 328.

Dibrachys cavus (boucheanus), hosts of: in Hungary, 577; in Switzerland, 100; in U.S.A., 577, 580, 724.

Dicerca aenea, on poplar and ash in

Russian Union, 429.

Dichlorethyl Ether, as intermediary solvent for derris in oil sprays, 574, 575; as a soil fumigant, 323, 747; use of, against peach borer, 323.

Dichocrocis punctiferalis, on ginger and turmeric in Ceylon, 359; on papaya in Queensland, 246.

Dichomeris ianthes, predacious enemies of, on Indigofera endecaphylla in Seychelles, 433.

Dichomeris marginella (on juniper), 296; in Czechoin Britain, slovakia, 657.

dicincta, Mylabris.

Dicranura vinula, on apple in Britain, 612; on poplar and willow in Germany, 461.

Dicrodiplosis guatemalensis, sp. n., predacious on mealybugs Guatemala, 692.

Dictyoploca japonica, parasite of, in Japan, **310.**

dictyospermi, Chrysomphalus.

Dicyphus, status and scope of, 688. Dicyphus minimus, on tobacco in U.S.A., 393.

Dicyphus nicotianae (nocivus) (see Engytatus tenuis).

didyma, Hadena (see Trachea secalis). Diesel Oil, 82.

differentialis, Melanoplus.

difficilis, Eusattus.

Toxoptera Digitaria sanguinalis, graminum transmitting mosaic to sugar-cane from, in U.S.A.,

Diglycol Oleate, effecting inversion of arsenical spray mixtures, 174; as spreader for nicotine thiocyanate, 704.

Diglycol Stearate Wax, as spreader and adhesive for cubé, 647.

dignoides, Telenomus.

dignus, Telenomus (Phanurus).

Dihydrodeguelin, toxicity of forms of, to insects, 617.

Dikraneura cedrelae, sp. n., Cedrela odorata in Porto Rico, 288.

dilatata, Paropsis.

Dill (see Anethum graveolens).

Dillenia indica, insecticidal properties of, 360, 361.

Diloba coeruleocephala, on almond in Palestine, 514.

Dilophus febrilis (vulgaris), new Nematode parasite of, in Denmark, **49**.

Dimethyl Sulphate, characterisation of oils by solubility in, 99.

dimidiata, Leis; Rondania.

dimidiatus, Carpophilus; Tyrophagus (Coelognathus, Tyroglyphus) (see T. putrescentiae).

dimmocki, Pikonema (Nematus). Dimmockia kampalana, sp. n., parasite of Pyralid in Uganda, 610.

Dinarmus, associated with Eumarchalia gennadii in Crete, 71.

diniana, Enarmonia (Semasia).

diniasalis, Pyrausta.

Dinitro-cresol, increasing toxicity of oil emulsions to Aphids, 62; toxicity of, to Cydia pomonella, 576: dusts containing, against Bupalus piniarius, 313; method for determining, 600. (See Detal.)

Dinitro - ortho - cyclohexylphenol, dusts containing, against thrips, 508; oil sprays impregnated with: against Aphids, 272; against Coccids, 25, 270, 272, 549, 562; against Eucosma ocellana, 562; causing injury to pine, 270, 549.

Dinoderus bifoveolatus, in stored cassava tubers in Madagascar, 455; effect of infestation by, on rotenone content of cubé roots,

396.

Dioctes (see Angitia).

diomphalia, Lachnosterna (Holotrichia).

dionysius, Phyllognathus.

Dioryctria abietella, parasite of, in Czechoslovakia, 529.

Diorymerellus laevimargo (Orchard Weevil), in U.S.A., 550, 692; danger of introduction of, into Hawaii, 692; feeding habits and control of, 550.

Diospyros kaki (see Persimmon). Diparopsis castanea (Red Cotton Bolworm), in Portuguese E. Africa, 199: in S. Africa, 199, 365; in Gambia, 199; in Nigeria, 667; in Nyasaland, 113, 199, 367; in S. Rhodesia, 199, 366; in Sudan, 511; in Tanganyika, 199; distribution of wild foodplants of, 199, 365; bionomics of, 365; measures against, 113, 200, 366, 367, 511.

Diphtera alpium (see Moma).

Diplazon (see Bassus).

Diploschema rotundicolle, bionomics of, on Aleurites and Citrus in Brazil, 502.

Diplosis, bionomics of, on pear in

Korea, 605.

Diplostichus tenthredinum, parasite of Diprion pini in Holland, 348. Diplotaxis atlantis, on strawberry in Connecticut, 18.

Diprion, introduction of European parasites of, into Canada, 103.

Diprion frutetorum, utilisation of Microplectron fuscipenne against, on Pinus sylvestris in Ontario, 538.

Diprion nipponicum, new parasites of, on pine in Japan, 295, 296.

Diprion pini (on pines), in Germany, 189; in Holland, 183, 348, 621; causes and detection of outbreaks of, 621; natural enemies of, 348; tests of dusts

against, 183, 348.

Diprion polytomum (on spruce), in Canada, 77, 103, 215, 329, 330, 487, 534, 538; in Czechoslovakia, 215; in U.S.A., 329, 330, 487; methods for estimating importance of injury by, 330; bionomics of, 215, 487, 534, 538; cytology of strains of, 215, 487; adult morphology of, 247; parasites and biological control of, 77, 103, 329, 487.

Diprion sertifer (on pines), Germany, 189; in Hungary, 198, 341; in Japan, 296; in Russian Union, 429; in Sweden, 198: in U.S.A., 580; parasites of, 198,

296, 341.

dipterophaga, Thysanus (Signi-

phora).

Dirhinus (parasite of Dacus spp.), introduced into Fiji from Hawaii, 59; in Zanzibar, 492; sitising house-flies, 59.

Dirhinus auratus, introduced into Ceylon against Dacus cucurbitae,

Dirphya, on coffee in N. Rhodesia, 500.

Dirphya usambica, on Rytigynia schumani in Tanganyika, disciguttus, Eutettix.

Discochaeta yponomeutae, parasite of Hyponomeuta in Ukraine, 302.

discolor, Myllocerus.

Disonycha laevigata, on beet in Jamaica, 490.

Disophrys, parasite of Acontia graellsi in Nyasaland, 114.

dispar, Calotermes; Lepidosaphes (Mytilaspis); Lymantria (Porthetria); Xyleborus. disparis, Pimpla.

Dissosteira longipennis, outbreaks of, in Oklahoma, 139, 580.

disstria, Malacosoma.

distigma, Chilocorus.

distincta, Galerucella; Zenillia.

distinguendus, Lariophagus.

diversana, Tortrix.

diversor, Echthromorpha (see immaculata).

diversus, Microdus (Bassus); Macrosteles (Cicadula).

Dociostaurus maroccanus, in Algeria, 685; in Cyprus, 108; losses and expenditure due to, in Jugo-slavia and Iran, 685; in Italy, 162, 592; in Libya, 260; in Spain, 698; colour variations and phases of, 162, 592, 593, 685; bionomics and ecology of, 72, 162, 685, 698; natural enemies of, 163, 593; measures against, 685, 698, 699.

Dock (see Rumex).

Sawfly (see Dock Ametastegia glabrata).

Dodecyl Thiocyanate, 712.

Dogwood (see Cornus).

Doirania leefmansi, parasite of Sexava in New Guinea Territory, 436.

Dolerus harukawai, on Juncus in Japan, **672.**

Dolichoderus bituberculatus, in Netherlands Indies, 262.

Dolichos, Brachyplatys testudonigra on, in Nyasaland, 114.

Dolichos lablab, Bruchids on, in Burma, 7, 8; Heliothis armigera on, in S. Rhodesia, 366; cotton pests on, in Sudan, 513; negative experiment with poisoned, against termites, 512.

Doloessa viridis, in copra and stored rice in Malaya, 31.

Dolvcoris baccarum (on cereals), bionomics of, in Germany, 51, 52; in Switzerland, 91; characters of, 51.

Dombeya, Lamiid on, in Nyasaland, 114.

Dombeya rotundifolia, Dysdercus nigrofasciatus on, in S. Africa, 365, 366.

domestica, Musca; Thermobia.

domesticus, Glycyphagus; Gryllulus (Gryllus); Opilo; Sclerodermus.

dominans, Charops.

Dominica, Citrus pests in, 751; summaries of plant quarantines in, **229, 491.** dominica, Rhizopertha.

Dominican Republic (Santo Domingo), beneficial insects and biological control in, 10, 41; summary of plant quarantines in, 229.

Doralis (see Aphis).

Doraphis populi, Lycaenid predacious on, on poplar in Japan, 673.

Doratifera vulnerans, measures against, on Eucalyptus and apricots in New South Wales, 111.

rsalis, Acarapis; Chalepus; Dacus (Chaetodacus) ferrugineus; dorsalis, Deltocephalus; Scirtothrips; Shirakia (see S. schoenobii).

dorsana, Cydia (Laspeyresia).

orientalis, Dorylus in ginger rhizomes in Ceylon, 359.

doryphorae, Doryphorophaga.

Doryphorophaga spp. (parasites of Leptinotarsa decembineata), imported into France from N. America, 594.

Dorysthenes huegeli, on apple in India, **409.**

Douglas Fir (see Pseudotsuga taxifolia).

Dracocephalum moldavica, effect of oil of, on insects and mites, 249, 250.

Draeculacephala mollipes, possibly causing galls on sugar-cane in Hawaii, 585; on narcissus in U.S.A., 288.

Dreft, spray formula containing,

dregii, Calidea.

Drevfusia (see Chermes).

Drosicha contrahens, on Citrus in India, 765.

Drosophila spp., in unsound tomatoes in U.S.A., 397, 398; method of testing insecticides on, 536.

druparum, Syntomaspis.

dubia, Anomala.

dubius, Dyscinetus.

Ducks, utilisation of, against rice pests, 498.

Dung, in baits for locusts, 698. duodecimpunctata, Diabrotica.

Duplaspis fraxini, gen. et sp. n., on Fraxinus excelsior in France, 248. duplex, Pseudaonidia.

duplicatus, Ips.

Dusting, types of equipment for, 164, 291, 347, 418, 443, 482, 483, 643, 682; use of aeroplanes for, 19, 398, 426, 433, 682, 748; compared with spraying, 443, 467, 522, 564, 647, 726, 734.

Dusts, action of, desiccating insects, 252; apparatus for testing contact, 253; machine for mixing, 357; use of oil-impregnated, 357, 426, 478, 653, 734.

Dutch Elm Disease (see Ceratos-

tomella ulmi).

Dutox, 447, 736. (See Barium Fluosillicate.)

Dyera costulata, pests of, in Malaya, 109, 764; Minthea rugicollis in timber of, 764.

Dyk's Method, for estimating prevalence of *Lymantria monacha*, **659**, **665**.

Dyscinetus bidentatus (see Euetheola).

Dyscinetus dubius (geminatus), on sugar-cane in Br. Guiana, **369.**

Dysdercus (Cotton Stainers), in S. Africa, 365; in N. and S. Rhodesia, 366; susceptibility of varieties of cotton to, in Nigeria, 666; in Nyasaland, 367; in Tanganyika, 113; bionomics of, in Uganda, 372–375, 376; Nematospora associated with, 375, 376; measures against, 113, 375.

Dysdercus cardinalis (on cotton and other plants), in Sudan, 512; in Tanganyika, 642; in Uganda, 372, 374; bionomics of, 372.

Dysdercus cingulatus, in Indo-China, 186, 206; baits for, on cotton and kapok, 186; low toxicity of extracts of Stemona to, 206.

Dysdercus fasciatus (on cotton), in S. Africa, 365, 366; in N. Rhodesia, 367, 500; in S. Rhodesia, 90, 366; in Sudan, 512; in Tanganyika, 642; in Uganda, 372–375; date of planting cotton in relation to, 366, 367; bionomics and alternative food-plants of, 365, 366, 367, 372–374, 512, 642; parasites of, 90, 375.

Dysdercus howardi, selenium treatment of cotton against, in Trinidad, 66.

Dysdercus intermedius, on Sterculia in S. Africa, 365; date of planting cotton in relation to, in S. Rhodesia, 366.

Dysdercus mendisi, sp. n., on cotton in Brazil, **268.**

Dysdercus melanoderes, on cotton

in Belgian Congo, **315.**Dysdercus mimus, food-plants of,

in Peru, **533.**

Dysdercus nigrofasciatus, bionomics of, in S. Africa and Uganda, 365, 366, 372-375; food-plants of, in Sudan, 512; natural enemies of, 375.

Dysdercus orientalis var. pulcher, on cotton in Tanganyika, 112.

Dysdercus ruficollis, measures against, on cotton in Peru, **533.**

Dysdercus superstitiosus, in Belgian Congo, 315; in Nigeria, 666, 667; in N. and S. Rhodesia, 366; incidence of, in Sudan, 512; in Uganda, 372–375; date of planting and susceptibility of varieties of cotton to, 366, 666, 667; bionomics and alternative food-plants of, 366, 367, 372–374, 512; parasite of, 375.

Dysdercus ugandanus, parasite of,

in Uganda, **375.**

E.

Earias cupreoviridis (on cotton), in China, 498; in India, 408, 409.

Earias fabia, on Hibiscus and cotton in India, 243, 408, 409; bionomics and control of, in Java, 766; parasites of, 243.

Earias huegeli, on cotton in Queensland, 494.

Earias insulana, on Hibiscus and cotton in India, 243, 408, 409; food-plants of, in Sudan, 511; parasites of, 243.

Earth Mite, Red-legged (see Halo-tydeus destructor).

Earwig, European (see Forficula auricularia).

ebeninum, Monomorium carbonarium.

ebenus, Ligyrus.

Eberhardia, classification and new species of, 94, 95.

Eberhardia agilis, Can., in Italy, 95. Eberhardia agilis, Mich. (see E. michaeli).

Eberhardia alberti, sp. n., 94; on decaying fungi in England, 95.

Eberhardia (Tyroglyphus) krameri, type of Cosmoglyphus, 95; new mite recorded as, 95.

Eberhardia michaeli, type of genus, 95; in decaying cabbage in England, 95.

Eberhardia oudemansi, sp. n., in ants' nests in Java, 95.

Eberhardia redikorzevi, sp. n., in mouldy grain in Russia, 95.

Eberhardia rhizoglyphoides, sp. n., in flax seeds in Russia, 95.

Echnocnemus bipunctatus, warmth accelerating development of hibernating, **54**.

Echthromorpha immaculata (diversor), parasite of Tirathaba trichogramma in Fiji, 334.

Echthromorpha tirathabae, parasite of Tirathaba trichogramma in Fiji, 334.

echinopus, Rhizoglyphus.

Economic Entomology, organisation and development of: in Mexico, 248; in Porto Rico, 116; in U.S.A., 136; literature of, in America, 367, 772.

Ectatomma ruidum, destroying

Ectatomma ruidum, destroying Sacadodes pyralis in Colombia,

403.

Ectopiognatha, parasite of Perkinsiella saccharicida in Hawaii and Mauritius, 454.

Mauritius, 454.

Ectropis bhurmitra (see Boarmia).

Egg-plant, Epilachna vigintioctopunctata on, in Ceylon, 358; pests of, in India, 242, 361; pests of, in Jamaica, 490; Empoasca benedettoi on, in Palestine, 77; Dacus cucurbitae on, in Philippines, 181; thrips on, in Sierra Leone, 240; Asphondylia on, in Tanganyika, 642; pests of, in U.S.A., 13, 64, 217, 395, 744; insects and virus diseases of, 77, 759; relation of, to Pentatomid predacious on Leptinotarsa, 13.

Egypt, cotton pests in, 457, 458, 685; miscellaneous pests in, 456, 457; new Aleurodid on Zizyphus spina-christi in, 440; Thermobia aegyptiaca damaging books in, 706; Stromatium fulvum in timber in, 591; locusts in, 112, 682, 685, 687; beneficial insects and biological control in, 212, 450, 456, 457, 592; beneficial insects introduced into other countries from, 275, 458;

summaries of plant pest legislation in, 134, 385, 682, 692.

Egyptian Grasshopper (see Anacridium aegyptium).

Elachertus monachae, bionomics of, in Germany, 233.

Elachertus tabashii, sp. n., parasite of Naranga aenescens in Korea, 768.

Elaeis guineënsis (see Oil Palm).
Elasmus, parasite of Earias spp. ir

Elasmus, parasite of Earias spp. in India, 243.

Elasmus albopictus, parasite of Schoenobius bipunctifer in China, 109.

Elasmus maculatus, parasite of Apanteles americanus in Cuba, 171.

Elasmus nephantidis, parasite of Nephantis serinopa in Cochin, 452.

Elasmus platyedrae, failure to introduce, into Texas against Platyedra gossypiella, 275.

Elasmus punctulatus, parasite of Apanteles americanus in Cuba, 171.

Elasmus zehntneri, parasite of Scirpophaga in India, 243, 602; bionomics of, 602.

Elaterids (see Wireworms).

electa, Zonosemata.

electo, Colias.

Electric Cables, insects damaging, 232, 262.

Electricity, uses of, against pests, 251, 488.

elegans, Heterorrhina; Zonocerus. elegantulus, Cylas formicarius.

Eleodes spp., association of, with ants in Kansas, 292.

Eleusine coracana, pests of stored and field, in India, 204, 532; Acridids on, in Ceylon, 358.

Eleusine indica, Carolinaia setariae on, in Porto Rico, 39.

Elionurus adustus, relation of Schistocerca paranensis to, in Argentina, 147.

Elis annulata (see Campsomeris).

elisus, Lygus.

Ellice Islands, summary of plant quarantines in, 134.

elliotti, Aulocara.

ello, Erinnyis. Ellopia fiscellaria lug

Ellopia fiscellaria lugubrosa, on Abies in U.S.A., 581.

Ellopia somniaria, outbreaks of, not related to sun-spots, 486.

Elm (*Ulmus*), pests of, in Czechoslovakia, **657**, **658**; pests of, in Finland, **201**; *Lecanium corni* on, in Germany, 251; Scolytids on, in Holland, 348; Ceratostomella ulmi in, in Italy, 119; pests of, in Japan, 265, 452; Alsophila pometaria on, in New Brunswick, 539; pests of, in Russian Union, 75, 530; Galerucella luteola on, in Switzerland, 164; pests of, in U.S.A., 13, 15, 18, 20, 126, 136, 225, 325, 543, 546, 551, 576, 725, 749; identity of woolly aphis on, in N. America, 484, 485; relation of beetles to Ceratostomella ulmi in, 20, 325, 547, 551, 656, 657, 658, 749; susceptibility of species of, to C. ulmi, 656; reviews of data on C. ulmi in, 119, 749; Pyrausta nubilalis experimentally fed on, 475.

Elm Leaf Beetle (see Galerucella luteola).

Elm Scale, European (see Gossy-paria spuria).

elongata, Hieroglyphus banian.

elongella, Prospaltella.

elutella, Ephestia.

elzneri, Sturmia. emigrata, Perisierola.

Emilia, relation of Thysanoptera and pineapple yellow spot to, in Hawaii, 213, 692.

Emmalocera depressella, on sugarcane in India, 243, 261, 499, 532; parasite of, 243, 532; susceptibility of varieties of sugar-cane to, 499.

Emphytus coryli, sp. n., on Corylus avellana in Germany, 560.

Empoasca, on Hibiscus and cotton in Netherlands Indies, 358, 766; new Crabro predacious on, in Ohio, 369; causing leaf-curl on Hibiscus sabdariffa, 358; susceptibility of varieties of cotton to species of, 261, 309, 666, 667, 766.

Empoasca benedettoi, possible relation of, to virus disease of potato and egg-plant in Palestine, 77.

Empoasca biguttula (on cotton), in China, 498; in Formosa, 309.

Empoasca devastans, on cotton in India, 261.

Empoasca erigeron, on narcissus in U.S.A., 288.

Empoasca fabae, in Bermuda, 626; in U.S.A., 15, 140, 216, 271, 288, 399, 443, 564, 565, 609; varietal susceptibility of apple to, 271; on dahlia and narcissus, 15, 288;

effects of, on lucerne and groundnuts, 216; on potato, 140, 399, 564, 565; on rhubarb, 443; on watermelon, 626; bionomics of, 609; measures against, 15, 140, 399, 443, 564, 565, 609.

Empoasca facialis (on cotton), in Belgian Congo, 315; in Nigeria,

666, 667.

Empoasca flavescens, on potato in Germany, 68; not transmitting potato viruses, 69.

Empoasca libyca, measures against, on vines etc. in Spain, 700; on cotton in Sudan, 513.

Empoasca papayae, sp. n., associated with bunchy-top of papaya

in Porto Rico, 288.

Empoasca solana, acquiring virus of beet curly-top in California, 445; on potato in Hawaii, 214.

Empusa infesting Laphygma exempta in Nyasaland, 114; infesting Zonocerus elegans in Tanganyika, 642.

Empusa grylli, not experimentally infesting Schistocerca paranensis in Argentina, 147; infesting Nomadacris septemfasciata in Angola, 688.

Empusa muscae, infesting Melanostoma in Britain, 101.

Enaria melanictera, parasite of, in Madagascar, 176.

Enarmonia caryana, on pecan in Florida, 587.

Enarmonia diniana, on Pinus cembra, in Austria, 664.

Enarmonia prunivora, on apple in Br. Columbia, **59.**

Encarsia formosa (parasite of Trialeurodes vaporariorum), establishment of, in Australia, 449, 582; in Britain, 623; breeding and utilisation of, in Canada, 623; in New Zealand, 582.

Encarsia nigricephala, sp. n., parasite of Bemisia in Porto Rico, 41. Encarsia pergandiella, parasite of

Aleurodes azaleae in California, 748.

Endospermum, pests of, in Fiji, 703. Endospermum malaccense, Uraniid on, in Malaya, 110.

enecator, Trichomma. Engytatus, scope of, 688.

Engytatus geniculatus (Cyrtopeltis varians), on tomato in Hawaii, 214, 752; predacious habits of, 214; synonymy of, 689.

Engytatus (Cyrtopeltis) tenuis, apparently not transmitting tobacco

leaf-curl in India, 88, 89; probably transmitting virus disease of tomato in Libya, 260; synonymy of, 688.

Enicospilus nigrimarginalis, sp. n., parasite of Orgyia postica in Formosa, 119.

enigma, Oedaleonotus.

Ensina sonchi, bionomics and control of, on Taraxacum koksaghyz in Ukraine, 618, 619.

Entedon apionidis, sp. n., parasite of Apion xanthostylum in Tanganyika, 610.

ganyika, UIU.

Entedon laricinellae (see Chryso-

charis).

Entedon perturbans (parasite of Bruchids), distribution of, in Africa, 610.

Entedon pictus (see Cirrospilus).

Entomology, book on recent advances in, 47; glossary of, 11; bibliographies of: in Canada, 440; in Finland, 120, 695. (See Economic Entomology.)

entomophaga, Monieziella (Histio-

gaster).

Entomophthora sphaerosperma, infesting Dysdercus in Uganda, 375; sprays preventing infestation of Typhlocyba pomaria by, in Virginia, 557.

Entomoscelis adonidis, on crucifers in Saskatchewan, 170.

Eotetranychus (see Tetranychus).

Ephedrus nitidus, parasite of Aphids in Ontario, 537, 538. ephemeraeformis, Thyridopteryx.

Ephestia (in stored products), in Germany, 460; in Trinidad, 11; tests of methyl bromide against, in U.S.A., 389.

Ephestia cautella, used for breeding Trichogramma japonicum in Japan, 515, 670, 671; danger of introduction of, into Australia, 435; imported into Holland from Dutch Guiana, 529; in New Guinea Territory, 435; in stored products, 315, 435, 529; effects of temperature and humidity on eggs of, 670, 671; characters of eggs of, 196; leaves of Melia azadirachta as repellent for, 315.

Ephestia elutella, bionomics and suggested biological control of, in France 523, 524; in Holland, 183, 599; in U.S.A., 328, 748; in stored cacao, 301, 523, 599; in tobacco, 328, 523; in lima beans, 748; known only from

stored products, 301; characters of eggs of, 196; measures against, 328, 599; derris dust ineffective against, 183.

Ephestia figulilella (in dried fruit), in Australia, 449; in California, 29, 30, 118, 324, 389; feeding on fruits in the field, 30, 118, 324, 449; bionomics of, 324; recovery of larvae of, paralysed by Microbracon hebetor, 29; measures

against, 118, 324, 389.

Ephestia kuehniella, 662; measures against, in stored cereals in U.S.A., 130, 322, 326; mating of, 440; effect of temperature on, 118, 322; effect of sting of Microbracon hebetor on, 12; normal host of Nemeritus canescens, 1; other parasites reared on, 102, 103, 276; used for rearing Podisus maculiventris, 596; disease of, 34; characters of eggs of, 196.

Ephialtes, parasite of Cydia molesta

in Italy, 154.

Ephialtes extensor, 154; utilisation of, against Cydia pomonella in U.S.A., 323, 747.

Ephialtes sanguinipes (see Pimpla).

ephippella, Argyresthia.

Epiblema spp., alternative hosts of parasites of Cydia molesta in Ontario, 509, 510.

Epibootania nonvitta, sp. n. (nonvittata, nom. nud.), reared from galls on *Citrus* in New South Wales, **502**.

Epicaerus cognatus (Potato Weevil), bionomics and control of, in Mexico, 11; intercepted in Louisiana, 11.

Epicampocera succincta, parasite of Pieris rapae in Britain, 342.

Epicampoptera marantica, on coffee in Tanganyika, 113.

Epicauta adspersa, on potato, etc., in S. America, 290, 291.

Epicauta aterrima, on potato, etc., in S. America, 290, 291.

Epicauta atomaria, on potato, etc., in S. America, 290, 291.

Epicauta designata, destroying locust eggs in Nyasaland, 114.

Epicauta gorhami, measures against, in Japan, 265.

Epicauta maculata, on beet, potato, etc., in Montana, 44.

Epicauta velata, on cereals in Nyasaland, 114.

epicnapterus, Pimpla (Iseropus). Epicometis hirta, soil conditions in relation to, in Russia, 429.

Epidiaspis leperii (piricola), measures against, on fruit trees in Chile and Italy, 221, 351.

epigonus, Pleurotropis (see P. metallica).

epijarbas, Deudorix.

Epilachna, on potato and eggplant in Mysore, 361; toxicity of plant extracts to, 360, 361.

Epilachna chrysomelina, bionomics of, on melon in Morocco, 417.

Epilachna varivestis (corrupta) (Mexican Bean Beetle), in U.S.A., 13, 270, 285, 327, 333, 545, 566, 644, 651, 704; Podisus maculiventris not experimentally attacking eggs of, 13; measures against, 270, 285, 327, 333, 566, 651; tests of nicotine thiocyanate sprays on, 704.

Epilachna vigintioctopunctata, foodplants of, in Ceylon, **358.**

Epinotia laracana, bionomics of, on hawthorn in Ontario, 509; alternative host of Glypta rufiscutellaris, 509.

Epipolaeus caliginosus, on hops in Britain, 613.

Epitetranychus (see Tetranychus). Epiterimerus vitis, measures against,

on vines, **521.**Epitrix, on tobacco in U.S.A., **393.**Epitrix cucumeris, on potato in

U.S.A., 285, 400, 564, 643, 648; on tobacco in Connecticut Valley, 132; measures against, 132, 285, 564, 565, 643, 648.

Epitrix parvula, on Capsicum in Jamaica, 490; on tobacco in U.S.A., 132, 183, 393, 645, 744; on potato in Colorado, 643; measures against, 132, 133, 183, 643, 645.

Epiurus (see Pimpla).

Epochra canadensis, bionomics and control of, in Oregon, 60, 61.

equestris, Merodon.

Eragrostis, relation of Schistocerca paranensis to, in Argentina, 147.

Evannis defoliaria, on oak in Poland, 256.

erecta, Phorocera.

Eremnus angustirostris, sp. n., on maize in S. Africa, 106.

eremocitri, Aonidiella.

Eremocitrus glauca, new Coccid on, in Queensland, 226.

Eretmocerus aleurolobi, sp. n., parasite of Aleurolobus marlatti in Japan, 296.

Eretmocerus serius, controlling Aleurocanthus woglumi in Bahamas, 491.

Ergates faber, life-cycle of, in trees and timber in Germany, 301.

Erica, insects on, in Britain, 296, 418.

ericae, Nysius.

eridania, Xylomyges (Prodenia).

erigeron, Empoasca. erineus, Eriophyes tristriatus.

Erinnyis alope, on papaya and Manihot in West Indies, 171,

491; parasite of, 171. Erinnyis ello, parasite of, on papaya and Manihot in Cuba, 171.

Eriobotrya japonica (Loquat), new Eriophyid on, in California, 650; Trypetids on, in India, 242; Jassid on, in Japan, 208.

eriobotryae, Eriophyes. Eriocampoides (see Caliroa).

Eriochloa punctata, Pseudococcus brevipes on, in Brazil, 139.

Eriococcus azaleae, on azalea and rhododendron in U.S.A., 286, 547; measures against, 286.

Eriococcus coriaceus, bionomics of, on Eucalyptus in New Zealand, 752; in Australia, 752; natural enemies and biological control of, 753.

Eriococcus devoniensis, measures against, on bell heather in Britain, 296.

Eriophyes, on Salvia in California, 173; on Nephelium spp. in China and Formosa, 674; on couch grass in Queensland, 494.

Eriophyes cactorum, sp. n., or Opuntia in California, 650.

Eriophyes eriobotryae, sp. n., on Eriobotrya japonica in California, 650.

Eriophyes gossypii, on cotton in Porto Rico, 41; distribution and importance of, 41.

Eriophyes neosalviae, sp. n., on Salvia leucophylla in California, 650.

Eriophyes oleivorus (see Phyllocoptruta).

Eriophyes pyri, bionomics of races of, on fruit trees etc. in British Columbia, 171; on pear in California, 19, 748; measures against, 171, 748.

Eriophyes sheldoni, sp. n., on lemon

in California, 173. 650.

Eriophyes spinulifera, sp. n., in Cecidomyiid galls on Artemisia in California, 650.

Eriophyes tenuis, on couch grass in W. Australia, 494.

Eriophyes tristriatus erineus, on walnut and lemon in California, 650.

Eriophyes tulipae, sp. n., on imported tulip bulbs in California, 650.

Eriopis connexa (predacious on Toxoptera graminum), parasite of, in Argentina, 289.

Eriopyga cynica, on apple and peach in Indiana, 126.

Eriopyga incincta, bionomics and distribution of, in U.S.A., 135.

Eriosoma americanum, on elm and Amelanchier in U.S.A., 484.

Eriosoma crataegi, bionomics and status of, in N. America, 484, 485.

Eriosoma lanigerum (Woolly Apple Aphis), in Britain, 527; in Canada, 484, 485; in Caucasus, 74; in Cyprus, 108, 260; in Czechoslovakia, 657; in Germany, 2, 233, 247; in Haiti 44; in Line 347; in Haiti, 41; in India, 409; predacious enemies of, in Japan, 771; in Libya, 260; in Malta, 469; in Poland, 98; tentatively recorded on quince in Réunion, in Rumania, 192; Spain, 696; in U.S.A., 41, 65, 484, 485; biological and morphological differences between E. crataegi and, 484, 485; susceptibility of varieties of apple to, 2; utilisation of Aphelinus mali against, 41, 65, 74, 98, 108, 192, 233, 347, 469, 485, 696, 771; spraying against, 260.

Eriosoma rileyi, on elm in U.S.A., 484.

Eriosoma ulmi, on currant and elm in Finland, 201.

Eritrea (see Africa, Italian East). Ermine Moths (see *Hyponomeuta*). *Ernobius mollis*, measures against, damaging timber in Britain, **463**.

erosa, Cosmophila. errans, Anomala.

eruditus, Cheyletus.

Erycia basifulva, parasite of Tirathaba trichogramma in Fiji, 334. Erythmelus, parasite of Capsids in U.S.A., 328.

Erythmelus psallidis, sp. n., parasite of Psallus seriatus in U.S.A., 227. Erythrina, new weevil on, in Brazil, 106; Oxyplax ochracea on, in Ceylon, 639; pests of, in India,

149, 292; pests of, in Netherlands Indies, 292, 607; Ferrisiana virgata on, in New Guinea Territory, 495; new Bruchid on, 292; as shade against coffee mealybugs, 182.

erythrinae, Specularius. erythrocephala, Calliphora.

Erythroneura, new Crabro predacious on, in Ohio, 369.

Erythroneura comes, on vines in U.S.A., 19, 286, 323; measures against, 286, 323.

Erythroneura pallidifrons, on ornamental plants in Britain, 350.

Espartillo Grass (see Elionurus adustus).

Estonia, miscellaneous pests in, 462.

esuriens, Diaprepes.

Eterusia virescens, parasites of, on tea in India, 149.

Ether, for extracting toxic principles of plants, 36, 37, 183, 206, 258, 361, 547, 566, 614, 766. Ethyl Acetate, in baits for moths,

Ethyl Acetate, in baits for moths, 236; for extracting insecticidal principles of plants, 36, 258; ethyl oxyhydrate probably an impure form of, 137.

Ethyl Alcohol (see Alcohol).

Ethyl Oxyhydrate, in baits for Cydia pomonella, 137.

Ethylene Dichloride, toxicity of, to pests of stored cereals, 326; and carbon tetrachloride, books, etc., treated with, against termites, 223.

Ethylene Oxide, experiments with, against pests of stored products, 326, 413, 540, 571, 572; use of, against pests of stored grain in Algeria, 413; vacuum fumigation of potatoes with, against Leptinotarsa decembineata, 597; post-fumigation effects of, on Lasioderma serricorne, 396; curves for expressing toxicity of, to insects, 633; fire hazard of, 313, 413; and carbon dioxide (see also Cartox and T-gas), 396, 413, 540, 571, 572, 597; other fumigants compared with, 326, 389, 633.

Etiella zinckenella (attacking leguminous plants), in Porto Rico, 224; in shelter belts in Russia, 432; parasites imported into U.S.A. against, 329.

Euaresta aequalis, introduced into Australia against Xanthium pungens, 450.

Eucalymnatus perforatus, on cinnamon in Seychelles, 433.

INDEX.

Eucalymnatus tessellatus, on coconut

in Sevchelles, 433.

eucalypti, Rhicnopeltella; Rhinocola. Eucalyptus, Gonipterus scutellatus on, in S. Africa, 203; pests of, in Australia, 8, 86, 111, 209, 334, 752, 753, 754; Helopeltis antonii on, in India, 149; pests of, in New Zealand, 752-754; G. scutellatus on, in Nyasaland, 203.

Eucalyptus Weevil (see Gonipterus

scutellatus).

Eucelatoria australis, parasite of Anomis luridula in Peru, 627.

Eucolaspis brunnea, on Eucalyptus in New Zealand, 754.

Eucosma lechriaspis, bionomics of, on apple etc. in Japan and Korea,

Eucosma macropetana, on Eucalyptus in New Zealand, 754.

Eucosma ocellana (on apple), bionomics of, in Caucasus, 477; in Czechoslovakia,657; in New York, 562; on pear, 477; measures against, 478, 562.

Eucosma phylloscia, sp. n., on coffee in Uganda, 151.

Eucosma schistaceana (on sugarcane), in China, 243; in Formosa, 263, 673, 768; in Loochoo Islands, 437; susceptibility of varieties of sugar-cane to, 263; bionomics of, 673, 768; effect of submersion on, 673.

uderus gossypii, parasite of Sphenoptera gossypii in India, **89.** Euderus

Euderus pempheriphila, sp. n., parasite of Pempheres affinis in India,

Eudia pavonia, on poplar and willow in Germany, 461.

Euetheola bidentata, on sugar-cane in Br. Guiana, 369.

Euetheola humilis, on sugar-cane in Brazil, 502.

Eugenia spp., Cerambycids on, in Netherlands Indies, 295.

eugenii, Anthonomus.

Eugenol, tests of, in baits for Cydia, 135.

Eulachnus, producing honey-dew in Central Europe, 459.

Eulan, Plodia interpunctella not affected by, 460.

Eulecanium (see Lecanium).

Eulia mariana, in light-traps in orchards in Nova Scotia, 535.

Eulimneria crassifemur, parasite of Pyrausta nubilalis in Poland, 623. Eulimneria rufifemur, parasite of Pyrausta nubilalis in Poland, 623.

Eulophus, parasite of Cassida spp. in Italy, 528.

Eulophus longulus, parasite of Dacus oleae in Libya, 261.

damaging Eumaenas salaminia. papaya fruits in Queensland, 246.

Eumarchalia gennadii (on carob), bionomics and control of, in Crete, 71; in Cyprus, 458.

Eumicrosoma benefica, parasite of Blissus leucopterus in U.S.A., 78. Euonymus, Coccids on, in Louisiana, 41, 589.

Eupatorium pallescens, planted to favour Scoliids in Mauritius, 317.

Eupelmella vesicularis, bionomics of, attacking Microplectron fuscipenne in Hungary, 341; parasite of Mayetiola destructor in Morocco, 340.

Eupelmus, parasite of Apanteles americanus in Cuba, 171; parasite of Mayetiola destructor in Morocco, 340: parasite Anomis luridula in Peru, 627.

Eupelmus allyni, competing with Platygaster in Mayetiola destructor

in U.S.A., 121.

Eupelmus urozonus, associated with Eumarchalia gennadii in Crete, 71; parasite of Cydia funebrana in Switzerland, 100.

Euphorbia hypericifolia, Bemisia on, in Porto Rico, 41.

Euphorbia ligularia, Orthezia praelonga on, in Grenada, 752.

euphorbiae, Aphthona; Celerio (Deilephila).

Euphoria longana (see Nephelium). Euphranta mikado, on Camellia in Japan, 311.

Euphyllura olivina (on olive), in Libya, 260; in Spain, 700.

Eupista nigricella (see Coleophora). Euplectrus euplexiae, parasite of Spodoptera mauritia in India, 241.

Euplectrus platyhypenae (parasite of armyworms), in Mexico, 586: establishment of, in Hawaii, 586.

euplexiae, Euplectrus.

Euproctis chrysorrhoea, auct. (see Nygmia phaeorrhoea).

Euproctis flava, on castor and rose in Ceylon, 359.

Euproctis fraterna, tests of plant extracts against, on apple in Mysore, 361.

Euproctis minor, on Tephrosia purpurea in Java, 110, 111.

Euproctis producta (rubricosta), on cotton and wild castor in Sudan, 513.

Eupteromalus leguminis, parasite of Bruchus pisorum in U.S.A., 654.

Eupteromalus nidulans, an imported parasite of Lepidoptera in U.S.A., 589, 590.

Eupteromalus viridescens, parasite of Angitia polynesialis in Virginia, 13.

Eupterote geminata, on castor in Ceylon, 359.

Eupterote mollifera, on moringa in India, 55.

Eupteryx atropunctata, on potato in Germany, 68; not transmitting potato viruses, 69.

Europe, keys to leaf-mines caused by insects in, 247; key to Coleopterous forest pests in, 312; keys to store and household pests in, 196.

Eurydema oleraceum, on crucifers in Finland, 201.

Eurydema ornatum, bionomics and control of, on rape in Czechoslovakia, 718.

Eurydinoteloides longiventris, parasite of Gasterocercodes brasiliensis in Brazil, 504.

Eurygaster, bionomics of, on wheat in Spain, 698; types of injury to cereals by, 51, 92.

Eurygaster austriaca, bionomics of, on cereals in Switzerland, 92.

Eurygaster maura (on cereals), in Czechoslovakia, 659; bionomics of, in Germany and Switzerland, 51, 52, 92; parasites of, 92; characters of, 51.

Eurytetranychus stenoperitrematus (see Paratetranychus).

Eurytoma, possibly parasitic on Hylesinus aculeatus in New Jersey, 394.

Eurytoma appendigaster, hyperparasite of Cydia funebrana in Switzerland, 100.

Eurytoma atripes, parasite of Mayetiola destructor in Morocco, 340. Eurytoma curculionum, parasite of

Lixus junci in Morocco, 417.

Eurytoma dentata, associated with

Eumarchalia gennadii in Crete,
71.

Eurytoma samsonovi, on apricot in India, 765.

Eurytoma tristis, parasite of Lixus junci in Morocco, 417.

Eurytoma walshi, parasite of Apanteles americanus in Cuba, 171. eurytomae, Syntomaspis.

Eusarcocoris (Eusarcoris) guttiger, bionomics of, on rice in China, 245.

Eusattus difficilis, association of, with ants in Kansas, 292.

Euscelis striatulus, measures against, on cranberry in New Jersey, 508.

Euscepes postfasciatus (batatae), measures against, on sweet potato in Barbados, 222, 446–448; bionomics of, 447; synonymy of, 576.

Euschistus euschistoides, causing black pit and kernel spot of pecan in Florida, 587.

Eutachina larvarum (see Tachina). Eutermes, method of testing sus-

ceptibility of timbers to, in Australia, 449; damaging lawns in Tanganyika, 113.

Eutetrapha tridentata, parasites of, on elm in Connecticut, 14, 15.

Eutettix discigntus, bionomics and control of, transmitting mulberry disease in Japan, 208.

Eutettix tenellus (Beet Leafhopper), in U.S.A., 127, 128, 134, 282, 283, 322, 445, 691, 743; experiments with curly-top of beet and, 445; other crops attacked by, 127; vegetation types in relation to, 322, 691; migrations of, 283; natural enemies of, 127, 128; measures against, 134, 282.

Euthrips insularis (see Frank-liniella).

Eutinophoea bicristata, on Citrus in Queensland, 494.

Euxoa longidentifera, on tobacco in Tanganyika, 642.

Euxoa ochrogaster, on cereals in Saskatchewan, 170.

Euxoa segetum, in Czechoslovakia, 657; new type of pseudograsserie in, in France, 164; in Italy, 528; in S. Rhodesia, 90; in Russian Union, 73, 160, 236, 422, 423, 424, 432, 472, 473; in Tanganyika, 642; on beet, 236, 423, 472, 473, 528, 657; on other plants, 160, 424, 473, 642, 657; bionomics and physiology of, 73, 422, 423, 424, 440, 472, 473; parasites and biological control of, 90, 236, 528; baits for, 160, 236.

Euxoa vestigialis, on spruce seedlings in Czechoslovakia, 657.

evanescens, Trichogramma. Evetria (see Rhyacionia). examinator. Pimbla. excessana, Tortrix, exclamationis, Feltia.

exempta, Laphygma (Spodoptera).

Exenterus abruptorius, bionomics of, parasitising Diprion sertifer in Europe, 197, 198; imported into U.S.A. from Canada against D. polytomum, 329.

Exenterus adspersus, establishment of, against Diprion polytomum in

Canada, 487.

Exenterus cingulatorius (see E. abruptorius)

Exeristes (see Pimpla).

exitiosa, Aegeria (Conopia).

exigua, Laphygma (Caradrina). exiguae, Hyposoter.

Exochomus flavipes, predacious on Aphis laburni in French Sudan,

Exochomus quadripustulatus, predacious on Chermes nordmannianae in Germany, 661.

Exochomus ventralis, establishment of, in Seychelles against Coccids,

exoleta, Calocampa (Noctua). Exorista aethiopica (see Tachina). Exorista mitis (see Zenillia).

exotica, Allograpta.

Experimental Data, analysis of, 94, 141, 165, 196, 633.

exsul, Conoderus (Monocrepidius).

Extar, 25.

extensor, Ephialtes. externus, Acarapis.

Eye-spotted Apple Bud Moth (see Eucosma ocellana).

F.

fabae, Aphis (Doralis); Empoasca. faber, Ergates. fabia, Earias. faceta, Antestia. facialis, Empoasca. fagi, Phyllaphis. Fagopyrum tataricum, Chaetocnema concinna on, in Poland, 255. Fall Cankerworm (see Alsophila pometaria). Fall Webworm (see Hyphantria cunea).

fallax, Tachina.

falsa, Antestia. Fannia canicularis, in unsound tomatos in U.S.A., 398. farinae, Tyroglyphus (Aleurobius).

farinalis, Pyralis. fasciata, Bogosiella.

fasciatipennis, Coloborhombus (Nothopeus).

fasciatus, Cleonus (Chromoderus); Dysdercus; Hercothrips; Oncopeltus.

Lachnus fasciatus, Shinji,

Cinara shinjii).

INDEX.

fasciculatus, Araecerus (Araeocerus). Fatsia japonica var. moseri, mites

on, in Germany, 660.

Fatty Acids, possibly effecting inversion of arsenical spray mixtures, 174; wetting and spreading properties of mixtures of sodium hydroxide with, 320; identification of, combined with pyrethrolone, 694.

fausta, Rhagoletis. febriculosus, Merisus. febrilus, Dilophus.

exclamationis, Feltiameasures against, in forest nurseries in Russia, 432.

femoralis, Hercinothrips (Hercothrips, Heliothrips); Philodicus; Rhaphidopalpa.

femorata, Brachymeria; Chrysobothris; Diapheromera.

femur-rubrum, Melanoplus.

fenestralis, Angitia; Sciara (Neosciara).

Fergusobia, subgen. n., 86.

Fergusonina, association of Anguillulina with, on Eucalyptus in Australia, 86.

Ferminia fusca, in granaries in Russia, 95, 96; bionomics of, 96. Fernando Po, summary of plant

quarantines in, 134. Ferns, pests of, in Britain, 152, 712,

Ferric Hydrate, as treatment for arsenic poisoning in bees, 255.

Ferric Hydroxide (precipitated), as an adhesive for lead arsenate

sprays, 750.

Ferrisiana (Ferrisia) virgata, in Hawaii, 212; bionomics of, on coffee and Leucaena glauca in Java, 182, 606, 607; food-plants of, in New Guinea Territory, 495; in Panama, 212; on Inga in Porto Rico, 41; on Solanum arundo in Tanganyika, 757; new Encyrtid parasites of, 41. 212; measures against, 182.

Ferrous (Iron) Sulphate, uses of, in sprays, 481, 534.

ferruginatus, Bibio.

ferruginea, Coelinius (Coelinidea). ferrugineum, Tribolium (see castaneum).

ferrugineus, Dacus (Chaetodacus); Rhynchophorus.

ferus, Nabis.

festata, Plusia (Phytometra).

festina, Stictocephala.

festiva, Xystrocera.

Festuca glauca, new Coccid on, in Czechoslovakia, 560.

Festuca ovina, Aelia acuminata on, in Germany, 51.

Festuca rubra, Geomyza tripunctata on, in Britain, 382.

Ficus spp., Pseudaonidia on, in Japan, 296; insects on, in W. Indies, 41. (See Fig.)

ficus, Chrysomphalus.

Fidia viticida (Grape Rootworm), measures against, in U.S.A., 323.

Field Plots, co-ordination of experiments in, 141, 356, 630.

Fig, Pseudococcus nipae on, in Hawaii, 586; pests of, in India, 601, 765; Eutettix disciguttus on, in Japan, 208; Dacus oleae associated with, in Italy, 161; pests of, in Palestine, 406; pests of, in U.S.A., 292, 722; Ephestia figulilella breeding on fruit of, 324.

figulilella, Ephestia.

Fiji, coconut pests in, 211, 212, 334, 497; miscellaneous pests in, 58, 497, 702; biological control of noxious insects and weeds in, 58, 59, 87, 334, 497, 702; introduction of beneficial insects into other countries from, 87, 145, 204, 212, 703; summary of plant quarantines in, 385.

Fiji Disease, of sugar-cane, Perkinsiella saccharicida a vector of,

fijiensis, Opius.

Filbert (see Corylus). filiformis, Orthacris.

Filippia oleae, measures against, on

olive in Italy, 350.

Finland, miscellaneous pests in, publications 201, 518; entomology in, 120, 695; summaries of plant quarantines in, 201, 385.

Fir, Douglas (see Pseudotsuga taxi-

folia).

Fir, Silver or White (see Abies).

(see Cryptocephalus Fire-beetle incertus).

Firebrat (see Thermobia domestica). Fireworm, Black-headed (see Rhopobota naevana).

Fireworm, Yellow-headed (see Peronea minuta).

fiscellaria, Ellopia.

Fish, danger to, of derris dusts, 383, 529.

Fish Oil, repellent to ants, 223; as an adhesive and spreader for sprays, 399, 488, 547, 548, 552, 652, 723, 724, 725, 750; surface tension affecting toxicity of emulsions of, to Lepidosaphes gloveri, 702; formulae containing, 399, 548, 652, 723, 724.

Fish-oil Soap (see Soap, Fish-oil).

fiskei, Brachymeria.

Fixator-A, a form of bentonite, 554. 555, 733.

Flabellomicrosiphum, in Utah, 119. Flame-throwers, against Lepidoderma albohirtum, 151; against locusts, 611, 686.

flammea, Meliana; Panolis. flava, Cosmophila (Anomis); Euproctis (Nygmia); Hoplocampa; Sipha.

flavescens, Empoasca (Chlorita).

flavicollis, Calotermes.

flavidipennis, Lonchaea.

flavipes, Exochomus; Apion; Reticulitermes.

flaviscutellaris, Cyrtomorpha.

flaviventris, Apanteles.

flavo-orbitalis, Cremastus.

flavopicta, Hylemyia (Chortophila). flavostigma, Chrysopa.

flavovarius, Tetrastichus.

flavus, Stomatothrips; Tychius.

Flax (Linseed), Toxoptera graminum on, in Argentina, 289, 290; Dasyneura lini on, in India, 181, 242; Blapstinus substriatus on, in Montana, 27; flea-beetles on, in Hungary and Poland, 197, 255; pests of, in Asiatic Russia, 6, 308, 351, 474, 601; mite in seeds of, in Russia, 95.

Flax Weevil (see Ceuthorrhynchus sareptanus).

Flea-beetle, Hemp or Hop (see Psylliodes attenuata).

Flea-beetle, Tobacco (see Epitrix parvula).

Flea-hopper, Cotton (see Psallus seriatus).

fletcheri, Lecanium; Opius.

Flinders Island, Halotydeus destructor in, 582.

floccifera, Pulvinaria.

floralis, Nemorilla.

orida, new Anobiid infesting books in, 329; Citrus pests in, 325, 650, 727, 728; cotton pests Florida, new in, 41, 132; Coccus mangiferae

and its fungus parasite on mango in, 587; miscellaneous pests in, 727: Homaledra sabalella on palms in, 22; pests of pecan in, 587; pests of solanaceous plants in, 64, 132, 144, 217; sugar-cane pests in, 28, 137, 326, 390; Reticulitermes flavipes damaging turnips in, 285; Naupactus leu-coloma in, 131, 580; beneficial insects and biological control in, 22, 137, 390, 391, 727.

Florida Flower Thrips (see Frankliniella cephalica bispinosa).

Florida Red Scale (see Chrysomphalus ficus).

floridanus, Stigmaeus,

Flour, pests in, and their control, 30, 95, 96, 118, 130, 165, 311, 321, 326, 389, 571; effect of injury to grain by Pentatomids on quality of, 92, 521; absorption of fumigants by, 389; effect of grain fumigants on quality of, 414, 415; in baits, 40, 224, 705; carrier and adhesive for dusts, 211, 639, 652, 705, 736; as an adhesive for sprays, 16, 443; as an emulsifier for oils, 185, 187.

Flour Beetle (see Tribolium confusum).

Flour Beetle, Red (see Tribolium castaneum).

Flour Mills, fumigation of, 38, 118,

Fluorine Compounds, problem of residues of, on sprayed fruit and vegetables, 25, 62, 216, 327, 401. Fogweed (see Atriplex expansa).

foliae, Pimpla (Epiurus).

forbesi, Aspidiotus; Cremastus.

Forest Entomology, summaries and bibliographies of papers on, 154, 368, 625.

Forest Fires, relation of pests to, 80, 385.

Forest Tent Caterpillar (see Malacosoma disstria).

Forestit, methods of dusting with, against forest pests, 69, 235; apparently harmless to bees, 69.

Forests, pests of, in S. Africa, 68, 203; pests of, in Australia, 8, 86, 111, 209, 334, 449, 494, 582, 583 pests of, in Austria, 215, 659, 663, 664, 715; Melolontha hippocastani in, in Belgium, 193; pests of, in Bermuda, 220, 626; pests of, in Brazil, 145, 502; pests of, in Britain, 296, 527, 625, 631, 633, 711; pests of, in Burma,

338-340, 762, 765; pests of, in Canada, 77, 103, 215, 293, 330, 487, 509, 510, 534, 538, 539; pests of, in Czechoslovakia, 215, 657, 658, 659, 715; bark-beetles in, in Estonia, 462; pests of, in Finland, 201, 518; pests of, in France, 248, 316; pests of, in Germany, 2-4, 50, 69, 154, 189, 235, 251, 252, 301, 312, 344, 419, 660, 634, 664, 762; pests of, in Holland, 183, 299, 347, 348, 621, 656; pests of, in Hungary, 197, 198, 527; pests of, in India, 531, 763, 764, 765; pests of, in Indo-China, 185, 186, 187; pests of, in Japan, 265, 295, 296, 311, 437, 438, 452, 516, 604, 669; Lymantria dispar in, in Jugoslavia, 759: pests of, in Korea, 767; pests of, in Latvia, 90, 721, 722; pests of, in Malaya, 110, 453, 763; Dendrolimus spectabilis in, in Manchuria, 672; pests of, in New 752-754; Zealand, 58, 266, Gonipterus scutellatus in, in Nyasaland, 203; pests of, in Poland, 4, 154, 256, 523; Dikraneura cedrelae in, in Porto Rico, 288; new Coccids in, in Portugal, 315; pests of, in Rumania, 193. 721; pests of, in Russian Union, 73, 74, 75, 158, 304, 308, 422, 426, 428, 429, 430, 431, 432, 433, 530, 665; pests of, in Sakhalin, 516; Vesperus in, in Spain, 699; pests of, in Sweden, 197, 198, 611, 634; pests of, in Switzerland, 164, 459; termites in, in Tanganyika, 113; pests of, in U.S.A., 14, 15, 17, 18, 20, 80, 126, 136, 215, 225, 226, 270, 293, 325, 326, 329, 330, 332, 369, 386, 392, 394, 399, 441, 487, 542, 543, 546, 547, 548, 549, 550, 551, 576, 580, 581, 588, 650, 696, 723, 725, 745; book on pests of, in U.S.A., 400; key to Coleoptera injurious to, in Central Europe, 312; fauna of litter of, 193, 523; insectivorous birds in, and their protection, 429, 625, 633, 634; effect of defoliation of, on rainfall and floods, 385; (shade trees), discussion of insecticides for, 560.

Forficula auricularia, decrease of, in California, 18; toxicity of " Detal" dusts to, 345.

Formaldehyde (Formalin), toxicity of, to cheese mites, 539; fumigation of mushroom houses with, against pests, 579.

Formica fusca, attacking Eutettix disciguttus in Japan, 208.

Formica rufa, attacking Panolis flammea in Germany, 235.

formicarius, Cylas.

Formosa, banana pests in, 263; pests of Citrus in, 207, 438, 674; cotton pests in, 309; miscellaneous pests in, 53, 208, 310, 517, 769; sugar-cane pests in, 263, 294, 673, 768, 769; new Aphids in, 119; Melaphis chinensis producing tannin galls in, 264; food-plant preferences of Coccids of, 262; Dacus spp. in, 206, 207, 263, 436, 517, 667; injurious mites of, 436, 674; Prodenia litura in, 207; beneficial insects in, 119, 207, 674, 768; Dermestid parasitic on Mantid eggs in, 514; introduced toad destroying Bufo melanostictus in, 317.

formosa, Encarsia.

formosanus, Phassus (see P. signifer);

Termes (Odontotermes).

fornicata, Phytodecta.

fornicatus, Xyleborus.

Fornicia ceylonica, parasite of Limacodids in Ceylon, 639; parasite of, **639**.

foveicollis, Cleonus; Rhaphidopalpa (Aulacophora).

Fowls, destroying noxious insects, 61, 181, 128, 132, 520, 658.

fragaefolii, Capitophorus.

fragariae, Capitophorus; Tylo-

derma.

France, Pegomyia spp. on beet etc. in, 314; orchard pests in, 192, 259, 343, 416, 464, 525, 526; miscellaneous pests in, 156, 229, 301; pests of sycamore in, 316; vine moths in, 415, 619; pests of stored products in, 95, 412, 414, 415, 501, 523; termites in building timber in, 314; Coccids of, 248, 303, 696; Leptinotarsa decemlineata in, 5, 368, 412, 415, 594, 595, 597; beneficial insects and biological control in, 415, 524, 594, 595; utilisation of bacteria against *Polychrosis bo*trana in, 619; pseudograsserie in Euxoa segetum in, 164; new species of Nosema attacking C. pomonella in, 464; summaries of plant quarantines in, 229, 692, 751.

Frankliniella, factors affecting damage to cotton by, in U.S.A.,

277.

Frankliniella cephalica bispinosa, measures against, on Citrus and ornamental plants in Florida, 727.

Frankliniella fusca, on cotton in U.S.A., 277, 644; measures

against, 644.

Frankliniella insularis, F. lycopersici recorded as, in S. Australia, 318; food-plants of, in W. Indies, 175, 222; measures against, 222.

Frankliniella lycopersici, sp. n., food-plants of in Australia, 318; vector of spotted wilt of tomato,

318.

Frankliniella paucispinosa, measures against, transmitting virus disease of tobacco and tomato in Argentina, 581.

Frankliniella robusta (see Kako-

thrips pisivorus).

Frankliniella runneri, on cotton in

Mississippi, 277.

Frankliniella tritici, in U.S.A., 277, 278, 286, 644; on cotton, 277, 278, 644; measures against, 286, 644.

Frankliniella vaccinii. measures against, on blueberry in Maine, 333.

Frankliniella williamsi, on cotton in S. Caroline, 278.

fraterculus, Anastrepha.

fraterna, Euproctis; Phragmatiphila (Nonagria) typhae.

fraxini, Duplaspis; Hylesinus; Stereonychus (Cionus).

fraxinifolii, Prociphilus.

Fraxinus (see Ash).

pests of, in excelsior, Fraxinus Czechoslovakia, 658; new Coccid on, in France, 248.

French Colonies, summaries of plant quarantines in, 385, 692.

French Ray Grass (see Arrhenatherum).

frischi, Dermestes.

frisius, Cylindroiulus (see C. oweni).

frit, Oscinella.

froggatti, Brontispa (see B. longissima); Typhlocyba.

frontalis, Aenasius; Dendroctonus; Platymetopius.

frugiperda, Laphygma.

Fruit (Dried), pests of, 30, 324, 389, 449, 461, 573, 599.

Fruit-flies (see Trypetids).

Fruit-fly, Cherry (see Rhagoletis cerasi).

Fruit-fly, Mediterranean (see Ceratitis capitata).

Fruit-fly, Melon (see Dacus cucurbitae).
Fruit-fly, Mexican (see Anastrepha ludens).
Fruit-fly, Olive (see Dacus oleae).
Fruit-tree Red Spider (see Paratetranychus pilosus).
frutetorum, Diprion; Holocremnus. fugax, Solenopsis. fulgidus, Scelio. fuliginosum, Gymnosoma. fuliginosus, Coeliodes; Lasius. fullawayi, Diachasma (Opius). fullo, Polyphylla.

fullonia (fullonica), Othreis (Ophideres). fulvicollis, Lema. fulvicornis, Lema.

fulvifrons, Zodion. fulvum, Stromatium.

Fumaria (Fumea) pinicola, sp. n., on Pinus patula in Transvaal, **68.** fumiferana, Harmologa (Cacoecia).

Fumigants, apparatus for determining absorbtion by, and toxicity to insects of, 567, 741; curves for expressing time-concentration relations of, 633.

Fumigation, general accounts of, against pests of stored grain, flour, etc., 38, 202, 321; types of equipment for, 20, 21, 39, 85, 116, 118, 176, 362, 389, 519, 593, 663, 695.

fumipennis, Hercothrips; Microdontomerus.

funebrana, Cydia (Grapholitha, Laspeyresia).

funeralis, Desmia. funesta, Oxythyrea.

Fungi, Entomogenous, 39, 46, 78, 81, 87, 101, 114, 123, 136, 146, 147, 152, 153, 179, 193, 239, 264, 294, 348, 370, 374, 375, 378, 395, 420, 433, 434, 456, 468, 504, 557, 587, 590, 591, 607, 642, 659, 681, 682, 688, 715.

Fungi, Injurious, 110, 119, 350, 406, 588, 625, 635, 644, 666, 707, 733, 744; relation of insects to, 20, 28, 43, 60, 178, 191, 249, 325, 375, 376, 385, 386, 454, 457, 540, 547, 551, 587, 632, 634, 656, 657, 658, 707, 749, 751, 753, 757.

Fungus, Édible, Scatopse fuscipes on, in Japan and Korea, 438. (See Mushrooms.)

furfura, Chionaspis.

Furniture, pests in, and their control, 190, 400, 591.

Fusarium, infesting Aspidiotus perniciosus, 715.

Fusarium coccinellum, infesting Aonidiella aurantii in S. Africa, 456.

fusca, Busseola; Ferminia (Glycyphagus); Formica; Frankliniella.

fuscipenne, Microplectron. fuscipennis, Thrips.

fuscipes, Rhynocoris; Scatopse. fuscispinus, Carpocoris pudicus.

G.

gabonator, Phosphorus. gahani, Pseudococcus.

Galerucella distincta, bionomics of, on strawberry in Japan, 771.

Galerucella luteola (xanthomelaena) (Elm Leaf Beetle), in Switzerland, 164; in U.S.A., 18, 126; measures against, 18, 164.

gallarum, Byrsocrypta.

Galleria mellonella (Large Wax Moth), destroying combs of honeybees in Brazil, 655; Chelifer cancroides predacious on, in Germany, 664; race of Nemeritis canescens parasitising, in U.S.A., 1; nutrition of larvae of, 184.

Gallobelicus (see Engytatus).

Gallobelicus crassicornis (see Engytatus tenuis).

Gambia, food-plants of *Diparopsis* castanea in, 199; summary of plant quarantines in, 229.

Gambrus stokesi, introduced into U.S.A. from Australia against Cydia spp., 450.

gamma, Plusia (Phytometra).

Ganaspis, parasite of Ceratitis spp. in Uganda, 492.

Garden Centipede (see Scutigerella immaculata).

Gardinol, contact performance of, 45; composition of, 45.

gasteralpha, Pulvinaria (see P. iceryi).

Gastrallus laticollis, damaging books in Netherlands Indies, 706.

Gasterocercodes brasiliensis, sp. n. (on cotton), in Argentina, 503; in Brazil, 115, 502-504, 505; bionomics of, 502-504; measures against, 504, 505.

Gasterocercodes gossypii, G. brasiliensis recorded as, in Brazil, 115; on Malachra in Peru, 533.

Gastrimargus musicus, bionomics and distribution of, in Australia, 449, 584.

Gastropacha quercifolia, on apple in Britain, 613.

Gaylussacia baccata, Cryptocephalus incertus on, in Massachusetts, **646**.

gei, auct., Macrosiphum (see M. solanifolii).

geisha, Vanessa io.

Gelatine, in baits for ants, 223; improving suspension of homologues of Paris green, 285.

Gelechia malvella (see Platyedra).

gelechiae, Microbracon.

Gelis laricellae, sp. n., parasite of Coleophora laricella in Austria, 660.

gemellatus, Silvanus.

geminata, Atacta; Eupterote; Solenopsis.

Solenopsis.

geminatella, Parornix (Ornix). geminatus, Dyscinetus (see D. dubius).

gemini, Phaulacridium. gemmatalis, Anticarsia.

Gendarussa vulgaris, insecticidal properties of, **361.**

geniculata, Fieb., Cyrtopeltis. geniculatum, Copidosoma.

geniculatus, Reut., Engytatus (Cyrtopeltis).

Geniocerus rapo (see Tetrastichus).

Genista, **51, 234.** genitalis, Phorbia.

gennadii, Eumarchalia (Asphondylia, Schizomyia).

Geocoris spp., predacious on Eutettix tenellus in Utah, 128.

Geomyza combinata, G. tripunctata confused with, in Britain, 382.

Geomyza tripunctata, bionomics of, on cereals and grasses in Britain, 382.

Georgia (U.S.A.), Cydia pomonella in, 30; Epitrix parvula on tobacco in, 132; Pseudococcus boninsis on sugar-cane in, 137; beneficial insects and biological control in, 30, 137.

Georgia Clay, as a carrier for cubé, 133.

Geranium, Hepialus lupulinus on, in Switzerland, 689. (See Pelargonium roseum.)

Germalus pacificus, attacking Teleonemia lantanae in Fiji, 87.

germanica, Blattella.

germanus, Xyleborus (Xylosandrus). Germany, beet pests in, 314, 419, 461, 520, 717; Depressaria nervosa on caraway in, 719; cereal pests in, 51, 52, 251, 521, 619, 717; forest pests in, 2-4, 50, 69,

154, 189, 233, 234, 235, 251, 252, 301, 312; 344, 346, 419, 461, 633, 660, 664, 665, 717, 762; pests of hazel in, 234, 560; pests of hops in, 522, 523; lucerne pests in, 461, 520; miscellaneous pests in, 52, 152, 248, 468, 576; onion pests in, 39; orchard pests in, 2, 50, 52, 100, 101, 156, 194, 233, 234, 250, 251, 346, 347, 420, 421, 522, 660, 716, 717, 718, 719, 761, 762; insects found in fruit-tree bands in, 689; pests of small fruits in, 194, 234, 344, 660: pests of ornamental plants in, 52, 101, 248, 496, 522, 660, 717; Leptinotarsa decemlineata potato in, 120, 252, 347, 368, 418, 719; other potato pests in, 68, 461, 520, 717; Bostrychids boring in lead sheathing of cables in, 232; pests of stored products in, 37, 38, 48, 95, 164, 300, 347, 439, 460, 461, 519, 662; pests of timber in, 48, 91, 195, 233, 300, 301, 599, 662; vine pests in, 4, 154, 419, 521, 601, 661, 664; ants associated with Aphids in, 195; review of larvae of Chrysomelids in, 716; Galleria mellonella in bee-hives in, 664; Melolontha spp. in, 250, 251, 253, 345, 346, 468, 520, 599, 717; other injurious Lamellicorns in, 194, 717; millepedes in, 461, 468; beneficial insects and biological control in, 196, 233, 234, 235, 347, 420, 523, 576, 620, 661, 662, 684, 763; entomogenous fungi in, 101, 152, 419, 468; pests intercepted in quarantine in, 101 plant pest legislation in, 91, 229, 719, 751; standardisation of tar distillates in, 99; list of officially tested insecticides in, 636; legislation dealing with poisonous fumigants in, 101, 347; history of HCN fumigation in, 120.

gerstaeckeri, Platypus.

Gherkin, Halticus saltator on, in Germany, 52.

ghesquierei, Agrilus.

Gibbium psylloides, food of, in buildings in Germany, 519.

gibbus, Bruchophagus.

giffardi, Ceratitis; Hedylus;

Opius; Tetrastichus. giffardianus, Tetrastichus.

gifuensis, Macrocentrus.

giganteus, Ceroplastes. gigas, Pantophthalmus; Sirex. Gilbert Islands, summary of plant quarantines in, 134.

Ginger, pests of, in Ceylon, **359**; (stored), pests in, in Jamaica, **491**.

Gisekia pharnaceoides, Laphygma exigua on, in India, 55.

glabrata, Ametastegia.

glabrisetus, Peritetranychus.

gladioli, Taeniothrips (see T. sim-

plex).

Gladiolus, Taeniothrips simplex on, in Bermuda, 626; Pseudococcus maritimus on, in Hawaii, 213; pests of, in U.S.A., 65, 398, 552; insecticides causing injury to, 398, 399, 552, 553.

Gladiolus Thrips (see Taeniothrips

simplex).

glaucus, Papilio.

gleadowi, Brontispa.

Glenea cantor, biomomics and control of, on kapok in Indo-China, 186.

Gliricidia, Frankliniella insularis on, in St. Vincent, 175.

Gliricidia maculata, Myllocerus discolor on, in Ceylon, 358; relation of Diaprepes spp. to, in St. Lucia, 402.

globata, Microgaster.

Globitermes (see Amitermes).

glomeratus, Apanteles.

Glossonotus crataegi, measures against, on apple in New York, 269, 574.

gloveri, Lepidosaphes.

Glue, uses of, in arsenical sprays, 285, 548; improving suspension of homologues of Paris green, 285.

Glycerine, facilitating emulsion of extracts of Melia azedarach, 180.

glycinae, Phalaenoides.

glycinivorella, Cydia (Grapholitha). glycinivorellae, Microdus; Pimpla (Epiurus).

Glycyphagus cadaverum, not found in Russia, 95.

Through a garage

Glycyphagus destructor (see Lepidoglyphus).

Glycyphagus domesticus, in stored cereals etc. in Russian Union, 95, 96.

Glycyphagus fuscus (see Ferminia). Glycyphagus michaeli (see Lepidoglyphus).

Glycyphagus ornatus, in stored cereals in Russian Union, 95.

Glyphodes unionalis (see Marga-ronia).

Glypta (?) pedata, parasite of Cydia molesta in Italy, 154.

Glypta rufiscutellaris (parasite of

Cydia molesta), alternative hosts of, in Ontario, 509, 510; introduced into Italy and Australia from U.S.A., 154, 448.

Glypta simplicipes, parasite of Tortrix rosaceana in Utah, 141.

Glyptomorpha deesae, parasite of sugar-cane borers in India, 243, 408, 532; bionomics of, 532.

Glyptomorpha nicevillei, bionomics and value of, in Madras, 408.

Glyptomorpha smenus, parasite of Sphenoptera gossypii in Mysore, 204.

Gmelina arborea, pests of, in Burma, 339.

Gnathocerus cornutus, susceptibility of, to derris, 183.

Gnophothrips piniphilus, sp. n., on pines in U.S.A., 329.

Gnorimoschema (see Phthorimaea).
Goa, new Bruchid on Erythrina in,
292.

Godetia, Haltica oleracea on, in Poland, 255.

godmani, Asynonychus (Pantomorus).

Gold Coast, new parasite of Retithrips aegyptiacus in, 610; summary of plant quarantines in, 691. goldingi, Bemisia.

Golf Greens (see Turf).

Gonia cinerascens (see Cnephalia). Goniozus, associated with cotton pests in Mexico, 275.

Gonipterus scutellatus (Eucalyptus Weevil), in S. Africa, 203; in New Zealand, 753; in Nyasaland, 203; parasites and biological control of, 204, 753.

Gooseberry, new mite on, in Central Asia, 530; pests of, in U.S.A., 60, 281; Lymantria dispar not surviving on, 759.

Gooseberry Maggot (see Epochra canadensis).

gorhami, Epicauta.

Goryphus nursei, parasite of Clania crameri in Madras, 499.

Gossyparia spuria, measures against, on elm in U.S.A., 18, 126.

gossypiella, Platyedra (Pectino-

phora).

gossypii, Aphis; Bucculatrix; Eriophyes; Euderus; Gasterocercodes; Heterospilus; Monolepta (Candezea); Phenacoccus; Sphenoptera.

gossypiperda, Bemisia. Gossypium (see Cotton). gotama, Mycalesis.

Goulac (see Sulphite Lye).

gowdeyi, Haplothrips.

Gracilaria theivora, parasites of, on tea in India, 149.

gracillimum, Monomorium.

graellsi, Acontia (Xanthodes).

Grain Bug, Say's (see Chlorochroa sayi)

Grain Moth, Angoumois (see Sitotroga cerealella).

Grain Moth, Lesser (see Rhizopertha dominica).

Grain Weevil (see Calandra granaria).

Gram (see Cicer arietinum).

graminum, Toxoptera.

granaria, Calandra (Sitophilus).

granarium, Macrosiphum (see M. avenae; Trogoderma.

grandicollis, Ips.

grandis, Anthonomus; Lebia: Synonycha.

Grape Leaf-folder (see Desmia funeralis)

Grape Leafhopper (see Erythroneura comes)

Grape Rootworm (see Fidia viticida).

Grape-berry Moth (see Polychrosis viteana).

Grape-vine Thrips (see Rhipiphoro-

thrips cruentatus).

Grapefruit, termite attacking, in California, 219; fruit-piercing moths associated with fungi on, in Dominica, 751; Anastrepha ludens on, in Mexico, 325; pests of, in Palestine, 406, 636; heat treatment of fruit of, against Dacus dorsalis in Formosa, 207.

Grapes, compulsory sterilisation of imported, against Ceratitis capitata in U.S.A., 229; juice of, in baits for Trypetids, 108; Ephestia figulilella on mummified, 449.

Graphium ulmi (see Ceratostomella). Graphocephala coccinea, on narcissus

in U.S.A., 288.

Graphocephala versuta, on cotton in Texas, 278.

Graphogaster maculatus, sp. parasite of Psalidium maxillosum in Russia, 94.

Grapholitha (see Cydia).

Graptosaltria nigrofuscata, measures against, on pear in Japan, 604.

Grass, Blue (see Poa).

Grass, Couch (see Agropyrum repens). Grass, French Ray (see Arrhenatherum).

Grass, Rye (see *Lolium*). Grass, Timothy (see Pheum pratense).

Grass Mealybug (see indica)

Grasselli Spreader-Sticker, 735, 736. Grasses, pests of, in Australia, 111, 209, 267, 449, 494, 582, 584, 703; pests of, in Britain, 341, 382; Cirphis unipuncta on, in Canada, 145, 441: new Coccid on, in Czechoslovakia, 560; Lamellicorns on, in Germany, 194, 717; pests of, in Loochoo Islands, 437. 516; Aphids on, in Porto Rico,

39, 40; pests of, in Russian Union, 257, 425, 426; pests of, in U.S.A., 15, 23, 44, 78, 122, 127, 139, 140, 226, 279, 282, 580, 589, 723; relations of locusts and grasshoppers to, 23, 44, 139, 140, 147, 162, 198, 282, 426, 494, 584, 680, 681, 683, 703, 769; damaged by termites, 113, 449; cereal and sugar-cane pests on, 39, 40, 51, 63, 67, 76, 135, 218, 240, 241, 290, 294, 326, 360, 427, 428, 577, 659, 728. (See Turf.)

Grasshopper, Clear-winged Camnula pellucida).

Grasshopper, Dark-winged (see Stauroderus scalaris).

Grasshopper, Egyptian (see Anacridium aegyptium).

Grasshopper, Siberian (see Aeropus sibiricus).

Grasshoppers (see Locusts).

Greece, Dacus oleae on olive in, 71; Bostrychids boring in lead sheathing of cables in, 232; list of pests in, 70; summary of plant quarantines in, 134.

Greenhouse Fumigation, 355, 488, 626, 711.

Greenhouse Red Spider (see Tetranychus telarius).

Greenhouse Thrips (see Heliothrips haemorrhoidalis).

Greenhouse Whitefly (see Trialeurodes vaporariorum).

gregalis, Blapstinus (see B. substriatus).

gregaria, Schistocerca.

Grenada, miscellaneous pests in, 752; mealybug parasite from, 212.

gressoria, Sitona.

Gretchena bolliana, on pecan in Florida, 587.

Grey-streaked Moth (see Prodenia litura).

Greyback Beetle (see Lepidoderma albohirtum).

grioti, Charips (Xystus).

grisator, Sthenias.
grisea, Sitona.
grisella, Achroia (Meliphora).
griseoalba, Acanthopsyche.
grisescens, Lydella stabulans.
griseus, Xylotoles (Xyloteles).
Ground-nut Oil (see Oil, Ground-nut).

Ground-nuts, Aphis laburni and virus diseases of, in E. and W. Africa, 114, 239, 531, 592; Heliothis armigera on, in S. Africa, 364; Prodenia litura on, in Formosa, 207; pests of, in U.S.A., 132, 609; review of literature on insects and diseases of, 386; methyl bromide not impairing germination of, 389.

grylloides, Hysteropterum.
Gryllotalpa, baits for, in Egypt, 457.
Gryllotalpa africana, light-traps of doubtful value for, in Egypt, 457; in forest nurseries in Korea, 767.

Gryllotalpa gryllotalpa, in Egypt, 457; on beet in Italy, 528; on vegetable seedlings in Malta, 469; in forest nurseries in Russia, 432; measures against, 432, 457, 528.

Gryllotalpa gryllotalpa var. cophta, in Egypt, 457.

Gryllulus assimilis, on melon and cotton in California, 748.

Gryllulus desertus, baits for, on beet in Italy, 528.

Gryllulus domesticus, bionomics of, in Germany, 37.

Gryllus (see Gryllulus).

Guatemala, beneficial insects in, 493, 692; summary of plant quarantines in, 134; Pyrophorus bellamyi introduced into Hawaii from, 493.

guatemalensis, Dicrodiplosis.

Guava (Psidium guayava), Aonidiella aurantii on, in S. Africa, 455; Dacus spp. on, in Ceylon, Bonin Islands and Fiji, 359, 672, 702; pests of, in India, 55, 361, 765; Pseudococcus nipae on, in Mexico, 586; D. cucurbitae on, in Philippines, 181; new weevil on, in St. Lucia, 405; Apate monacha on, in Sierra Leone, 240; Helopeltis bergrothi on, in Sudan, 513.

gudimiri, Prionomerus. guerini, Rodolia (Novius).

guexi, Synaphaeta.

Guiana, British, pests of sugar-cane and rice in, 221, 369, 491; coconut pests in, 369, 491; beneficial insects and biological control in, 221, 369, 491; summaries of plant pest legislation in, 134, 491; rotenone-yielding plants in, 319; Metagonistylum minense introduced into Trinidad from, 10.

Guiana, Dutch, rotenone-yielding plants in, 319; pests imported into Holland in butter-nuts from, 529; summary of plant quaran-

tines in, 229.

Guinea, Spanish, summary of plant quarantines in, 134.

gularis, Aphomia (Paralipsa). Gulf Wireworm (see Heteroderes laurenti).

Gum Copal, Manila, in banding mixture against ants, 93.

gurgistanus, Agriotes. gurnevi, Coccophagus.

Gutierrezia, Macrosiphum on, in Utah, 247.

guttata, Parnara.

guttiger, Eusarcocoris (Eusarcoris). guttulosa, Austracris.

Gymnosoma fuliginosum, parasite of Chlorochloa sayi in Montana, 44. Gymnosoma rotundatum, parasite of Aelia rostrata in Spain, 698.

Gynaikothrips uzeli, new parasite of, on Ficus nitida in Porto Rico, 41.
Gynerium, Aphids on, in Porto Rico, 40.

Gypsum, as a carrier for dusts, 399, 534, 537, 652, 726, 735; talc preferable to, as a carrier for pyrethrum, 726.

Gypsy Moth (see *Lymantria dispar*). Gyranusa (parasite of mealybugs), attempted introduction of, into Hawaii from Mexico, **586**.

H.

Haber's Formula, 633.

Habrobracon (see Microbracon).

Habrocytus lixi (parasite of Lixus spp.), in Italy, 528; bionomics of, in Morocco, 417.

Habrocytus thyridopterigis, hyperparasite of Thyridopteryx in Illinois, 173.

Habrolepoidea tarsalis, parasite of Bruchus brachialis in Pennsylvania, 580.

Hadena didyma (see Trachea secalis). Hadrobregmus destructor, sp. n., in wood in museums in Alaska, 214.

Haematopinus suis, Opilo domesticus predacious on, in Germany, 662.

Haematozylon campechianum, Orthezia praelonga on, in Grenada, 752.

Haemoglobin, as a spreader for oil emulsions, 280.

haemorrhoidalis, Heliothrips.

haeussleri, Phaeogenes.

Haiari, identity and constituents of plants known as, in S. America, 319.

Haiti, parasites and biological control of insects in, 41, 223; summary of plant quarantines in, 385. halli, Lepidosaphes.

Halotydeus destructor, bionomics and control of, in Australia, 85, 150,

582.

Haltica ampelophaga, tests of rotenone dusts against, on vines in Algeria, 371.

Haltica oleracea, on Godetia and strawberry in Poland, 255.

Haltica pagana, measures against, on strawberry in Tasmania, 209. Haltica torquata, on blueberry in Maine, **333.**

Halticus saltator, food-plants of, in

Germany, 52.

Halyomorpha picus, on Citrus in Ceylon, 359.

hamada, Castalius.

hamatus, Schistoceros.

hambletoni, Heterospilus; Rhabditis.

Hambletonia pseudococcina, ported into Porto Rico against Pseudococcus brevipes, 329.

Hamitermes (see Amitermes).

hampei, Stephanoderes.

Hapalia machaeralis, parasites and biological control of, on teak in India and Burma, 339, 688, 764, 765.

Haplophyton cimicidum, extracts of, in bait-sprays for Anastrepha

spp., 505.

aculeatus, attacking Haplothrips eggs of Cecidomyiid on wheat in Japan, 670.

Haplothrips gowdeyi, possibly a vector of pineapple yellow spot in Hawaii, 213.

hargreavesi, Brachyurothrips.

Harmologa fumiferana, on Pinus ponderosa in U.S.A., 580.

Harmologa oblongana, on apricot and plum in New Zealand, 58.

Harmonia axiridis (see Coccinella). Harpalus, attractiveness of cereals to, in Crimea, 356.

Harpalus rufipes, in Britain, 384, in Finland, 201; 763:

strawberry, 201, 384; outbreak of, 763; measures against, 384, 763.

harpax, Sarcophaga (Thyrsocnema). harti, Aspidiotus.

harukawai, Dolerus.

Harvester Termite (see Hodotermes).

hastulifer, Agrilus.

havilandi, Calotermes (Cryptotermes). Hawaii, fruit-flies in, 84, 213, 325, 752; miscellaneous pests in, 213, 584, 586, 692; pests and diseases of pineapple in, 143, 213, 570, 692; sugar-cane pests in, 454, 493, 585, 589; Megamelus proserpina on taro in, 212; vegetable pests in, 213, 214, 325, 692, 752; beneficial insects and biological control in, 84, 143, 212, 213, 214, 454, 492, 493, 497, 586; introduction of beneficial insects into other countries from, 59, 137,

hawaiiensis, Kentronothrips Podothrips lucasseni); Taenio-

thrips.

Hawthorn (see Crataegus).

Hazel (see Corylus).

273, 359, 627.

Heat, uses of, for treating timber against *Hylotrupes* and *Lyctus*, 155, 190, 379, 707; fruit treated with, against Trypetids, 207, 325, 751: against mushroom pests, against pests of stored products, 8, 31, 160, 455, 539, 768; susceptibility of eggs of Bruchus obtectus to, 284; affecting germination of seeds, 8; regulations for treatment of cottonseed with, against pests in U.S.A., 134, 229. (See Solar Heat, Steam and Water, Hot.)

Heather, Lochmaea suturalis on, in

Britain, 418.

Heather Beetle (see Lochmaea suturalis).

hebetor, Microbracon (Habrobracon).

hederae, Aspidiotus.

Hedylus giffardi, parasite of Ceratitis punctata in Sierra Leone, 492: introduced into Hawaii, 492.

Heinsia pulchella, coffee Lycaenid on, in Sierra Leone, 239.

Helianthus annuus (see Sunflower). Helianthus ciliaris, Isophrictis on, in Mexico, **275**.

Helicobia australis, not a parasite of grasshoppers in Australia, 629.

Helicteres isora, alternative host of parasite of *Hapalia* on, in India. 765.

helioba, Phthorimaea.

Heliothis, on tobacco in U.S.A., 393. Heliothis armigera, in S. Africa, 152, 364, 511; in Argentina, 362; in Belgian Congo, 315; in Hawaii, 752; in India, 409; in Nyasaland, 367; in Queensland, 494; in N. and S. Rhodesia, 366; in Russia, 472; in Sudan, 511; in U.S.A., 19, 65, 117, 142, 144, 271, 304, 326, 327, 489, 509, 564, 643, 644, 744, 745; on cotton, 142, 315, 362, 364, 366, 367, 409, 472, 494, 511, 512; on maize, 65, 144, 271, 304, 326, 364, 366, 489, 509, 511, 564, 643, 644, 745; resistant strains of maize against, 326; transport of, in maize ears, 564; on leguminous plants, 152, 327, 364, 366, 509, 511; on tomato, 19, 117, 364, 644, 752; relative attractiveness of different plants to, 144, 250, 364, 366, 494; bionomics of, 364, 366, 472; metabolism of, 440; cotton nectar as adult food of, 197; Trichogramma parasitic on, 744: measures against, 65, 142, 152, 271, 327, 362, 489, 494, 509, 511, 745, 752; pH and buffer value of blood of, 386; characters of, 117.

Heliothis assulta, measures against, on tobacco in Indo-China, 185.

Heliothis obsoleta (see H. armigera). Heliothis phloxiphaga, food-plants of, in California and Vancouver, 117, 118, 293; characters of, 117. Heliothis virescens, cubé of little value against, on tobacco in U.S.A., 133.

Heliothrips femoralis (see Hercinothrips).

Heliothrips haemorrhoidalis, on orange in California, 508; on tea in Nyasaland, 114; foodplants, of, in Russian Union, 354, 358; bionomics of, 354; measures against, 354, 508.

Heliotropium undulatum, attraction of, for Schistocerca gregaria, 628.

Helix aspersa, establishment of Lampyris noctiluca against, in Australia, 450.

Hellula undalis (on crucifers), bionomics of, in Japan, 605; in Virginia, 13.

hellulae, Angitia (see A. polyne-sialis).

Helopeltis, measures against, on tea in India, 148.

Helopeltis antonii, food-plants of, in India, **55**, **149**; measures against, **55**.

Helopeltis bergrothi, in Belgian Congo, 315; in Nigeria, 666, 667; in Nyasaland, 114; in Sudan, 513; in Uganda, 376; on cotton, 315, 376, 513, 667; type of injury by, 513; varietal susceptibility of cotton to, 376; on other plants, 114, 513.

Helopeltis sanguineus, on cotton in

Belgian Congo, 315.

Helops, soil conditions in relation to, in Russia, 429.

Helorimorpha, probably parasitic on Antestia in Tanganyika, 179.

helxines, Crepidodera (Chalcoides). hemerobiella, Coleophora.

Hemerocampa oslari, on Abies in California, **581.**

Hemerocampa pseudotsugata, on Pseudotsuga taxifolia in N. America, 293, 581; urticating hairs of, 293.

Hemerophila pariana (see Anthophila).

Hemiliopsis aequalis, sp. n., in Panama and Colombia, 405.

Hemiliopsis spondias, sp. n., on Spondias lutea in Brazil, 405.

hemipterus, Coloborhombus (Nothopeus); Hemiteles.

hemisphaerica, Saissetia (Lecanium) (see S. coffeae).

Hemiteles albipalpus var. austriacus, n., parasite of Coleophora laricella in Austria, 660.

Hemiteles areator, parasite of Cydia molesta in Italy, 154.

Hemiteles hemipterus, probably hyperparasite of Cydia funebrana in Switzerland, 100.

Hemiteles inimicus (with var. longisetosus), hyperparasite of Cydia funebrana in Switzerland, 100.

Hemiteles pulchellus, parasite of Coleophora laricella in Austria, 660.

Hemiteles tricoloripes, reared from Cydia molesta in Italy, 154; hyperparasite of C. funcbrana in Switzerland, 100.

Hemithea costipunctata simplex, on Tephrosia purpurea in Java, 110.

Hemlock (see Tsuga).

Hemp, pests of, in Poland 622, 719; pests of Russian Union, 159, 308, 470, 475, 476, 477.

Hemp Flea-beetle (see Psylloides attenuata).

heparana, Tortrix (Pandemis).

Hepialus lupulinus, on geraniums in Switzerland, 689.

Heptamelus ochroleucus, habits of, on ferns in Britain, 152.

heracleana, Depressaria. herbsti, Chlorophorus.

Hercinothrips bicinctus, on banana

in Queensland, 494.

Hercinothrips femoralis, on banana, etc., in Bermuda, 626; experiment with curly-top virus and, in California, 445.

Hercothrips fasciatus, food-plants and control of, in California, 19,

Hercothrips femoralis (see Hercinothrips)

Hercothrips fumipennis, on cotton and Crotolaria in Sudan, 513.

Hercothrips sudanensis, on cotton in Sudan, 513.

herculeanus, Camponotus.

herrichi, Animula

herricki, Platygaster.

Herring Oil, 702. (See Fish Oil.) Herse cingulata, measures against,

on sweet potato in Jamaica, 145. hesperidum, Coccus (Lecanium). hesperus, Lygus; Reticulitermes.

Hessian Fly (see Mayetiola destruc-

laurenti, Heteroderes

measures against, on cotton and potato in U.S.A., 327.

Heterogenea asella, measures against, on maple and lime in Russia, 433.

Heterographis bengalella, measures against, on Anona squamosa in Indo-China, 185.

Heteronychus licas, Carabid predacious on, on sugar-cane and maize in Natal, 411.

Heteronychus plebejus, bionomics of, in Madagascar, 66, 455.

Heteronychus tristis, on sugar-cane and maize in Natal, 411.

Heterorrhina elegans, on Citrus in

Ceylon, 359.

spp., parasites odes brasiliensis of Heterospilus Gasterocercodes Brazil, 504; hosts of, in Connecticut, 14; bionomics of, 14, 504.

Heterotoma meriopterum, oviposition of, on currant in Germany, 234.

Heterotylenchus aberrans, gen. et sp. n., associated with Hylemyia antiqua in Denmark, 49.

Heterusia (see Eterusia).

Hevea (see Rubber).

Hexachlorethane, use of, against Heliothis armigera, 745.

Hexamermis, parasite of grasshoppers in U.S.A., 124.

Hexamitodera semivelutina, on clove trees in Celebes, 295.

hibisci, Onebala.

Hibiscus spp., relation of Dysdercus spp. to, in Africa, 365, 366, 367, 374, 512; alternative food-plants of other cotton pests, 83, 243, 309, 503, 511, 513, 690, 766; Hypomeces squamosus on, in Indo-China, 186.

Hibiscus calycinus, Diparopsis castanea unable to mature on, 199.

Hibiscus cannabinus, effect of environment on infestation of, by insects and fungi, 666.

Hibiscus sabdariffa, pests of, in Netherlands Indies, **358.**

Hibiscus tiliaceus, Anomala xanthopleura on, in Loochoo Islands, 516.

hicoriae, Trigonura.

Hidari irava, new parasite of, in Malaya, 688.

hiemalis, Platygaster.

Hieracium, Nodonota puncticollis on, in New York, 288; Paroxyna tessellata on, in Ukraine, 619.

Hieroglyphus annulicornis, bionomics and control of, in Formosa, 769.

Hieroglyphus banian, Mermithid parasite of, on rice in India, 55.

Hieroglyphus banian var. elongata, on sugar-cane in Bombay, 261.

hilare, Acrosternum. Hippiscus californicus, in California,

19. hippocastani, Melolontha.

Hippodamia convergens, in U.S.A., 128, 575; predacious on Macrosiphum onobrychis, 128; massing of, on mountains, 575; technique for rearing, on frozen Aphids, 649; tests of insecticides on, 390.

Hippodamia parenthesis, predacious on Macrosiphum onobrychis

in Utah, 128.

Hippodamia quinquesignata, predacious on Macrosiphum onobrychis in Utah, 128.

Hirsutella, infesting Teleonemia lantanae in Fiji, 87.

Hirsutella exoleta, synonymy and hosts of, 152.

hirta, Epicometis; Oemona.

hirticornis, Otiorrhynchus telus); Thripobius.

Lachnosterna (Phyllohirticula, phaga).

hirtiventris, Lachnosterna (Phyllo-phaga).

Hispa armigera, measures against, on rice in China, 498; on other plants, 498.

Hispaniella howardi, sp. n., parasite of Parlatoria in Haiti, 41.

Hister americanus, predacious on Tyloderma fragariae in Kentucky, 737.

Histiogaster entomophaga (see Monieziella).

Histiostoma sapromyzarum, attacking Gryllulus domesticus in Germany, 38.

Hockeria unicolor, parasite of Lixus scabricollis in Italy, 528.

Hodotermes, bionomics and control of, in S. Africa, 48.

Hodotermes mossambicus, damaging grasses in Tanganyika, 113.

Hofmannophila, considered distinct from Borkhausenia (q.v.), 412. Hog Plum (see Spondias lutea).

Holarrhena antidysenterica, relation of, to parasite of teak pests in India, 765.

Holland, Paratetranychus pilosus on apple etc. in, 464, 465; instructions for spraying fruit trees in 343: Contarinia nasturtii on crucifers in, 466; forest pests in, 183, 299, 347, 348, 529, 621; Scolytids and elm disease in, 656; Leptinotarsa decem-lineata in, 120; miscellaneous pests in, 299; pests of mush-rooms in, 655, 656; pests of stored products in, 183, 599; pests imported into, in butternuts from Dutch Guiana, 529; beneficial insects in, 348, 465, 529, 576; new mite possibly imported into California on tulip from, 650.

Holly, Pulvinaria floccifera on, in France and Switzerland, 156; Phytomyza ilicicola on, in U.S.A., 541, 546.

Holly Leaf-miner (see *Phytomyza ilicicola*).

Hollyhock (see Althaea rosea).

Hollyhock Seed Moth (see *Platyedra malvella*).

Holochlora nawae, bionomics of, in China, 244.

Holocremnus frutetorum, parasite of Diprion pini in Holland, 348. hololeuca, Porthesia (see Acyphas

leucomelas). hololeucus, Niptus. holosericea, Maladera. Holotrichia (see Lachnosterna).

Homaledra sabalella, bionomics and control of, on palms in Florida, 22.

Homalodisca triquetra, on cotton in Texas, 278.

Homalopoda cristata, parasite of Ceroplastes giganteus in Haiti, 41.

Homona coffearia (Tea Tortrix), in Ceylon, 337, 360, 639; in India, 149; in Java, 639; parasites and biological control of, 149, 337, 639.

Homorocoryphus vicinus, swarm movements of, in Belgian Congo, 687.

Honduras, summary of plant quarantine restrictions in, 134.

Honduras, British, summaries of plant quarantine restrictions in, 134, 491, 692, 751.

Honey, in bait for ants, 202; in bait-spray for Trypetids, 619.

Honey-dew, importance of Aphids producing, in Central Europe, 459, 460.

Hongay (see Pongamia glabra).

Hoop Pine (see Araucaria cunning-hami).

Hop Flea-beetle (see Psylliodes attenuata).

Hopkins Host-Selection Principle, 1. Hoplocampa (on plum), measures against, in France, 5; in Palestine, 514

Hoplocampa flava (Plum Sawfly), measures against, in Britain, 191, 614, 708; bionomics of, 708.

Hoplocampa pyricola, warmth accelerating development of hibernating, 54.

Hoplochelus rhizotrogoides, bionomics of, on sugar-cane in Madagascar, 66, 67.

Hops, Epipolaeus caliginosus on, in Britain, 613; pests of in Germany, 522, 523; Vanessa io geisha on, in Japan, 673; pests of, in Poland, 622, 623, 719; Psylliodes attenuata on, in Ukraine, 477.

Hordeum pusillum, Eriopyga incincta on, in U.S.A., 135.

Horismenus apantelivorus, parasite of Apanteles americanus in Cuba, 171.

Horismenus cockerelli, parasite of Apanteles americanus in Cuba, 171.

Horistonotus uhleri, naphthalene ineffective against, in S. Carolina, 327.

Horned Squash Bug (see Anasa armigera).

Hornet (see Vespa orientalis).

hortensis, Chaetocnema.

horticola, Anomala (Phyllopertha).

hortuellus, Crambus.

hortulanus, Bibio.

hottentotta, Villa (Anthrax).

House Cricket (see Gryllulus domesticus).

House Longhorn Beetle (see Hylotrupes bajulus).

House-flies (see *Musca domestica*). Household Pests, key to, in Central Europe, 196; control of, 247, 248, 347.

hova, Cuphocera.

howardi, Dysdercus; Hispaniella;

Scello.

howitti, Aphodius.

hubbardi, Ataxia.

Huckleberry, Black (see Gaylussacia baccata).

huebneri, Hyalarcta.

huegeli, Dorysthenes (Lophosternus); Earias.

humeralis, Dacus (Tridacus).

humerella, Sophronia.

Humidity, relations of mites to, 74, 96, 97, 159, 160, 193, 396, 422, 465, 470, 471; relations of locusts to, 6, 72, 105, 114, 147, 410, 416, 583, 591, 678, 679, 681, 755, 756; relations of other insects to, 27, 37, 38, 61, 72, 113, 152, 158, 159, 165, 178, 190, 193, 195, 209, 219, 231, 232, 235, 236, 251, 302, 351, 352, 354, 356, 373, 374, 375, 380, 381, 399, 422, 423, 424, 425, 426, 446, 454, 457, 458, 474, 475, 476, 487, 511, 515, 519, 528, 532, 545, 546, 561, 595, 598, 607, 608, 624, 627, 637, 657, 661, 662, 668, 670, 671, 707, 709, 725, 758, 761.

humilis, Euetheola (Ligyrus); Iridomyrmex; Opius.

humuli, Phorodon.

Hungary, Cydia pomonella in, 254; Melož attacking bees in, 344; miscellaneous pests in, 197, 343; parasites of insect pests in, 341, 527, 576, 589.

hunteri, Catolaccus.

hvacinthi, Rhizoglyphus.

Hyalarcta huebneri, on trees in New South Wales, 334; spray against, 335.

hyale, Colias.

hyalinalis, Psammotis.

hyalinata, Diaphania (Margaronia).

hyalinipennis, Oxycarenus.

Hyaliodes vitripennis, predacious on Typhlocyba pomaria in New York, 557.

Hyalomya cuthbertsoni, parasite of Dysdercus in S. Rhodesia, 90.

Hyalopterus arundinis (pruni), bionomics and control of, in California, 62; on plum in Finland, 201; on peach in Italy, 189, 758; on almond in Libya, 260; predacious enemies of, 189.

Hybalus cornifrons, bionomics and control of, on cereals in Sicily,

757.

Hybernia defoliaria (see Erannis). Hyblaea puera, factors affecting biological control of, on teak in India, 764, 765; in Burma, 765. Hydnocarpus wightiana, pests of, in Ceylon, 359.

Hydrangea, thrips on, in Sierra

Leone, 240.

Hydrochloric Acid, for removing spray residues from fruit, 140, 318, 402, 646; addition of

Vatsol to, 141. Hydrocyanic Acid. greenhouse fumigation with, against tomato leaf-miner, 711; against pests of nursery stock, plant material and fruit, 306, 307, 593, 663, 747, 750; against mushroom pests, 579; soil fumigation with, 579; soil fumigation with, against *Tipulo aino*, 54; tent fumigation with, 18, 53, 70, 85, 107, 116, 117, 176, 177, 202, 456, 542, 663, 700, 702, 703, 707; dosage charts and methods for applying, 70, 86, 117, 176, 202, 702, 703; against pests of stored products, 24, 322, 326, 396, 514, 515, 654, 741; problem of penetration of, into flour, 322, 389; vacuum fumigation with, 24, 593, 741, 750; ineffective against Stephanoderes hampei in seed coffee, 359; against Coleoptera in furniture and building timber, 155, 190, 591; tests of toxicity of, to insects, 304, 326, 633, 741; sorption of, by various insects, 567, 741; strains of Aonidiella aurantii resistant to, 324; methylene blue distinguishing Coccids killed by, 770; post-fumigation effects of, on Lasioderma serricorne, 396; plants injured by, 54, 117; instrument for analysing mixtures of air and, 439; history of fumigation with, in Germany, 120 : other fumigants compared

with, 389, 747. (See Calcium and Sodium Cyanides and Zyklon.)

Hydrogen Phosphide, as a fumigant, 38, 101, 519; forms and dangers of. 101.

Hydrogen Sulphide, as a fumigant, 160, 481.

Hydrogen-ion Concentration, soil, relation of insects to, 425, 429, 430: of insects and insecticides, effect of, on toxicity of insecticides, 386, 690, 741, 742.

Hvdrol, in spray formula, against

gladiolus thrips, 65.

Hydroxymercurichlorophenol, unsatisfactory against Phorbia brassicae, 647.

hyettus, Aenasius. hylas, Cephonodes.

Hylastes ater, practices against, on pine in Sweden, 634.

Hylastes cunicularius, practices against, on pine in Sweden, 634. Hylastes obscurus, on clover in Utah, 126.

Hylastes palliatus, on Pinus sylves-

tris in Estonia. 462.

Hylastes rufipes, on elms in U.S.A., 14, 15, 325, 551, 576; a vector of Dutch elm disease, 325, 551; parasites of, 14, 15.

Hylastinus (see Hylastes).

Hylecoetus cossis, on conifers in

Japan, **517**.

Hylemyia antiqua (Onion Fly), in Britain, 298; new Nematode associated with, in Denmark, 49; in Germany, 39; in Ohio, 399; in Ontario, 536; bionomics of, 298, 399; measures against, 298.

Hylemvia brassicae (see Phorbia).

Hylemyia brunnescens, on carnation in Britain, 349.

Hylemyia cilicrura (see Phorbia).

Hylemyia flavopicta, bionomics and control of, on crucifers in Sakhalin, 770; in Japan, 770.

Hylemyia seneciella, introduced into Australia to destroy ragwort, 450.

Hylesia nigricans, new parasite of, in Argentina, 772.

Hylesinus fraxini, intercepted in Vancouver in crates from Britain, 483; studies on nutrition of, 631.

Hylesinus oleiperda, measures against, on olive in Spain, 700.

Hylesinus vestitus (Pistachio Bud Borer), in Sicily, 119.

Hylobius abietis (on pines), in Germany, 312; in Latvia, 90, 722: in Sweden, 634: measures against, 634, 722.

Hylobius pinastri, practices against,

on pine in Sweden, 634.

Hylobius radicis, bionomics of, in Connecticut, 18.

Hylotrupes bajulus (in building timbers), in Britain, 191, 378; in Germany, 48, 91, 195, 599, 662; in Sweden, 155; in Turkey, 530; bionomics of, 91, 155, 195; Clerid predacious on, measures against, 155, 379.

Hylurgopinus (see Hylastes).

Hylurgops (see Hylastes).

Hymenarcys nervosa, type of injury to peach by, in Delaware, 444.

hymeniae, Cremastus (see C. flavoorbitalis).

hyoscyami, Pegomyia.

Hyoscyamus niger, Aphids and virus diseases of, in Britain, 32, 342, 615; Ptinus tectus feeding on dried, in Germany, 164.

Hypera, measures against,

lucerne in Russia, 76.

Hypera meles, on high buildings in Ū.S.A., **215.**

Hypera punctata, on high buildings in U.S.A., 215.

Hypera variabilis (on lucerne), in Central Asia, 238; in Holland, 299; bionomics and spread of, in U.S.A., 18, 326, 747; natural enemies and biological control of, 18, 238.

Hyperaspis (predacious on Pseudococcus nibae), in Mexico, 586: introduced into Hawaii, 586.

Hyperaspis silvestrii (predacious on Pseudococcus nipae), in Mexico, 586; introduced into Hawaii,

Hyperaspis trilineata, imported into Porto Rico against sugar-cane

mealybugs, 223.

Hyperaspis usambarica, established against Coccids in Seychelles, 433. Hypericum perforatum, insects intro-

duced into Australia to destroy, 450.

Hyphantria cunea, parasites of, in Canada and U.S.A., 133, 291; on pecan in Florida, 587; tests of insecticides on, 388.

Hypocassida subferruginea,

beet in Italy, 528.

Hypodiranchis, parasite of Meromyza americana in Kansas, 123. hypogaea, Diarthronomyia.

Hypogastrura armata, measures against, on mushrooms Holland, 656.

Hypomeces squamosus, bionomics of, on kapok etc. in Indo-China, 186; measures against, on cotton in Java, 766; toxicity of extracts of Stemona to, 206.

Hyponomeuta padellus (including malinellus or apple race), bionomics of, and legislation respecting, in Cyprus, 107, 259; in Czechoslovakia, 656; in Germany, 50; in Italy, 712, 713; in Ukraine, 159, 302; on apple, 107, 159, 259, 302, 656, 712, 713; on plum, 50, 259; parasites of, 203; parasites of, 250, 250; parasites of, 250; par 302; measures against, 50, 259, 713.

Hypopteromalus, associated with Hylesinus aculeatus in New Jersey, 394.

Hyposerica castanea, on cassava in

Madagascar, 455.

Hyposoter exiguae, parasite of armyworms in Hawaii, 586.

Hyposoter pilosulus (parasite of Hyphantria), oviposition and sex ratio in, in Colorado, 291.

Hypoxylon sertatum, 46.

Hypsipyla robusta, on Cedrela toona in Burma, 340; on mahogany in India, 765; in Queensland, 502; parasites of, 502, 765.

Hysteroneura setariae (see Carolinaia).

Hysteropterum grylloides, on olive in Spain, 700.

Hystrix patula, new Coccid on, in New York, 226.

I.

Ialine, soil treatment with, against Euscepes batatae, 447.

ianthes, Dichomeris. ianthinus, Aenasius.

Icerya pilosa, Coccinelid predacious on, in India, 243.

Icerya purchasi, imported into Cyprus, 107; food-plants of, in Formosa and Australia, 263; in Libya, 260; on Aleurites in Nyasaland, 114; in W. Indies, 223, 491; on Citrus, 107, 491; utilisation of Rodolia cardinalis against, 223, 260, 491; symbiont of, 560.

Icerya seychellarum, natural enemies of, on fruit trees in Seychelles,

433.

icervoides, Phenacoccus.

Ichneumonids, list of, in Germany, 420; prepupal stage in, 197, 198.

Idaho, biological control of apple pests in, 65, 323, 747; Bruchus pisorum and its natural enemies in, 65, 653, 654; miscellaneous pests in, 65, 581, 747; Aphids in, 369, 696; vegetation types in relation to Eutettix tenellus etc. in, 322, 691.

Idatiella, key to species of, 247.

Idatiella albisignata, sp. n., on apple in New Zealand, 247.

Idiocerus (Mango Hoppers), toxicity of plant extracts to, in Mysore, 361.

Idioderma varia, on Citrus in Jamaica, 490.

idiota, Pentodon.

Igepon T, as a spreader, 547, 548.

Ilex (see Holly).

ilicicola, Phytomyza. illepida, Argyroploce.

Illinoia (see Macrosiphum).

Illinois, cereal pests in, 124, 125, 391; Gossyparia spuria on elm in, 126; orchard pests in, 125, 544, 545, 587, 588, 652, 653, 734; termite in building timbers in, 748; Mermis subnigrescens parasitising honey-bee in, 746; insect parasites of pests in, 173, 330, 369, 544, 653.

imbili, Stomatoceras.

immaculata, Echthromorpha; Scutigerella.

implicita, Lachnosterna (Phyllophaga).

impressicolle, Alissonotum.

impressifrons, Clivina.

Imperata arundinacea, Alissonoimpressicolle associated tum with, in Assam and Burma, 294.

inaequalis, Coelophora. Inareolata (see Angitia).

incanus, Brachyderes.

inceptaria, Itame.

Schoenobius incertellus, (see S. bipunctifer).

incertus, Cryptocephalus; Zatropis.

incincta, Eriopyga.

inconsequens, Taeniothrips. inconspicua, Sturmia.

inconspicuella, Sturmia. inconspicuoides, Sturmia.

Incurvaria rubiella, on loganberry in Britain, 297.

indagatrix, Pimpla (Epiurus).

Indarbela dea, bionomics and control of, in Assam, 601, 602.

Indarbela quadrinotata, bionomics and control of, in Assam, 601, 602. India, cotton pests in, 41, 55, 89, 261, 408, 409, 666; pests of flax and Cajanus in. 181, 242, 451; forest pests in, 68, 531, 763, 764, 765; relation of undergrowth to biological control of teak pests in, 764, 765; fruit pests in, 51, 87, 242, 340, 409, 602, 765; miscellaneous pests in, 241, 409, 497, 592; Pyralids on Moringa in, 55, 405; rice pests in, 55, 240, 360; sugarcane pests in, 8, 55, 241, 243, 532; pests of tea and its shade trees in, 148, 149, 693; locusts in, 112, 628, 629, 666, 677, 678, 683, 686; insects and virus diseases of plants in, 55, 88, 89, 242; book on lac cultivation in, 311; beneficial insects in, 55, 88, 89, 149, 241, 242, 243, 532, 764, 765; catalogue of Chalcidoidea of, 440; Dirhinus auratus introduced into Ceylon from, 359; Bruchid imported into Uganda from, 610. (See also under the various Provinces.) Moth (see Indian Meal Plodiainterpunctella). Indiana, cereal pests in, 126, 326, 580, 730; Cydia pomonella on apple in, 126, 320, 442, 652, 732; Hylastes rufipes and Dutch elm disease in, 325; miscellaneous pests in, 126. indica, Antonina. indicata, Lamprosema. indicum, Syntomosphyrum. indicus, Neocatolaccus. Indigofera cordifolia, effect of, on development of Schistocerca gregaria, 628. Indigofera endecaphylla, Lycaenid on, in Java, 110; Dichomeris ianthes on, in Seychelles, 433. Indo-China, miscellaneous pests in. 184, 185, 206, 261; pests of kapok and rubber in, 185-187; bionomics of Prodenia litura in, 205; termites in, 186, 187, 188. Indo-Malayan Region, keys genera of Dactylopiinae in, 607. indobrassicae, Siphocoryne (see Rhopalosiphum pseudobrassicae). infausta, Aglaope. inferens, Sesamia. inferior, Miselia. infuscatellus, Chilo. Inga spp., insects on, in Porto Rico, 41, 223; injured by thallium.

223.

inimica, Polyamia. inimicus, Hemiteles; Pantomorus. Inocarpus edulis, pests of, in Fiji, 497. inops, Sitona. Inostemma, parasite of wheat Cecidomviid in Japan, 670; key to Palaearctic species of, 526. Inostemma contariniae, sp. n., parasite of Contarinia medicaginis in Hungary, 527. Inostemma opacum, parasite of Contarinia medicaginis in Sweden, inquisitor, Calosoma; Pimpla. Insecticides, methods of testing, 36, 135, 253, 276, 297, 312, 321, 388, 405, 536, 572, 575, 610; organic compounds as potential, 524, 575, 576, 651; bibliographies on materials of vegetable origin for, 694; retention of, from emulsions and emulsionsuspensions, 370; relation of pH etc. of insects to toxicity of, 386, 690, 741, 742; affecting action of light on plants, 525; standardisation of, in European countries, 98, 99, 155, 636; publications and patents relating to, in U.S.A., 48, 304; general papers on, 533, 560, 651, 695, 703. Insects, influence of, on conservation of natural resources, 385; effect of environment on infestation of crops by, 666; nutrition of wood-eating, 248; action of dusts desiccating, 252; method for recording cardiac activity in, 439; preparation of specimens of, 183, 357; Dermestids infesting specimens of, 250. insidiator, Acanthostoma. insidiosus, Orius. insignis, Axionicus; Closterocerus; Orthezia. insulana, Earias. insulare, Melitomma. insularis, Aenasi's; Frankliniella (Euthrips). intacta, Scirpophaga. integerrima, Datana. integra, Lineodes. intercessor, Microbracon (Bracon). intermedia, Brachymeria. intermedius, Dysdercus. interpunctella, Plodia. interruptor, Cremastus.

interstitialis, Baris.

intricatus, Scolytus.

io, Vanessa.

Iowa, Blissus spp. in, 225, 283; pests of Gladiolus in, 65; grasshoppers in, 23, 123; Lachnosterna spp. in, 225.

iowensis. Blissus.

Ipobracon, parasite of Platyedra gossypiella in Brazil, 704; in Peru, 391.

Ipobracon rimac (parasite of Diatraea saccharalis), in Peru, 390, 391; establishment of, in U.S.A., 391.

Ipobracon tucumanus (parasite of Diatraea saccharalis), in Argentina, 391; introduced into U.S.A. 391.

Ipomoea, Cylas formicarius on, in Loochoo Islands, 669.

Ipomoea batatas (see Sweet Potato). Ipomoea biloba, Euscepes batatae on, in Barbados, 446.

Ipomoea cathartica, thrips migrating to banana from, in Bermuda, 626.

Ipomoea cordofana, cotton pests on, in Sudan, 512, 513.

ipomoeae, Blosyrus.

Îps amitinus, on conifers in Estonia, 462.

Ips bidentatus, on pine in Estonia, 462.

Ips calligraphus, aseptic rearing of, 28.

Ips chalcographus, on conifers in Estonia, 462.

Ips confusus, oil sprays against, on pines in California, 394.

Ips duplicatus, on conifers in Estonia, 462.

Ips grandicollis, associated with blue-stain fungi on Pinus resinosa in Minnesota, 386; aseptic rearing of, 28.

Ips pini, associated with bluestain fungi on Pinus resinosa in

Minnesota, 386.

Ips proximus, on pine in Estonia, 462.

Ips quadridens, on pine in Estonia,462.Ips typographus, on spruce in

Sweden, **634.**Iran (see Persia).

Iraq, summary of plant quarantines in, 692.

irava, Hidari.

iridescens, Levuana.

iridis, Bregmatothrips.

Iridomyrmex humilis, measures against, in California, 129; relation of, to Aphids in greenhouses in Germany, 195. Iris, Rhadinoceraea reitteri on, in Italy, 598; Cnephasia conspersana on, in Scilly Isles, 527; thrips on, in U.S.A., 273; insecticide causing injury to, 274.

Iris Thrips (see Bregmatothrips iridis).

Iron Sulphate (see Ferrous Sulphate).

irrorata, Anticarsia; Aulacizes.

irroratus, Phlepsius.

Isaria exoleta (see Hirsutella).

Isaria farinosa, hosts and identity of, 152.

Ischnaspis longirostris, biological control of, on coconut in Seychelles, 433.

Ischnodemus saccharivorus, bionomics and control of, in Loochoo Islands, 437.

Iseropus (see Pimpla).

ishiharai, Chermes.

Isoberlinia doka, Lyctus in timber of, in Sudan, 514.

Isocrates vulgaris (see Asaphes).

Isoglyphus, new subgenus of Caloglyphus (q.v.), 94.

Isophrictis, on Helianthus ciliaris in Mexico, 275.

Isostasius punctiger (parasite of Contarinia tritici), in Germany, 620; in Sweden, 155, 231; bionomics of, 231.

Italian Locust (see Calliptamus italicus).

Italian Millet (see Setaria italica). italicus, Calliptamus.

Italy, beet pests in, 161, 184, 461, 527, 528; pests in chestnuts for export to U.S.A. from, 384; Citrus pests in, 70, 713; review of research on elm disease in, 119; sawfly on Iris in, 598; Cecidomyiid on lucerne in, 189; pests of maize and Sorghum in, 528, 712, 713; miscellaneous pests in, 713, olive pests in, 161, 261, 350; orchard pests in, 5, 154, 189, 350, 593, 594, 636, 712, 713, 714, 758; Macrosiphum solanifolii transmitting mosaic of solanaceous plants in, 759; locusts in, 162, 163, 592; Acarapis on honey-bees in, 343; Dermestid attacking silkworm eggs and pupae in, 153; Coccids of, 496; Pemphigini of poplars in, 696; beneficial insects and biological control in, 153, 154, 163, 189, 410, 527, 528, 593; fungus infesting Orgyia antiqua in, 152; summary of plant quarantines in, 691; introduction of other beneficial insects into countries from, 154, 233; probable spread of Cydia molesta into Switzerland from, 163.

Itame inceptaria, on blueberry in New Hampshire, 20.

Ithone mirabilis (see Aiolocaria). Ithycerus noveboracensis, on peach in Connecticut, 17.

Itonida seminis (see Cecidomyia). Itoplectis (see Pimpla).

Jacaranda, termite damaging, in Tanganyika, 113.

jaceae, Macrosiphum. jacobaeae, Tyria.

Jamaica, termites attacking building timbers in, 220; miscellaneous pests and biological control in, 145, 490; search for natural enemies of Tomaspis in, 369; summary of plant pest legislation in, 491.

Lachnosterna; Projamaicensis, toparce sexta.

janata, Achaea. janus, Aspongopus.

Japan, Crioceris spp. on asparagus in, 497; pests of Camellia in, 311, 438; pests of crucifers in, 605, 769, 770; forest pests in, 265, 295, 296, 311, 437, 438, 452, 516, 604, 669, 673, 771; pests of fruit trees in, 53, 54, 296, 368, 438, 604, 671, 673, 771; list of tree pests in, 208; pests of Juncus in, 672; pests of leguminous plants in, 53, 264, 265, 310, 605, 669, 673; miscellaneous pests in, 265, 266, 438, 605, 669, 673; pests of mulberry in, 208, Pyrausta phoenicealis on 604; Perilla in, 771; rice pests in, 53, 54, 167, 265, 309, 439, 669, 670, 671, 672, 770, 771; Coleoptera on strawberry in, 603, 771; Cecidomyiids on wheat in, 670; pests of stored products in, 514, 668, 770; termite in imported timber in, 55; insects transmitting plant diseases in, 167, 208, 265, 439; Aleurodids of, 496; Aphids of, 452, 496; Melaphis chinensis producing tannin galls in, 264; foodplants of adult Lamellicornis in, 54, 516, 605; beneficial insects and biological control in, 54, 108, 142, 208, 295, 296, 439, 515, 516, 605, 669, 670, 671, 672, 771; species of silkworms and their insect enemies in, 247, 310, 514, 515; Dermestid parasitic on Mantid eggs in, 514; risk of establishment of Dacus cucurbitae in, 667, 668; summary of 692: plant quarantines in, Rhizoglyphus echinopus imported into Formosa from, 674; Thripoctenus brui introduced into Hawaii from, 143.

Japanese Beetle (see Popillia japo-

Japanese Empire, Braconids of, 247. Japanese Mandated Islands, Coccids in, 48; Hispids on palms in, 336; Trypetid apparently introduced into Bonin Islands from, 672.

Japanese Plum (see Prunus salicina). japonensis, Cryptotympana.

japonica, Aserica (Autoserica); Athalia; Delomerista; Dictyoploca; Popillia; Vanessa (Pyrameis) cardui.

japonicum, Trichogramma.

japonicus, Chermes; Macrocentrus; Semanotus.

Jarring, to dislodge pests from trees, 186, 193, 621, 690, 708, 727. Jassids, of N. America, 367.

javana, Brontispa (see B. longissima).

javanica, Tessaratoma.

javanus, Plaesius. Javel Water, tests of, against cheese mites, 539.

Jeffrey Pine (Pinus jeffreyi). jeffreyi, Dendroctonus.

Jelutong (see Dyera costulata).

jezoensis, Castalius hamada; Dendrolimus.

jonesi, Platyphytoptus. Jowari (see Sorghum). jucundus, Aphelinus.

juglandis, Acrobasis. Juglans regia (see Walnut).

Jugoslavia, Lymantria dispar in, 759; Polychrosis botrana 720; Phytodecta fornicata in, 760; losses and expenditure due to locusts in, 685; beneficial insects in, 192, 759, 760; summaries of plant quarantines in, 134, 229, 385, 692.

Julodis caillaudi, on date palm in Sudan, 513.

junci, Lixus.

juncivorus, Tomostethus.

Juncus effusus decipiens, pests of,

in Japan, 672.

jungi, Austroicetes (see A. cruciata). Juniper (Juniperus), Dichomeris marginella on, in Britain and Czechoslovakia, 296, 657.

Juniperus bermudiana, pests of, in

Bermuda, 220, 626.

Juniperus chinensis, Coccid on, in Bermuda, 626.

junodi, Acanthopsyche. juvenilis, Ceromasia.

K.

Kaburagia rhusicola, gen. et sp. n., in galls on Rhus from China, 265.

kaempferi, Platypleura.

Kainit, against Eurydema ornatum, 718; ineffective as a repellent for Melolontha, 346.

Kakothrips pisivorus, on peas in Britain, 527.

Kale, Meligethes aeneus on, in Britain, 296.

Kalo, 735, 736. (See Cryolite.) Kalotermes (see Calotermes).

kampalana, Dimmockia.

Kansas, pests of cereals and grasses in, 121-123, 135, 728; Psallus seriatus on cotton in, 136; miscellaneous pests in, 184; beneficial insects in, 123, 136; Enaresta aequalis introduced into Australia from, to destroy Xanthium, 450.

Kaolin, as a carrier for dusts, 133,

197, 268, 755.

Kapok, cotton stainers on, in Africa, 374, 512; pests of, in Indo-China, 185, 186, 187; new Coccid on, in Java, 607.

Karaya Gum, form of nicotine bentonite prepared with, 555.

kariyai, Apanteles.

Kentronothrips, a subgenus Podothrips, 457.

hawaiiensis Kentronothrips Podothrips lucasseni).

Kentucky, Tyloderma fragariae on strawberry in, 736; Epitrix parvula on tobacco in, 183; key to Lachnosterna in, 304; beneficial insects in, 330, 737.

Kenya Colony, locusts in, 679; Pseudococcus kenyae in, 592; parasites of Trypetids in, 492.

kenyae, Pseudococcus.

Kerosene, for extracting insecticidal principles of plants, 35, 36, 725, 726; atomised sprays of

pyrethrum in, against leafhoppers 134; fumigants combined with, against borers and pests in soil, 262, 295, 431, 449, 748; orchard crates dipped in, against *Cydia* pomonella, 125; and creosote for treating timber against Lyctus, 514; and fuel oil, warehouses sprayed with, against rice pests, 218; rice fields treated with, against Parnara guttata, 498.

Kerosene Emulsion (see Oil Emulsions).

kewensis, Aleuroplatus.

Kieselguhr, as a carrier for anabasine dust, 236; chloropicrin adsorbed on, for soil fumigation,

Kiln-sterilisation, of timber against Lyctus, **190, 707.**

King Island, Halotydeus destructor in, 582.

kirkpatricki, Microbracon.

Kola (see Cola nitida). kolae, Balanogastris.

Kolofog, 25. (See Bentonite-sul-

Kophene cuprea, bionomics of, on banana in Madras, 499.

Korea, cotton pests in, 309, 674; orchard pests in, 243, 605, 671; miscellaneous pests in, 264, 438, 767, 768; list of tree pests in, 208; studies on Lamellicorns in, 436; Scolytoidea of, 207; beneficial insects in, 108, 264, 309, 310, 768; Melaphis chinensis producing tannin galls in, 264; introduction of Tiphia popilliavora into U.S.A. from, 324.

koreana, Miridiba.

Korror (see Japanese Mandated Islands).

Kouzka Beetle (see Anisoplia austriaca).

krameri, Eberhardia (Cosmoglyphus, Tyroglyphus).

kraunhiae, Pseudococcus. Kryocide, **736.** (See Cryolite.)

kuehniella, Ephestia. Kurrajong (see Brachychiton).

taxicornis, habits Labidostomis and control of, on vines in Sicily, 70.

laburni, Aphis. lacca, Laccifer.

Laccifer lacca (Lac), pests of host-tree of, in Ceylon, 359; book on cultivation of, in India, 311.

Lachniella, producing honey-dew in Central Europe, 459.

Lachnopus aurifer, on Citrus in

Jamaica, 490.

Lachnosterna, in U.S.A., 324, 545, 696, 729; species and importance of, in Iowa, 23, 225; key to larvae of, in Kentucky, 304; measures against, 324, 696, 729.

Lachnosterna diomphalia, bionomics and control of, in Korea, 436,

Lachnosterna hirticula, in Iowa, 225. Lachnosterna hirtiventris, in Iowa,

Lachnosterna implicita, in Iowa, 23, 225: tests of dusts against,

Lachnosterna jamaicensis, on eggplant in Jamaica, 490.

Lachnosterna rugosa, in Iowa, 225. Lachnosterna smithi, biological control of, on sugar-cane in Mauritius and Barbados, 176, 222,

316, 317. titanis, measures Lachnosterna against, in Korea, 436.

Shinji Lachnus fasciatus, (see Cinara shinjii).

Lachnus laricis (see Cinara).

Lachnus pichtae (see Cinara).

Lachnus salignus, on pear Britain, 613.

Lacon murinus, soil conditions in relation to, in Russia, 429.

Lacon variabilis, on sugar-cane in Queensland, 151.

lacordairei, Rhadinosomus.

Lactuca scariola, destruction against Hercothrips fasciatus in California, 19, 129.

Laemophloeus, tests of fumigants on, in U.S.A., 389, 741.

Laemophloeus minutus, 741.

Laemostenus complanatus, predacious on Cydia pomonella Victoria, 169.

laetatorius, Bassus (Diplazon).

laetus, Xylotoles.

laevigata, Disonycha.

laevigatus, Alphitobius.

laevimargo, Diorymerellus.

laevis, Longitarsus (see L. succineus).

Lagerstroemia speciosa, Coloborhombus fasciatipennis on, in Java,

lahorensis, Rhopalosiphum (see R. rufomaculatum).

Lamachus albopictus, sp. n., parasite of Diprion nipponicum in Japan, 296.

Calotermes; lamanianus, termes (Schedorhinotermes).

laminatus, Praxibulus.

Lampides boetica (see Cosmolyce).

Lampronia rubiella (see Incurvaria). Lambrosema indicata, new Braconid parasite of, in Cuba, 330; on Derris elliptica in Porto Rico, 224. lamprosemae, Stantonia.

Lampyris noctiluca, establishment of, against Helix aspersa Australia, 450.

Lamtoro (see Leucaena glauca). lanigera, Oregma (Ceratovacuna). lanigerum, Eriosoma.

Lansium domesticum, new Coccid on, in Java, 607.

Lantana camara, Teleonemia lantanae introduced into Australia against, 86.

Lantana crocea, Teleonemia lantanae destroying, in Fiji, 86.

lantanae, Teleonemia.

lapathi, Cryptorrhynchus. Laphygma exempta, recorded as Spodoptera mauritia in Hawaii, 692; measures against, in Natal, 411; in Nyasaland, 114; in Queensland, 151; in S. Rhodesia, 90; in Sierra Leone, 240; on cereals and grasses, 114, 240; on sugar-cane, 151, 411; natural enemies of, 90, 114, 151.

Laphygma exigua, measures against, in S. Africa, 152; in Arizona, 580; in Hawaii, 214; natural enemies of, in India, 55; in Libya, 261; in Sudan, 513; on cotton, 513, 580; on lucerne, 152, 261; on other plants, 55, 214.

Laphygma frugiperda, on cotton in Argentina, 362; on rice in Br. Guiana, 369.

lapidarius, Biscirus. laracana, Epinotia.

Larch (Larix), Coleophora laricella on, in Austria, 659; Barypithes pellucidus on, in Britain, 625; Aphids on, in Czechoslovakia, 657, 715; Otiorrhynchus hirticornis on, in Germany, 344, 345; Lymantria dispar on, in Japan, 604; Anomala pallidipennis on, in Korea, 767; fauna of litter of, in Rumania, 193; Chermes on, in Switzerland, 459; pests of, in U.S.A., 441, 543, 745; not a favourable food-plant for Dendrolimus spectabilis, 672; effect of

soil treatment with lead arsenate on, 441.

Larch Case-Bearer (see Coleophora laricella).

lardarius, Dermestes. laricella, Coleophora.

laricellae, Gelis (Pezomachus).

laricinellae, Chrysocharis (Entedon). laricis, Cinara (Lachnus).

Lariophagus distinguendus, parasite of Bruchus brachialis in N. Carolina, 24.

Larix (see Larch).

Larra americana, introduced into Porto Rico from Brazil against Scapteriscus vicinus, 223.

Larvaevora (see Tachina) larvarum, Tachina (Eutachina).

Lasioderma serricorne, damaging books in Netherlands Indies, 706; in stored tobacco in U.S.A., 363; 396; studies on oviposition of, 164, 165; experiments with low temperatures against, 363; effects of fumigants on, 396, 567, 741.

Lasius, ecology and relation to Aphids of species of, in Germany,

Lasius brunneus, 195.

Lasius fuliginosus, 195. Lasius niger, 195, 196.

Lasius niger alienus, associated with Aphids on rubber-producing plants in Ukraine, 106, 107.

Laspeyresia (see Cydia).

Laspeyresia caryana and prunivora (see Enarmonia).

Latania loddigesii, Homaledra sabalella on, in Florida, 22.

lataniae, Aspidiotus.

Oncometopia; lateralis, Astycus;

Phytomyza (Napomyza).

Lathyrus odoratus (see Sweet Pea). sativus, Pachytychius Lathyrus strumarius on, in Morocco, 641. laticollis, Gastrallus; Prionus.

laticostalis, Margaronia.

latiferreana, Cydia (Mellissopus).

latiusculus, Listronotus.

latreillei, Tropidacris.

Corymbites (Selatosomus); latus, Tarsonemus.

Latvia, Coleoptera on alsike clover and crucifers in, 256, 257; forest pests in, 90, 721.

laurenti, Heteroderes.

Laurus camphora (see Cinnamomum).

Laurus nobilis, Aspidiotus perniciosus not developing on, 305.

Lau yl Rhodanate (Thiocyanate), 272, 274. (See Thiocyanates.)

Lavender (Lavandula), Sophronia humerella on, in Germany, 301; insecticidal properties of oil of, 249, 250.

Lawns (see Turf).

Lead, insects boring in, 232, 262; problems of residues of, on sprayed fruit, 79, 140, 175, 320, 368, 402, 490, 555, 556, 646, 750.

Lead Arsenate, against cereal pests, 207, 265, 309, 441, 768; and molasses, maize swabbed with, against Heliothis, 494; against pests of forest and shade trees, 226, 270, 332, 335, 393, 441, 546, 547, 548, 550, 582; against pests of iris, 274, 598; against pests of malvaceous plants, 358, pests of malvaceous plants, 358, 766; against olive pests, 350; against orchard pests, 5, 9, 15, 16, 19, 25, 44, 46, 56, 62, 79, 99, 100, 108, 111, 125, 140, 144, 149, 150, 168, 169, 174, 175, 243, 250, 254, 255, 259, 260, 287, 291, 299, 318, 320, 332, 333, 343, 401, 442, 443, 444, 449, 479, 488, 489, 490, 500, 510, 525, 544, 554, 555, 556 500, 510, 525, 544, 554, 555, 556, 557, 558, 587, 588, 594, 614, 615, 647, 652, 653, 658, 713, 718, 723, 724, 725, 732, 734, 748, 758; effect of, on parasites of orchard pests, 555, 558; against pests of nut trees, 138, 324, 500, 547; against pests of small fruits, 45, 61, 270, 282, 443, 563, 646, 735; against Tineid on palms, 22; against pests of solanaceous plants, 210, 581; against sweet potato pests, 207, 222, 447; against vine pests, 70, 111, 201, 335; against pests of other crops, 151, 207, 266, 713; against pests in soil, 210, 324, 393, 403, 441, 447, 727, 729; effect of phosphates on toxicity of, to larvae of Popillia, 324; tests of toxicity of, to insects, 30, 205, 226, 388, 572, 741, 742; in baits, 182, 270, 393, 441 540, 705; in bait-sprays, 61; dusting with, 210, 226, 443, 705, 766; carriers for, 125, 226, 443, 705, 734, 766; oil-impregnated dusts of, 125, 653, 734; aeroplane dusting with, 748; formulae for, in sprays, 5, 9, 15, 16, 22, 44, 45, 46, 56, 62, 111, 125, 138, 144, 149, 151, 168, 169, 174, 175, 210, 222, 259, 282, 287, 324, 332, 333, 335, 350, 358, 393, 401, 443, 447, 479, 489, 490, 510, 546, 547, 548, 550, 554, 555, 563, 582, 588, 594, 598, 614, 615, 646, 647,

653, 723, 724, 733, 742, 743, 758; adhesives and spreaders for, 16, 21, 151, 168, 174, 175, 222, 310, 320, 350, 393, 442, 443, 447, 488, 489, 510, 546, 547, 548, 550, 594, 614, 652, 653, 713, 723, 725, 750; 324, 444, 448, 489, 510, 588, 653, 723, 724, 725, 733, 743; and lime-sulphur, 25, 100, 250, 350, 489, 510, 614, 748; chemistry of mixtures of lime-sulphur and, 742; and nicotine, 168, 210, 250, 260, 442, 554, 555; and oils, 169, 174, 175, 254, 259, 270, 318, 402, 442, 488, 510, 554, 555, 588, 614, 652, 653, 708, 750; inverted spray mixtures of oils and, 174, 175, 488, 544, 653; addition of sodium silicate to sprays of, 402; and sulphur, 15, 443, 489, 724, 734; and zinc sulphate, 16, 125, 138, 174, 443, 448, 588, 724; and injury to plants, 16, 21, 79, 125, 138, 150, 210, 324, 393, 441, 442, 489, 500, 510, 544, 552, 588, 614, 647, 653, 705, 743; triplumbic form of, 500, 748; problems of residues of, on fruit, etc., 25, 140, 169, 175, 259, 318, 320, 401, 402, 442, 488, 489, 490, 510, 525, 555, 556, 614, 646, 647, 652, 653, 735, 750; estimation of deposits of, by weight and surface area of apples, 320: other insecticides compared with, 16, 25, 30, 62, 79, 108, 138, 149, 150, 182, 291, 310, 401, 402, 442, 443, 479, 544, 547, 548, 552, 555, 556, 647, 652, 653, 658, 729, 732, 733, 735; increased use of, in U.S.A., 651.

Leaf-cutting Ants (see Atta and Acromyrmex).

Leaf-cutting Bee (see *Melipona* testacea).

Leaf-mines (caused by insects), keys to, in central and northern Europe, 247; study of, in Rumania, 192.

Leaf-roll (see Virus Diseases).

Leafhopper, Apple (see *Typhlocyba froggatti*).

Leafhopper, Beet (see *Eutettix* tenellus).

Leafhopper, Blunt-nosed (see Euscelis striatulus).

Leafhopper, Grape (see Erythroneura comes).

Leafhopper, Peach or Plum (see Macropsis trimaculata).

Leafhopper, Potato (see Empoasca fabae).

Leafhopper, White Apple (see Typhlocyba pomaria).

Leather, Gibbium psylloides attacking, in Germany, 519.

Lebeckia retmoides, not containing rotenone, 182.

Lebeda nobilis, bionomics and control of, on Myrica rubra in China, 244.

Lebia grandis (predacious on Leptinotarsa decemlineata), imported into France from N. America, 595; rearing of, 595.

Lecanium, on almond in California, 18; on plum in Czechoslovakia, 659; on fruit trees in Seychelles, 433; natural enemies of, 18, 433, 659.

Lecanium armeniacum, aeroplane spraying against, on apricot in California, 19, 748.

Lecanium corni, factors affecting susceptibility of trees to, in Germany, 251.

Lecanium fletcheri, on arbor-vitae in Indiana, 126.

Lecanium hemisphaerica (see Saissetia).

Lecanium hesperidum (see Coccus). Lecanium nigrofasciatum, measures against, on peach in Pennsylvania, 26.

Lecanium perforatum (see Eucalymnatus).

Lecanium rufulum (pulchrum), on oaks in Czechoslovakia, 657.

Lecanium tessellatum (see Eucalymnatus).

Lecanium viride (see Coccus).

Lecanobius, imported into Peru against Saissetia oleae, 81.

lecheana, Tortrix (Cacoecia).

lechriaspis, Eucosma (Spilonota). lectularius, Cimex.

Ledra auditura, on Camellia in Japan, 438.

leefmansi, Doirania.

Leefmansia bicolor, parasite of Sexava in New Guinea Territory, 436.

lefroyi, Microbracon.

Legislation, review of, dealing with fumigants in Germany, 347. (See Plant Pest Legislation.)

leguminis, Eupteromalus.

leguminosae, Aphis (see A. laburni).

Leis conformis (predacious on Aphids), in Australia, 335, 450; introduced into Egypt, 450.

Leis dimidiata ab. quinquedecimspilota, bionomics of, in Florida, 727.

Lema fulvicollis [? fulvicornis], on turmeric in Ceylon, 359.

Lema melanopa, on cereals Czechoslovakia, 657.

Lema oryzae, measures against, on

rice in Japan, 309.

Lema trilineata, effect of rearing Pentatomids on, in Ohio, 13, 14. Lema tristis, bionomics of, in Man-

churia, 767.

Lemon, mites on, in California, 173, 650; pests of, in New Zealand, 266; little attacked by Ceratitis capitata in Palestine, 406; Coccids on, in Porto Rico, 41; Aspidiotus perniciosus on, Russian Union, 305; Aphids on, in Tanganyika, 36.

Lemon (Fruit), in baits for grass-

hoppers, 139.

Lentil, Pachytychius strumarius on, in Morocco, 641; Etiella zinckenella on, in Russia, 432.

Lentils (Stored), fumigation against Bruchus lentis, 414; practically immune from grain mites, 96.

lentis, Bruchus.

lepelleyi, Pseudococcus.

leperii, Epidiaspis (Diaspis).

Leperisinus (see Hylesinus).

lebida, Parasa.

Lepidaphycus bosqui, parasite of Eriopis connexa in Argentina, 289.

Lepidiota caudata, damaging pasture in Queensland, 267, 494; epizootic in, 267.

Lepidiota pinguis (see Leucopholis). Lepidium nitidum, relation of Eutet-

tix tenellus to, in California, 323. Lepidoderma albohirtum, natural enemies and control of, on sugarcane in Queensland, 56, 151.

Lepidoglyphus destructor, mics of, in stored cereals in Russian Union, **96, 97.**

Lepidoglyphus destructor mixtus, in stored cereals, etc., in Russian Union, 95.

Lepidoglyphus michaeli, in hay and grain in Russia, 95.

Lepidoptera, biological studies of, by means of light traps in Maine, 746: list of, producing silk, 68. Lepidosaphes, on Citrus in Trinidad,

Lepidosaphes beckii (pinnaeformis, auct.) (Citrus Mussel Scale), in Portuguese E. Africa, 377; in S. Africa, 377; in Brazil, 82; in Cyprus, 107, 706; in Indo-China, 185; in Palestine, 77, 636-638; in Peru, 81; in Porto Rico, 41; in U.S.A., 41, 690, 727; bionomics of, 41, 377, 378, 636-638; measures against, 82, 107, 185, 637, 690, 707, 727; characters of, 378.

Lepidosaphes citricola (see L. beckii). Lepidosaphes dispar (on cassava), measures against, in Madagascar, 454; predacious enemy of, in

Tanganyika, 112.

Lepidosaphes gloveri, new parasite of, on Euonymus in Louisiana, 41; on Citrus in Transvaal, 378; surface tension of oil emulsions affecting their toxicity to, 702; characters of, 378.

Lepidosaphes halli, eradication of, in California, 18, 747.

Lepidosaphes pinnaeformis, (see L. beckii).

Lepidosaphes sacchari, restricted to Miscanthus in Formosa, 263.

Lepidosaphes ulmi, in Chile, 221; on olive in Spain, 700; in Transcaucasia, 481; in U.S.A., 25, 44, 172, 547, 738; measures against, on fruit trees and cranberry, 25, 44, 172, 221, 481, 738; on beech, 547.

Lepidosaphes vermiculus, sp. n., parasite of, on coconut in Mauritius, 66.

lepidotus, Moropactus.

lepidus, Platypus.

saccharina, measures Lepisma against, in houses in Germany and U.S.A., 37, 224, 519; damaging papers, 519.

Lepra Explosiva, 581.

Leptacis tipulae (parasite of wheat gall-midges), in Germany, 620; in Sweden, 155, 231; bionomics of, 231.

brevipennis, damaging Leptidea wicker in Czechoslovakia, 658.

Leptinotarsa decemlineata (Potato Beetle), danger of introduction of, into Austria, 518; in Belgium, 120, 368, 719; in France, 5, 368, 412, 415, 594, 595, 597; in Germany, 120, 252, 347, 368, 418. 719: legislation against introduction of, into Germany,

91: in Holland, 120: in Luxemburg, 120, 368; in Saskatchewan, 170; in Switzerland, 70, 91, 120; in U.S.A., 14, 15, 215, 564, 643; conference on control of, in Europe, 467; breeding on wild species of Solanum, 13, 14, 415; taken on high buildings, 215; experimental rearing and utilisation of natural enemies of, 13, 14, 594, 595; tests of chemical manures against, 412; measures against, 252, 418, 467, 518, 564, 597, 598, 643: susceptibility of, to fluorine, 216; anatomy and physiology of, 368; Lilioceris lilii mistaken for, 520.

Leptobyrsa (see Stephanitis).

Leptocneria reducta, bionomics and control of, in New South Wales, 582.

Leptocoris augur, on Schleichera trijuga in Ceylon, 359.

Leptocorisa acuta, on rice in Madras, 360.

Leptoglossus membranaceus, on cucurbits and Citrus in Ceylon, 358, 359.

Leptoglossus oppositus, causing black pit and kernal spot of pecan in Florida, 587.

Leptoglossus phyllopus, causing black pit and kernal spot of pecan in Florida, 587.

Leptomastidea abnormis (see Tanaomastix).

Leptomastidea antillicola, sp. n., parasite of Ferrisiana virgata in Porto Rico, 41.

Leptostylus praemorsus, on Citrus in Dominica, 751.

Lepturoides, Omophlus.

Leskiopalpus diadema (see Stomatodexia).

Lespedeza, Anomala pallidipennis on, in Korea, 767.

Lespedeza capitata, Calomycterus setarius on, in Connecticut, 15.

Lethalate, as a wetter for sprays, 614, 615; composition of, 615.

Lethane (see Thiocyanates).

Lethane Spreader, 727.

Lethrus apterus, measures against, in nurseries in Russia, 352, 432; bionomics of, 352.

Lettuce, pests of, in U.S.A., 19, 488; as trap-crop for *Melolontha* 521; experimental feeding of insects on, 11, 147, 194; methyl bromide not impairing germination of seed of, 389.

Lettuce, Prickly (see Lactuca scariola).

Leucaena glauca, coffee mealybugs on, in Java and New Guinea Territory, 182, 495, 606.

Leucania (see Cirphis). leucarthros, Necremnus.

Leucodiaspis (Leucaspis), species of, in New Zealand, 303.

leucoloma, Naupactus. leucomelas, Acyphas. leuconotus, Anthores.

Leucopholis pinguis, on rubber in Ceylon, 358.

leucophthalmus, Philaenus.

Leucopis obscura (predacious on Chermes spp.), liberation of, in Australia, 449; imported into U.S.A., from Canada, 329.

Leucoptera coffeella, parasites a doubtful factor in control of, on coffee in W. Indies, 223.

leucopterus, Blissus.

Leucosomus pedestris (see Cleonus). leucosticha, Cirphis.

leucotreta, Argyroploce.

Levuana iridescens, on coconut in Fiji, 211.

librinocens, Neogastrallus.

Libya, Ceratitis capitata in, 5, 260; new Coccids in, 496; miscellaneous pests in, 260; Sphingid on vines in, 201; beneficial insects in, 260, 261, 410.

libyca, Empoasca.

Licania rigida, new weevil on, in Brazil, 106, 115.

licaniae, Conotrachelus.

licas, Heteronychus.

licoides (licus, auct.), Castnia.

Light, insecticides affecting action of, on plants, **525**.

Light (Artificial), insects attracted by, 454, 556, 572, 623, 723, 729, 732, 747; comparison of types of, 573, 723, 732, 747.

Light-traps, for cockchafers, 251, 545; of doubtful value for Gryllotalpa spp., 457; Ichneumonid caught in, 54; for leaf-hoppers, 54, 439; for Lepidoptera, 54, 55, 244, 245, 488, 669, 732; biological studies of Lepidoptera by means of, 535, 746; electrocuting types of, 251, 488.

ligneus, Otiorrhynchus. Lignin Pitch, **506.** (See Sulphite

Lye.)
ligustici, Otiorrhynchus (Brachyr-

ligustri, Phyllocoptes. Ligustrum (see Privet).

rhinus).

Ligyrus bituberculatus, on sugarcane and banana in Brazil, 502. Ligyrus ebenus, on sugar-cane in Br. Guiana, 369.

Ligyrus humilis (see Euetheola). Ligyrus tumulosus, on egg-plant in Jamaica, 490.

lilacinus, Pseudococcus.

lilii, Lilioceris.

Lilioceris lilii, on potato in Germany, 520; mistaken for Leptinotarsa decemlineata, 520.

Lilioceris merdigera, on onions in

Finland, 201.

Lily, Aphis gossypii transmitting rosette disease of, in Bermuda, 626.

Lima Bean Pod Borer (see Etiella zinckenella).

Lima Beans, Bruchids on, in Burma, Maruca testulalis on, in Porto Rico, 643; pests of, in U.S.A., 129, 270, 327, 328, 509, 566; problem of fluorine residues on, 327.

Lima Beans (Stored), pests in, 7,

748.

limacina, Caliroa.

limbatus, Adrastus.

Lime (Citrus), pests of, in W. Indies, 402, 448, 752.

Lime (Tilia), pests of, in Finland, 201: Lymantria dispar on, in Jugoslavia, 759; Heterogena asella on, in Russia, 433; experimentally attacked by Otiorrhynchus hirticornis, 345; susceptibility of timber of, to Codiosoma

spadix, 354. Lime, as a carrier for dusts, 142, 222, 236, 276, 333, 334, 397, 399, 443, 498, 564, 565, 566, 643, 644, 645, 646, 705; oil-impregnated dusts containing, 544, 631, 734; and adhesives, spraying with, against Popillia, 443, 551, 552; addition of, to arsenical sprays, 15, 16, 125, 291, 324, 401, 431, 433, 444, 448, 479, 484, 488, 510, 559, 566, 588, 653, 723, 724, 725, 733, 743; and potassium permanganate, 72; addition of, to other sprays, 186, 554, 556, 559, 701, 724; as a repellent, 251, 346, 443, 551; as a soil dressing against pests, 190, 718; for destroying ants, 589; with stored seeds against pests, 125, 642; use of, in connection with spray residues on apples, 318; formulae containing, 72, 125, 142, 186, 236, 276, 291, 324, 333, 334, 397, 399, 431, 443, 444, 448, 479, 488, 510, 551, 552, 554, 559, 564, 566, 588, 643, 644, 701, 723, 724, 733, 734, 743.

Lime-sulphur, against Aphis gossypii, 309; against Coccids, 44, 55, 172, 220, 221, 272, 350, 351, 455, 562, 663, 724, 728; value of, against certain Lepidoptera, 16, 29, 718; against mites, 16, 72, 171, 246, 297, 334, 471, 488, 521, 583, 674, 690, 693, 694, 724; against Paratrioza cockerelli, 579, 643; against thrips, 291, 727; empty mushroom houses sprayed with, against pests, 579; effects of, on natural enemies of pests, 16, 557, 558; as a repellent 350, 602; formulae for, 16, 29, 44, 72, 131, 149, 171, 172, 221, 234, 246, 272, 334, 350, 351, 488, 489, 510, 521, 579, 602, 614, 643, 693, 718, 724, 742; formulae containing, 29, 131, 149, 234, 272, 350, 489, 510, 614, 643, 742; and arsenicals, 25, 100, 250, 350, 489, 510, 614, 643; chemistry of mixtures of lead arsenate and, 742; decreasing effectiveness of derris, 690; and nicotine, 29, 131, 149, 150, 250, 291, 727; and oil emulsions, 272; phytocidal properties of oil emulsions in combination with, 708; and iron sulphate, pyrethrum unsatisfactory with, 534; and salt (see California Mixture); and sulphur, 727; and tar distillates, 234, 235; precipitation of, by addition of zinc sulphate, 488: reducing photosynthesis on apple leaves, 216.

Limonius aeruginosus, soil conditions in relation to, in Russia,

Limonius californicus (see Pheletes). Limonius canus (see Pheletes).

Lindorus lophanthae (predacious on Coccids), in N. Africa, 410; in Réunion, 501; history of distribution of, 410.

Atomaria; Brontispa; linearis, Spintherus.

lineata, Celerio (Deilephila); Navomorpha; Sitona.

lineatella, Anarsia.

lineaticollis, Antestia.

lineatus, Agriotes; Philaenus; Xyloterus.

Lineodes integra, bionomics of, in California, 744. lini, Dasyneura.

Linolic Acid, in baits for Cydia, 135.

Linseed (see Flax).

Linseed Oil (see Oil, Linseed).

Linseed-oil Soap (see Soap, Linseed-

lintneri, Tyroglyphus.

Linum perenne, Ceuthorrhynchus sareptanus on, in Siberia and Russian Far East, 351, 601.

Linum usitatissimum (see Flax).

Liothrips (Phloeothrips) oleae (on olive), in Italy, 350; in Libya, 261; in Spain, 700; measures against, 350, 700.

Liothrips urichi (destroying Clidemia hirta), utilisation of, in Fiji, 59; introduced into Solo-

mon Islands, 703.

Lipari Islands, Coccids of, 119.

Lippia citriodora, tests of oil of, against Aphids and mites, 249, 250.

Liriomyza (see Agromyza).

Listroderes obliquus, in U.S.A., 144, 691, 747; bionomics of, 691; fumigation of celery against, 747; on tobacco, 144.

Listronotus latiusculus, bionomics and control of, in U.S.A., 565,

566.

Litchee (Nephelium litchi), pests of, in China and Formosa, 674, 693. lithocolletidis, Pleurotropis.

Lithocolletis populifoliella, on poplar in Rumania, 193.

litura, Prodenia.

livida, Phaenocarpa.

livornica, Celerio (Deilephila) lineata.

lixi. Habrocvtus.

Lixus brevirostris, parasites of, on beet in Morocco, 417, 526.

Lixus junci (on beet), in Italy, 528: in Morocco, 416, 417, 526; bionomics of, 416; parasites of, 417, 526, 528.

Lixus scabricollis, parasites of, on beet in Morocco and Italy, 417, 526, 528.

Lizards, destroying noxious insects, 128, 269, 397.

lobipennis, Brachyexarna.

Lochmaea suturalis (Heather Beetle), bionomics and control of, Britain, 418.

Lockemann Method, for determination of arsenic content of sprays, 772.

Locris neumanni, on rice in Tanganyika, 113.

Locust, Asiatic (see Locusta migratoria).

Locust, Black-tipped (see Chortoicetes terminifera).

Locust, Desert (see Schistocerca gregaria).

Locust, Italian (see Calliptamus italicus).

Locust, Post-oak (see Dendrotettix quercus).

Locust, Red (see Nomadacris septemfasciata).

Locust Tree, Black (see Robinia

pseudacacia).

Locusta migratoria, in Central Asia, 77, 426; distribution of, in Australia, 584; in India, 628; in Sierra Leone, 240; Aspergillus flavus infesting, 591; bio-610, 628, nomics of, 629: measures against, 77, 426, 611.

Locusta migratoria capito, in Madagascar, 67, 416; damaging sugarcane, 67; effects of temperature

and humidity on, 416.

Locusta migratoria manilensis, in N. Borneo, 112; in Netherlands Indies, 685; birds destroying, in Philippines, 675.

Locusta migratoria migratorioides, in E. Africa, 679, 684, 687; in W. Africa, 112, 679, 684, 687; possibly in S. Africa, 684; outbreak centres and migrations of, 111, 679, 684, 687; effect of temperature on activity of, 147. 148: losses and expenditure due to, in Mozambique, 676.

Locustana pardalina, in Angola, 687; in S. and S.W. Africa, 676, 682, 684; outbreak centres of, 684: losses and expenditure due to, 676, 687; measures against,

682, 684.

Locusts and Grasshoppers, in E. Africa, 104, 111, 112, 113, 114, 115, 372, 642, 676, 678, 679, 682, 683, 684, 687; in S. Africa, 152, 411, 676, 681, 682, 683, 684, 687; in S.W. Africa, 676; in W. Africa, 112, 240, 315, 679, 684, 687, 755; in Algeria, 180, 239, 685; in Argentina, 146, 290, 655, 689 in S.W. Asia, 111, 112, 198, 199, 486, 683, 685, 687; in Australia, 111, 169, 440, 449, 450, 494, 583, 584, 629, 676, 686, 703; in N. Borneo, 112; in Brazil, 290; in Canada, 170, 440, 486, 540, 590, 676, 680, 686; in Ceylon, 358; in China, 498; in Cyprus, 107; in Egypt, 112, 682, 685,

687; in Formosa, 769; in India, 55, 199, 628, 629, 666, 677, 678, 683, 686; in Netherlands Indies, 603, 685; in Italy, 162, 163, 592; in Japan, 672; in Jugoslavia, 685; in Libya, 260; in Madagascar, 67, 416, 454; in Morocco, 180; in Philippines, 675; in N. and S. Rhodesia, 114, 410, 683, 687; in Rumania, 610; in Russian Union, 77, 246, 353, 357, 426; in Spain, 698; in Sudan, 6, 111, 112, 198, 679; in U.S.A., 18, 19, 23, 43, 44, 48, 123, 124, 128, 133, 138, 139, 246, 282, 326, 393, 398, 574, 580, 588, 608, 609, 676, 680, 681, 686, 748; in Uruguay, 290; international conference on, 676-688; colour variations and phases of, 6, 114, 146, 239, 410, 449, 590, 592, 593, 628, 677–687, 689; outbreak centres and ecology of, 6, 72, 147, 162, 169, 198, 199, 426, 449, 583, 584, 610, 611, 678-687, 698, 703, 755, 756; migrations of, 111, 112, 114, 146, 199, 583, 584, 628, 678, 679, 681, 683, 687; effects of temperature and humidity on, 104, 139, 147, 169, 410, 416, 583, 584, 628, 679; water-content of adults of, 184; correlation of outbreaks of, with sun spots, 486; potential reproductive capacity of, 163, 677; fungus and bacterial diseases of, 146, 147, 184, 591, 681, 682, 688; mites infesting, 147, 163, 169, 629; nematode parasites of, 55, 123, 124; insect enemies of, 44, 114, 139, 146, 147, 163, 169, 290, 450, 484, 588, 593, 629, 682, 688, 689; bats destroying, 629; birds destroying, 128, 169, 282, 498, 629, 655, 675, 682; doubtful danger to storks of feeding on poisoned, 180; estimation of abundance of eggs of, **357**; measures against (see also under Baits), **77**, **133**, **152**, **169**, **180**, **246**, **353**, **426**, **427**, 498, 611, 680, 681, 682, 684, 686, 687, 698, 769; aeroplanes for distributing poison against, 426; estimates of losses due to, 44, 170, 676, 677, 685, 686, 687, 688; anatomy and histology of, 440, 624; egg-pods of, 246; effects of hypertonic solutions and cyanide on eggs of, 304; bibliography of papers on, 112. Lodgepole Pine (see Pinus con-

torta).

Loganberry, pests of, in Britain,

Lolium spp. (Rye Grass), Dipterous pests of, in Britain, 382.

Lombardy Poplar (see Populus nigra italica)

Lonchaea aristella, on figs in Palestine, 406.

Lonchaea flavidipennis, on asparagus in Britain, 33.

Lonchaea pendula, on Capsicum annuum in Chile, 654.

Lonchocarpus, classification, constituents and insecticidal properties of species of, 224, 318, 319; dusting tests with, 347, 348. (See Barbasco, Cubé, Haiaris and Timbo.)

londinensis, Cylindroiulus.

longicaudatus, Biosteres (Opius). longicellana, Tortrix (Cacoecia).

longiceps, Orgilus.

longicollis, Odoiporus. longicornis, Diabrotica.

longidentifera, Euxoa (Agrotis).

longimacula, Anastrepha.

longior, Tyroglyphus (see phagus putrescentiae).

longipenis, Tetranychus (Apotetranychus).

longipennis, Dissosteira; cephala (see Brontispa longissima); Scirtothrips.

longipes, Anoplolepis (Plagiolepis). longivostris, Ischnaspis.

longiscapus, Aenasius.

longisetosus, Hemiteles inimicus.

longispinus, Pseudococcus (see P. adonidum).

longissima, Brontispa (Oxycephala). Longistigma caryae, technique for rearing Coccinellids on frozen, in Ohio, 649.

longistylus, Dacus.

Longitarsus laevis (see L. succineus). Longitarsus lycopi, on mint in Poland, 255.

Longitarsus parvulus, on flax in Poland, 255.

Longitarsus succineus, on chryssanthemum in Britain, 297.

Longitarsus waterhousei, on mint in Poland, 255.

longiventris, Eurydinoteloides.

longula, Sitona.

longulus, Eulophus.

of Rhagoletis Lonicera, relation cerasi to species of, in Switzerland, 156, 300; Acidia lucida on, 156.

Islands, Phyllocnistis Loochoo citrella on Citrus in, 438; Cylas formicarius in, 669; pests of sugar-cane and cereals in, 264, 437, 516; beneficial insects in, 142, 437; toad introduced into, against noxious insects, 605.

lophanthae, Lindorus (Rhizobius).
Lophocateres pusillus, in butternuts imported into Holland from Dutch Guiana, 529; in stored cassava tubers in Madacasgar, 455.

Lophosternus huegeli (see Dorysthenes).

Lophyroplectus nipponensis, sp. n., parasite of Diprion sertifer in Japan, 296.

Lophyrus (see Diprion).

Loquat (see Eriobotrya japonica).

loreyi, Cirphis.

lorisona, Virachola (Deudorix). Lorol, Sulphonated, as a spreader for derris sprays, 191, 382.

loti, Contarinia.

Lotis neglecta, predacious on Aonidiella aurantii in S. Africa, 456.

Louisiana, pests of chrysanthemum in, 395, 396; cotton pests in, 276, 279, 328; miscellaneous pests in, 286, 580; pecan pests in, 323, 324; pests of field and stored rice in, 218; sugar-cane pests in, 28, 137, 390, 391; Aphids transmitting sugar-cane mosaic in, 577; beneficial insects and biological control in, 41, 137, 227, 390, 391; Epicaerus cognatus intercepted in, on potatoes from Mexico, 11.

lounsburyi, Aphycus.

Loxostege bifidalis, bionomics of, on cotton etc. in Brazil, 333, 334.

Loxostege sticticalis, in Russian Union, 94, 157, 257, 303, 308, 354, 424, 472, 474; in Saskatchewan, 170; in U.S.A., 44, 126; on beet, 44, 94, 157, 308, 472; on cotton, 354; on other plants, 126, 170, 308; natural enemies of, 94, 303, 354; bionomics and physiology of, 44, 157, 257, 308, 424, 440, 472, 474.

lucasseni, Podothrips (Kentrono-

thrips).

Lucerne, pests of, in S. Africa, 151;
Astylus atromaculatus on, in
Argentina, 705; pests of, in
Australia, 84, 132, 494; Sitona
lineata on, in Czechoslovakia,
657; pests of, in Germany, 461,
520; Hypera variabilis on, in
Holland, 299; pests of, in
Hungary, 527; Contarinia loti

on, in Italy, 189, 190; Phytodecta fornicata on, in Jugoslavia, 760, 761; pests of, in Libya, 260, 261; *Prodenia litura* on, in Palestine, 638; pests of, in Russian Union, 76, 238, 356; Contarinia medicaginis on, in Sweden, 232; pests of, in U.S.A., 18, 19, 118, 126, 127, 135, 136, 139, 140, 216, 326, 489, 545, 580, 747; grasshoppers damaging, 19, 139, 140, 152; review of relation of insects to diseases of, 386: effects of Empoasca fabae on carotene content of, 216; Blapstinus substriatus experimentally fed on, 27; planted among cereals against Blissus leucopterus, 78.

Lucerne Gall-midge (see Conta-

vinia medicaginis)

Lucerne Seed Chalcid (see Brucho-phagus gibbus).

lucida, Acidia (Myiolia). lucifugus, Reticulitermes.

Lucilia coeruleiviridis, infesting drying fruit in California, 228.

Lucilia sericata, infesting drying fruit in California, 228; susceptibility of sexes of, to nicotine, 332.

ludens, Anastrepha.

Luffa cylindrica, Dacus cucurbitae on, in Formosa, 206.

lugubrosa, Ellopia fiscellaria. luigionii, Cleonus (Conorrhynchus).

luminosus, Pyrophorus.

lunata, Cydonia; Stomatorrhina.
Lupin (Lupinus), Pomphopoea sayi
on, in Connecticut, 17; pests of
in Germany, 248, 496.

in Germany, 248, 496.
Lupin Leaf Weevils (see Sitona gressoria and S. grisea).

Lupinus perennis, Cydia glycinivorella on, in Manchuria, 310.

lupulinus, Hepialus. lurida, Scotinophara.

luridula, Anomis. lutaria, Sialis.

luteola, Galerucella. lutescens, Amblypelta.

Luxemburg, Leptinotarsa decem lineata in, 120, 368.

Lycaena argiolus (see Lycaenopsis). Lycaena boetica (see Cosmolyce).

Lycaenopsis argiolus, on raspberry in Britain, 296.

lycopersicella, Phthorimaea (Gnorimoschema).

lycopersici, Frankliniella; Phyllocoptes.

Lycophotia margaritosa, on cotton in California, 19; on potato in Hawaii, 214.

Lycophotia muscosa, on tobacco in Tanganyika, 642.

lycopi, Longitarsus.

Lycoria pusilla (see Sciara).

Lyctus (in timber), in Britain, 190, 707; in Sudan, 514; methods of depletion of starch from timber against, 190, 635, 707; other measures against, 190, 514, 707.

Lyctus africanus, in stored cassava tubers in Madagascar, 455.

Lyctus brunneus, in stored cassava tubers in Madagascar, 455.

Lyda stellata, on Pinus sylvestris in Czechoslovakia, 657.

Lydamis, on Anna in Brazil, 115. Lydella stabulans grisescens, considered correct name for Caromasia senilis (q.v.), 730.

masia senilis (q.v.), 730.

Lygaeonematus (see Pristiphora).

Lygidea mendax, measures against, on apple in Nova Scotia, 534.

Lygropia, alternative hosts of parasite of Hapalia in India, 765.

Lygus communis, measures against, on apple in Nova Scotia, **534.**

Lygus elisus, food-plants of, in U.S.A., 19, 126, 127, 128; damaging peach fruits, 127; predacious on Eutettix tenellus, 128.

Lygus hesperus, on lucerne, etc., in Utah, 126, 127, 128; damaging peach fruits, 127; predacious on Eutettix tenellus, 128.

Lygus pabulinus, on apple and bush fruits in Germany and Britain, 234, 297; oviposition

of, 234.

Lygus pratensis, on chrysanthemum in Britain, 349, 709; on potato and cotton in U.S.A., 217, 279, 565; potato wilt caused by, 217; bionomics of, 710; measures against, 279, 565, 710.

Lygus pratensis oblineatus, causing cat-facing of peaches in Delaware,

444; on clover, 444.

Lygus simonyi, measures against, on cotton in Uganda, 376.

Lygus spinolae, measures against, on vines in Europe, **521**.

Lymantria (Porthetria) dispar, in Germany, 234; relation of sunspot numbers to outbreaks of, in Japan, 604; in Jugoslavia, 759; in Russian Union, 429, 432; in U.S.A., 14, 216, 332, 560, 589, 646, 723; on cranberry, 646; on forest and shade trees, 14, 234, 332, 429, 432, 604, 759; on fruit trees, 332, 604, 759; bionomics of, 216, 759; methods for determining feeding rates of larvae of, 156; natural enemies and biological control of, 234, 589, 759, 760; polyhedral disease in, 234; measures against, 332.

Lymantria monacha (in forests), in Czechoslovakia, 657, 659, 715; in Germany, 154, 233, 346, 419, 420; in Japan, 516; bionomics and control of, on spruce, 715; natural enemies of, 233, 419, 420, 715; surviving parasitism by Meteorus versicolor, 233; use of scent of females for estimating prevalence of, 659, 665, 715; review of recent studies on, 346.

Lymantria reducta (see Leptocneria). Lyonetia clerkella, on cherry in

Ukraine, 159.

lyropictus, Periphyllus (Chaitophorus).

Lysiphlebus testaceipes, parasite of Aphids in Porto Rico, 39.

lysizone, Zizera (Zizina) otis.

Lysol, ineffective as a repellent for *Melolontha*, **346**.

Lytta vesicatoria, measures against, in forests in Russia, 430, 431, 433; effect of, on ash, 430.

M.

macaensis, Polistes. machaeralis, Hapalia. machaon, Papilio. macleayi, Camptocladius.

Macrocentrus (parasite of Homona coffearia), in Java, 337, 639; establishment of, in Ceylon, 337, 639; bionomics of, 337.

Macrocentrus abdominalis, synonym of, **342**.

Macrocentrus abdominalis var. pallidipes, parasite of Pyrausta nubilalis, in Poland, 623.

Macrocentrus amicroploides, parasite of Tortrix rosaceana in Utah, 141.

Macrocentrus ancylivorus (parasite of Cydia molesta), introduced into Australia, 448; in Canada, 510, 535; imported into Italy, 154; in U.S.A., 16, 270, 399, 544, 647, 653, 724, 735; parasite of Ancylis comptana, 399, 735; utilisation of, 16, 535, 647, 724,

735; simultaneous propagation of Ascogaster carpocapsae and, 535; Epiblema strenuana and alternative host of, 510.

Macrocentrus delicatus (parasite of Cydia molesta), imported into Italy from U.S.A., 154; alternative host of, in Ontario, 510.

Macrocentrus gifuensis (parasite of Pyrausta nubilalis), in Manchuria, 296; establishment of, in U.S.A., 730.

Macrocentrus japonicus, parasite of Pyrausta diniasalis in Manchuria, 296.

Macroglenes penetrans, bionomics of, in Sweden, 231.

macropetana, Eucosma (Spilonota).

Macropsis trimaculata, in Canada,
173; in U.S.A., 173, 385, 560;
relation of, to virus diseases of
peach, 173, 385; bionomics of,
173, 560.

Macrosiphoniella (see Macrosiphum).

Macrosiphum, on cereals in Argentina, 31, 289; on tobacco in India, 88; classification and new species of, in U.S.A., 247, 496, 696; (Macrosiphoniella), of Europe, 440.

Macrosiphum avenae, Aphid recorded as, on cereals in Argentina, 31, 289.

Macrosiphum gei, auct. (see M. solanifolii).

Macrosiphum granarium (see M. avenae).

Macrosiphum jaceae, on cornflowers in Britain, 350.

Macrosiphum onobrychis (on lucerne and peas), in S. Africa, 152; in Russia, 355; in U.S.A., 12, 126, 128, 270, 285, 291, 327, 443, 489, 545, 649, 651, 740; relation of, to vein-mosaic of leguminous plants, 12; Hippodamia spp. predacious on, 128: technique for rearing Coccinellids 649: measures frozen, against, 128, 285, 291, 327, 355. 443, 651, 740; nicotine fumigation tests on, 21.

Macrosiphum pisi (see M. ono-brychis).

Macrosiphum rosae, ants associated with, in Germany, 195.

Macrosiphum sanborni, tests of nicotine fumigation on, on chrysanthemum in U.S.A., 20.

Macrosiphum solani, on potato in Britain, 527.

Macrosiphum solanifolii, on potato in Br. Isles, 48, 527; in Italy, 759; in U.S.A., 12, 286; measures against, on sweet-potato, 286; relation of, to virus diseases of plants, 12, 48, 759.

Macrosteles, synonomy and key to N. American species of, 78.

Macrosteles divisus (Cicadula sexnotata, auct.), in U.S.A., 78, 288, 386; transmitting yellow disease of aster, etc., 78, 386; on narcissus, 288.

Macrosteles (Cicada, Cidadula) sexnotatus, type of genus, 78; M. divisus recorded as, in America,

78.

Macrotermes bellicosus, damaging clove seedlings in Tanganyika, 642.

macula-alba, Ceuthorrhynchus. maculata, Epicauta; Nephrotoma (Pachyrhina).

maculatus, Bruchus (Callosobruchus); Elasmus; Graphogaster; Thripoctenus.

maculifera, Sugitania. maculipennis, Plutella. maculiventris, Podisus. maculosa, Nemorilla floralis.

maculosum, Anthidium.

Madagascar, pests of field and stored cassava in, 454; new Aleurodid possibly transmitting cassava mosaic in, 413, 455, 706; Papilio demodocus on Citrus in, 167; Cephonodes hylas on coffee in, 68; cotton pests in, 67; Locusta migratoria capito in, 67. 416; Parnara spp. on rice in, sugar-cane pests in, 66, 67; beneficial insects in, 67; attempted introduction of Tachinid into Mauritius from, 176; rotenone-containing plants in, 67, 182, 183.

Madeira, Coccids of, 560.

Madras, miscellaneous pests in, 409, 499; Cecidomyiid on Moringa in, 603; rice pests in, 360; pests and disease of Sorghum in, 408, 531; sugar-cane pests in, 408, 602; utilisation of parasites of pests of teak and mahogany in, and their introduction into Burma, 339, 765; other beneficial insects in, 408, 499, 602, 603.

Magdalis armicollis, on elm in U.S.A., 14, 15, 325; Ceratostomella ulmi cultured from, 325; parasites of, 14, 15.

Magdalis barbita, parasites of, on elm in Connecticut, 14, 15.

Magicicada septendecim, in U.S.A., 81, 395, 726; bionomics of, 726;

fungi infesting, 81, 395.

Magnesium Arsenate, dusting with, against insect pests, 333, 478, 691; tests of, against cowpea weevil, 572, 691; relation of pH of insects to toxicity of, 741, 742; and barium fluosilicate, 478.

custs, **76**, **77**; aeroplane dusting with, **77**. Magnesium Arsenite, against lo-

Magnesium Carbonate, 248: trees banded with, against ants, 93: beet seed treated with, against mites, 160; as a carrier for anabasine dust, 236.

Magnesium Oxide, as an insecticide.

Magnesium Silicate (see Talc).

Magnesium Sulphate, tests of, in baits for grasshoppers, 608, 609; and sodium cyanide, against 630; wireworms, increasing yield of beet, 631.

Mahogany, Hypsipyla robusta on, in India, 765.

Mahonia aquifolium, not attacked by Rhagoletis cerasi, in Switzerland, 156.

maidiradicis, Anuraphis.

maidis, Aphis (Rhopalosiphum); Peregrinus.

Maine, miscellaneous pests in, 332, 484, 487, 580; light-trap studies

of Lepidoptera in, 746.

Maize, pests of, in S. Africa, 106, 364, 411, 511; Aphids on, in Argentina, 289, 290; pests of, in Australia, 362, 494, 755; pests of, in Canada, 441, 486, 536; Agrotis ypsilon on, in Celebes, 337; Tettigoniid on, in China, 244: Acridid on, in Formosa, 769; pests of, in India, 241, 532; Hypomeces squamosus on, in Indo-China, 186; pests of, in Italy, 713; pests of, in Loochoo Loochoo Islands, 264, 437; Sesamia vuteria on, in Madagascar, 67; 437; Sesamia Lema tristis on, in Manchuria, 767; S. vuteria on, in Morocco, 500, 501; Carabid on, in New Zealand, 361; Pyrausta nubilalis on, in Poland, 622; pests of, in Porto Rico, 39, 40; Heliothis armigera on, in N. Rhodesia, 366: pests of, in S. Rhodesia, 203, 366, 608; insects on, in Spain, 695, 700; H. armigera on, in Sudan,

511; pests of, in Russian Union, 424, 475, 476, 530; pests of, in U.S.A., 14, 44, 63, 65, 78, 124, 125, 126, 140, 144, 271, 279, 285, 304, 326, 329, 367, 391, 489, 505, 509, 545, 564, 580, 643, 644, 648, 723, 726, 729, 730, 745; spiders and insects associated with, in Ohio, 590; insects and diseases of, 386, 532, 736; time of sowing and varieties of, in relation to pests, 14, 63, 125, 144, 391, 326, 536, 564, 713, 729; effect on Euxoa segetum of feeding on, 424; as a trap-crop, 494, 713; Ultrawet causing injury to, 285; products of, in baits for grasshoppers, 410, 608, 609, 682.

Maize (Stored), pests of, and their control, 11, 414, 608, 643, 644, 746; effect of fumigants on germination of, 390, 414; (meal), Oryzaephilus surinamensis in, 369.

Maize Weevil (see Calandra oryzae). majalis. Amphimallus (Amphi-

mallon).

major, Penthaleus; Thrips.

Malachite Green, for preventing discoloration of palm foliage by lead arsenate, 22.

Malachra, pests of, in Peru, 533.

Malacosoma americana, on fruit and shade trees in U.S.A., 16, 542, 696; measures against, 16, 542.

Malacosoma disstria, in New Brunswick, 227; in U.S.A., 29, 45, 227, 580; new parasite of, 227; value of lime-sulphur against, in orchards, 29; on forest trees, 45, 580.

Malacosoma neustria, on fruit trees and oak in Czechoslovakia, 657; question of using derris against, in Holland, 529.

Malacosoma neustria testacea, bionomics of, in Korea, 264; on fruit trees in Japan, 54; in lighttraps, 54.

Maladera holosericea, outbreak and control of, in Germany, 194; soil conditions in relation to, in Russia, 429.

Malaya, new Capsid on Areca in, 198; forest pests in, 109, 110, 295, 453; Lyctid in timber in, 763; Oryctes rhinoceros in, 434; insects associated with copra and stored rice in, 31; new species of Telenomus in, 89, 688; other beneficial insects in, 764; summary of plant quarantines in, 229; Spalangia introduced into Ceylon from, 359.

Malayan Powder-post Beetle (see Minthea rugicollis).

malayensis, Platypus suffodiens. Maleuterpes spinipes (Phytolymus),

Maleuterpes spinipes (Phytolymus), on Citrus in New South Wales, 755.

mali, Aphelinus ; Dasyneura ; Psylla. malinelus, Hyponomeuta padellus.

Mallow, Crepidodera rufipes on, in Britain, 612.

Malpighia glabra, Orthezia praelonga on, in Grenada, 752.

Malta, miscellaneous pests in, 469; summary of plant quarantines in, 134.

Malva (see Mallow).

malvella, Platyedra (Gelechia).

Mamestra pisi (see Polia).

mamillata, Agria.

Manchuria, Aphids of, 496; Cydia glycinivorella on soy beans and lupin in, 54, 310, 452, 606; Dendrolimus spectabilis on pines in, 672; Lema tristis on cereals in, 767; list of insect pests in, 304; Braconid parasites of Lepidoptera in, 296, 452.

mancus, Agriotes; Xyleborus. Mandarin (see Orange, Mandarin).

manducator, Alysia.

Manganese Arsenate, in sprays against thrips, 274, 552.

Manganese Arsenite, in sprays against gladiolus thrips, **398**, **399**. Mangel, pests of, in Britain, **296**, **527**; Cirphis unipuncta on, in

New Brunswick, 441.

mangifera, Symitha. mangiferae, Coccus.

Mango (Mangifera), Dacus ferrugineus dorsalis on, in Bonin Islands, 672; Coccids on, in Ceylon and Florida, 359, 587; pests of, in India, 361, 601; pests of, in Jamaica, 490; Anastrepha ludens on, in Mexico, 142; Dacus cucurbitae on, in Philippines, 181; new Noctuid on, in Seychelles, 405; Apate monacha on, in Sierra Leone, 240.

Mango (Fruit), heat treatment of, against *Dacus dorsalis* in Formosa, **207.**

Mango Hoppers (see Idiocerus).

Mango Shield Scale (see Coccus mangiferae).

Manihot, Erinnyis spp. on, in Cuba, 171.

Manihot utilissima (see Cassava). manihotis, Bemisia. manilae, Scolia (see Campsomeris marginella modesta).

manilensis, Locusta migratoria. maniosa, Solenopsis geminata. Manitoba, grasshoppers in, **540**.

Maple, Dasychira pudibunda on, in Germany, 69; pests of, in Russia, 75, 433; pests of, in U.S.A., 126, 226, 543.

maplei, Aenasius.

marantica, Epicampoptera (Meta-drepana).

margaritosa, Lycophotia.

Margaronia caesalis, on Artocarpus integer in Ceylon, 358.

Margaronia laticostalis, alternative host of parasite of teak pests in India, 765.

Margaronia hyalinata (see Diaphania).

Margaronia unionalis, on olive in Cyprus, 108.

marginalis, Orthotylus.

marginata, Mesogramma.

marginatus, Melanoplus; Syromastes (Mesocerus).

marginella, Campsomeris; Dichomeris (Nothris).

mariana, Eulia.

Marianne Islands (see Japanese Mandated Islands).

Marigold (see Calendula officinalis). maritimus, Pseudococcus.

Marl, Green-sand, effect of, on action of lead arsenate, **324**. marlatti, Aleurolobus; Aphelinus.

maroccanus, Dociostaurus.

Maruca testulalis (on leguminous plants), in Brazil, 642; in Java, 110; in Porto Rico, 643; parasites of, 111.

Maryland, pests of greenhouse and ornamental plants in, 541, 552, 553; Popillia japonica in, 444; Pyrausta nubilalis in, 730; parasites of Phytomyza ilicicola in, 541.

marylandensis, Tetrastichus.

Masicera silvatica, parasite of Aglaope infausta in Spain, 697.

maskelli, Pteroptrix.

Massachusetts, cranberry pests in, 645, 646; forests pests in, 547, 576; miscellaneous pests in, 646-648; *Pyrausta nubilalis* in, 648, 729, 730, 731; beneficial insects and biological control in, 647, 648, 729, 730; Nematodes in grasshoppers in, 123, 124.

Massospora cicadina, infesting Magicicada septendecim in Michigan, 81.

Mastotermes, measures against, damaging timber in Queensland,

materna, Othreis (Ophideres).

mathias, Parnara.

Matsucoccus, on pines in U.S.A.,

matsuyadorii, Pteromalus.

Mattesia dispora, possibly infesting Plodia interpunctella, 598.

maura, Eurygaster

mauritanicus, Tenebroides.

mauritia, Spodoptera.

Mauritius, Coccids on coconut in, 66, 176, 204; sugar-cane pests in, 176, 316, 317, 454; Technomyrmex albipes in, 204; beneficial insects and biological control in, 66, 176, 204, 316, 317, 454.

maxillosum, Psalidium

May-flies, nymphs of, damaging timber under water in Siam, 149.

Mayetiola destructor (on cereals), in Cyprus, 108; in Morocco, 340, 640; in U.S.A., 79, 121, 126, 129, 326, 329, 728; loss due to, in Indiana, 126; susceptibility of varieties of wheat and grasses to, 326, 728; bionomics of, 126, 640; parasites and biological control of, 121, 329, 340.

mayri, Monomorium gracillimum; Sphaerakis (Bruchobius).

mbila, Cicadulina. Meal Moth, Indian (see Plodia interpunctella).

Mealy Plum Aphis (see Hyalopterus arundinis).

Mealybug, Avocado (see Pseudococcus nipae).

Mealybug, Grass (see Antonina

indica). Mealybug, Grey Sugar-cane (see Pseudococcus boninsis).

Mealybug, Pineapple (see Pseudococcus brevipes).

Meat, in baits for ants, 222, 223; (cured) Piophila casei in, 38.

Mecaspis (see Cleonus). medicaginis, Contarinia.

Medicago denticulata, Smynthurus viridis on, in Victoria, 84.

Medicago falcata, Contarinia medicaginis on, in Sweden, 232.

Medicago lupulina, probably primary food-plant of Phytodecta fornicata in Jugoslavia, 760, 761.

Medicago sativa (see Lucerne).

mediocris, Agrilus.

Mediterranean Fruit-fly (see Ceratitis capitata).

Medlar, Lymantria dispar on, in

Jugoslavia, **759.**Megacephala carolina, predacious on Anomis luridula in Peru, **627.** megacephala, Pheidole.

megacephalus, Microphanurus (see M. basalis).

Megachile centuncularis, damaging plants in Germany, 522.

Megachile seychellensis, damaging Derris and roses in Seychelles, 433.

Megamelus proserpina (Taro Leafhopper), bionomics of, in Hawaii and Philippines, 212.

Megaselia plurispinosa, measures against, on mushrooms in Holland,

655.

Megatoma varius, bionomics and control of, in Japan, 514.

Melaloncha ronnai (parasite of honey-bees), bionomics of, in Brazil, 115, 655; bait for, 115.

Melampodothrips, new subgenus of Podothrips, 457.

Melanagromyza (see Agromyza).

melanictera, Enaria,

Melanocallis caryaefoliae, on pecan in Florida, 587.

melanocephala, Diabrotica. melanochra, Stathmopoda. melanoderes, Dysdercus. melanogaster, Drosophila.

melanopa, Lema.

Melanophila californica, oil sprays against, on Pinus ponderosa in California, 394.

Melanophila picta, on poplar and ash in Russian Union, 429.

Melanoplus, ecology and effects of cultivation on species of, in N. America, 680, 681; baits for, 23, 139, 540, 574.

Melanoplus bivittatus, in Canada, 680; in U.S.A., 19, 23, 139, 680,

Melanoplus devastator, in California, 19.

Melanoplus differentialis, in U.S.A., 23, 139, 574, 681; effects of hypertonic solutions and cyanide on eggs of, 304.

femur-rubrum, in Melanoplus Canada, 680; in U.S.A., 23, 580.

Melanoplus marginatus, in California, 19.

Melanoplus mexicanus, in Canada, 540, 680; in U.S.A., 23, 44, 139, 580, 680, 681.

Melanoplus packardi, in California, 19; in Canada, 680.

Melanostoma, Empusa muscae infesting, in Britain, 101.

Melanostoma mellinum, 101.

Melanotus tamsuyensis, on sugarcane in Loochoo Islands, 437. melanura, Nacerda.

Melaphis chinensis, bionomics and utilisation of, producing tannin-

galls in Far East, 264.

Melasoma, Coccinellid predacious on, in Korea, 310.

Melasoma adamsi, in Korea, 310. Melasoma populi, in Korea, 310.

Melasoma saliceti, trees attacked by, in Russian Union, 429.

Melasoma vigintipunctata, ir. Korea, 310.

Melcha ornatipennis, parasite of Scirpophaga in India, 243. meles, Hypera (Phytonomus).

Melhania ferruginea, cotton stainers on, in Uganda, 373.

Melia, Coccids on, in Nyasaland, 114.

Melia azadirachta (Neem), leaves of, as repellent for *Ephestia* in stored cacao, **315.**

Melia azedarach, Phassus signifer on, in Formosa, 517; Leptocneria reducta on, in New South Wales, 582; extract of, as repellent spray for locusts, 180.

Meliana flammea var. stenoptera, on Juncus in Japan, 672.

Melicocca bijuga, Apate terebrans on, in Jamaica, 491.

Meligethes aeneus (on crucifers), in Britain, 296; in Czechoslovakia, 656; in Holland, 299; in Hungary, 197; measures against, 197, 299.

Melilotus (see Sweet Clover).

Meliphora grisella (see Achroia).

Melipona, damaging bananas in
Trinidad, 10.

Melipona testacea cupira, damaging Citrus in Peru, 82.

Melitomma insulare, measures against, on coconut in Seychelles, 433

Melittia satyriniformis, on cucurbits in U.S.A., 269, 647, 725; measures against, 647.

mellinum, Melanostoma.

Mellissopus latiferreanus (see Cydia).

mellitor, Microbracon. mellonella, Galleria.

Meloë spp., attacking honey-bees in Hungary, 344.

Melolontha, in Germany, 251, 345, 346, 520, 599; birds destroying,

468, 520; fungus disease of, 193, 468; methods of ascertaining flight years of, 154; organised collection of adults of, 193, 346, 468, 599, 722; electrocuting light-trap for, 251; methods of protecting fields from oviposition by, 251, 346, 721; measures against larvae of, 345, 432, 520.

Melolontha hippocastani (in forests), in Belgium, 193; in Denmark, 468; in Germany, 251, 253, 312; in Latvia, 90, 722; in Poland, 154; in Russia, 158, 429, 430, 432; factors affecting distribution of larvae of, in soil, 158, 429, 430; bionomics of, 154, 193, 253; influence of restricted space on, 194.

Melolontha melolontha (vulgaris), in Denmark, 468; in Germany, 251, 253, 312, 346, 468, 717; in Latvia, 90; in Poland, 154; in Rumania, 721; in forests, 90, 154, 312, 721; bionomics of, 253, 717, 721.

Melon, Rhaphidopalpa foveicollis on, in Algeria, 371; Dacus spp. on, in Formosa, 206, 207, 667; D. cucurbitae on, in Hawaii, 325; pests of, in India, 765; Epilachna chrysomelina on, in Morocco, 417; Myiopardalis pardalina on, in Palestine, 77; D. cucurbitae on, in Philippines, 181; pests of, in U.S.A., 269, 647, 748.

Melon Fly (see Dacus cucurbitae). membranaceus, Brachytrypes; Leptoglossus.

mendax, Lygidea. mendesi. Dvsdercus.

mendicus, Cleonus (Conorrhynchus). Meraporus, parasite of Mayetiola destructor in Morocco, 340.

mercator, Oryzaephilus surinamensis. mercurius Proteides

mercurius, Proteides.

Mercury Compounds, uses of, against Diptera infesting vegetables, 298, 536, 647, 770. merdigera, Lilioceris.

meriopterum, Heterotoma.

Merisus spp. (parasites of Mayetiola destructor), competition of, with Platygaster herricki in U.S.A., 121.

Mermis nigrescens, parasite of Hieroglyphus banian in India, 55.

Mermis subnigrescens (parasite of grasshoppers), in U.S.A., 123, 124, 746; bionomics and utilisation of, 123, 124; parasitising honey-bee, 746.

Merodon equestris, on narcissus, in New York, 393.

Meromyza americana (Wheat Stem Maggot), in Canada and U.S.A., 121; bionomics of, in Kansas, 121-123.

meromyzae, Coelinius (Coelinidea): Microbracon.

Meroprion perseae, sp. n., on avocado in Brazil, 502.

Mesocerus marginatus (see Syromastes).

Mesogramma marginata, predacious on Thrips tabaci in Bermuda, 626.

Metadrepana marantica (see Epicampoptera).

Metagonistylum mixense (parasite of Diatraea), in Brazil, 82; in Br. Guiana, 10, 221, 369, 491; establishment of, in W. Indies, 10, 82, 121; of little value against D. canella, 221, 491; Thysanus dipterophaga parasitic on, 491; forms of, 82.

Metal Shields, against termites in buildings, 17, 724.

metallica, Pleurotropis.

Metarrhizium anisopliae, infesting Alissonotum impressicolle Burma, 294; compost-pit temperatures affecting control of Oryctes by, in Čeylon, 434; experiment with, infesting Magicicada septendecim in 395.

Metatetranychus ulmi, auct. (see Paratetranychus pilosus).

metathoracica, Amauromorpha.

Meteorological Conditions (see Humidity, Rainfall and perature).

Meteorus, parasite of Anomis luridula in Peru, 627.

Meteorus scutellator, parasite of Cirphis unipuncta in Manchuria, 296.

Meteorus trachynotus, parasite of Sparganothis sulfureana in New Jersey, 563.

Meteorus versicolor, establishment of, against Stilpnotia salicis in U.S.A., 590; Lymantria monacha surviving parasitism by, 233.

Methyl Alcohol, tests of, against cheese mites, 539.

Methyl Bromide, tests and uses of, as a fumigant, 18, 324, 388–390, 414, 539, 541, 571, 572, 746, 747; regulations dealing with fumigation of strawberry plants and potatoes with, against Popillia japonica in U.S.A., 384, 751; action of, on man and animals, 389, 414, 746; and carbon dioxide, 539, 571, 572; determination of small quantities of, in air, 695.

Methyl Formate. and carbon dioxide, fumigation with, against Tribolium castaneum, 571.

Methyl Iodide, tests of, as a fumigant, 575.

Methylene Blue, for distinguishing Coccids affected by HCN, 770.

(Phlogometiculosa, Brotolomia phora).

meticulosalis, Terastia.

Mexican Bean Beetle (see Epilachna varivestis).

Mexican Fruit-fly (see Anastrepha ludens).

mexicanus, Melanoplus.

Mexico, Anastrepha spp. in, 142, 325, 505; Epicaerus cognatus on potato in, 11; miscellaneous pests in, 28, 219, 248, 395; parasites and biological control of cotton pests in, 274, 275; parasites of mealybugs in, 212, 586: development of economic entomology in, 248; introduction of beneficial insects into Hawaii from, 586.

michaeli, Eberhardia; Lepidogly-

phus (Glycyphagus).

Michigan, Camnula pellucida in, 681; forest pests in, 80; orchard pests in, 79, 733, 743; Pyrausta nubilalis on maize in, 580, 730; Magicicada fungus infesting septendecim in, 81.

Micrapate brasiliensis, boring in lead sheathing of cables in Mon-

tevideo, 232.

microbolista, Odites.

Microbracon, parasite of Stictodiplosis moringae in Madras, 603.

Microbracon analcidis, parasite of Tyloderma fragariae in Kentucky,

Microbracon brevicornis, failure to establish, against Platyedra gossypiella in Mexico and Texas, 275, 276.

Microbracon celer, variety of, parasitic on Ceratitis nigra in Kenya,

492.

Microbracon cushmanni, parasite of Desmia funeralis in California, 19, 285; destroyed by calcium cyanide, 286.

Microbracon gelechiae, parasite of Sparganothis sulfureana in New

Jersey, 563.

Microbracon hebetor, parasite of Platyedra gossypiella in Brazil, 704; bionomics of, parasitising Ephestia elutella in France, 524; effects of sting of, on other species of Ephestia, 12, 29.

Microbracon intercessor (parasite of Lixus spp.), in Italy, 528; in Morocco, 417, 526; bionomics

of, 526.

Microbracon kirkpatricki, doubtful value of introduction of, against Platyedra gossypiella in Mexico and Texas, 275, 276.

Microbracon lefroyi, parasite of Earias spp. in India, 243.

Microbracon mellitor, parasite of Platyedra gossypiella and Anthonomus grandis in Mexico, 275; parasite of Sparganothis sulfureana in New Jersey, 563; introduction of Hawaiian strain of, into Texas against P. gossypiella, 276; rearing of, 275.

Microbracon meromyzae, parasite of Meromyza americana Kansas, 123.

Microbracon pembertoni, parasite Argyroploce illepida Hawaii, 497.

Microbracon piger, imported into U.S.A. against Etiella zinckenella,

Microbracon platynotae, hosts and utilisation of, in Mexico, 275.

Microcerotermes parvus, damaging trees in Tanganyika, 113.

Micrococcus, notice of list of species of. 641.

Micrococcus silvestrii, in ants' nests, 642; characters of, 641.

Micrococcus similis, on cereals in Morocco, 641, 642; characters of, **641.**

Microcryptus annulationus, sp. n., parasite of Diprion spp. in Japan, 295.

Microcryptus opacus, sp. n., parasite of Diprion nipponicum in Japan, 296.

Microdontomerus anthonomi, parasite of Bruchus pisorum in Idaho,

Microdontomerus fumipennis, parasite of Tortrix rosaceana in Utah.

Microdus, parasite of Diatraea saccharalis in U.S.A., 391.

Microdus diversus, liberation of, against Cydia molesta in Connecticut, 16, 724.

Microdus glycinivorellae, n., sp. parasite of Cydia glycinivorella

in Manchuria, 452.

Microdus pumilus (parasite of Coleophora laricella), in Austria, 660; establishment of, in U.S.A., 745.

Microdus stigmaterus (parasite of Diatraea saccharalis), in Peru, 391; establishment of, in U.S.A., 391.

parasite of Cydia Microgaster, molesta in Italy, 154.

Curt. bicolor, (see Microgaster Macrocentrus abdominalis).

Microgaster bicolor. Nees (see Apanteles circumscriptus).

Microgaster globata, parasite Pyrausta nubilalis in Poland, 623. micrographus, Pityophthorus (Pity-

ogenes).

Microphanurus basalis (parasite of Nezara viridula), bionomics of, in Egypt, 457; establishment of, in W. Australia, 458.

Microphanurus megacephalus M. basalis).

Microphanurus semistriatus, bionomics of, parasitic on Pentatomids in Switzerland, 92.

Microplectron fuscipenne (parasite of Diprion spp.), establishment and utilisation of, in Canada, 77, 487, 538; parasite of, in Hungary, 341; imported into U.S.A., 329; bionomics of, 77, 78.

Microtermes redenianus, soil fumigation against, damaging plants

in Tanganyika. 113.

Microthrips piercei, on carrot in Hawaii, 692.

Middlings (see Bran).

migratoria, Locusta.

migratorioides, Locusta migratoria. mikado, Brachymeria (Chalcis): Euphranta.

miles, Cerambyx.

militaris, Apanteles; Calotermes (Neotermes).

Milk (Dried), Attagenus piceus feeding on, in U.S.A., 568; as a spreader and adhesive for sprays, 489, 490, 548, 566, 743, 750; dust formula containing, 566.

Millepedes, plants attacked by, in Germany, 461, 468; measures against, 461.

Millet (see Panicum miliaceum). Millet, Italian (see Setaria italica).

Mimela testaceipes, bionomics and control of, in Korea, 436.

mimus, Dysdercus.

Mindarus abietinus, on silver fir in Germany, 252, 253,

minense, Metagonistylum.

Mineola vaccinii, on cranberry in U.S.A., 172, 563, 646; bionomics of, 172; measures against, 172, 646.

minimus, Dicyphus; Scymnus (see Stethorus punctillum); Spheno-phorus (Calendra).

Minnesota, Camnula pellucida in, 681; Nematode parasite of grasshoppers in, 123; Eriosoma crataegi in, 484; new Eulophid parasite of Malacosoma disstria in, 227; insects and fungi in Pinus resinosa in, 43, 386; Lygus pratensis on potato in, 217.

minor, Cremastus; Euproctis; Myelophilus (Blastophagus)

Mint, Longitarsus spp. on, in Poland, 255.

Mint (Dried), Borkhausenia pseudo-spretella in, in France, 412.

Minthea obsita, in stored cassava in Madagascar, 455.

Minthea rugicollis, in timber in Burma and Malaya, 339, 763; in stored cassava in Madagascar, 455; bionomics and distribution of, 763; measures against, 764. minuscula, Clania.

minuta, Charips; Peronea. minutissimum, Trichogramma (see T. minutum).

minutum, Trichogramma. minutus, Platygaster (see P. hiemalis).

mirabilis, Aiolocaria (Ithone).

Miridiba koreana, measures against, in Korea, 436.

Miscanthus, Lepidosaphes sacchari restricted to, in Formosa, 263.

Miselia inferior, on tobacco in

Tanganyika, 642.

Mississippi, cotton pests in, 277, 278, 328, 329; grasshoppers in, 574; Naupactus spp. in, 580; new parasites of Psallus seriatus in, 227.

Missouri, pests of cucurbits in, 269; miscellaneous pests in, 23, 79; Nematode parasite of grasshoppers in, 123.

mistum, Succingulum.

Mites, attacking insects, 38, 84, 95, 112, 128, 147, 162, 163, 169, 343, 629, 655; attacking other mites, 15, 95, 97, 160; natural enemies of, 71, 465, 674, 712, 724; classification and new species of, 94, 95, 103, 163, 173, 436, 465, 470, 530.

mitis, Zenillia (Exorista).

mixtus, Lepidoglyphus (Glycyphagus) destructor.

Mnium, Melaphis chinensis on, in Korea, 264.

modesta, Campsomeris marginella. modestor, Opius perproximus.

moestum, Anacridium.

Mohelnaspis moravica, gen. et sp. n., on Festuca glauca in Czecho-

slovakia, 560.

Molasses, in baits, 28, 51, 108, 125, 168, 170, 209, 236, 270, 282, 393, 423, 473, 488, 506, 540, 544, 574, 682, 685, 686, 713, 727; in baitsprays, 50, 51, 61, 71, 325, 505, 619, 700, 713; addition of, to other sprays, 131, 138, 169, 222, 447, 484, 546, 550, 647, 682; other uses of, with insecticides, 494, 644, 645; effect of, on insecticide extracts of plants, 361, 505; causing injury to begonias, 484; types of, 71, 506, 700.

Mole-cricket (see Gryllotalpa).

Mole-cricket, West Indian Scapteriscus vicinus).

molesta, Cydia (Grapholitha, Laspeyresia).

molestae, Angitia (Dioctes). molitor, Tenebrio.

mollifera, Eupterote.

mollipes, Draeculacephala.

mollis, Ernobius.

Moma alpium, measures against, on oak in Russia, 443.

Momordica charantia, Epilachna vigintioctopunctata on, in Ceylon,

monacha, Apate; Lymantria.

monachae, Agria phaga); Elachertus. (Pseudosarco-

xanthophilum, Monalonion cacao in Brazil, 643.

moniezi, Caloglyphus.

Monieziella entomophaga, in stored cereals in Russian Union, 95.

Monoamylamine Oleate, contact performance of, 45.

Monochamus centralis, on coffee in Tanganyika, 113.

Monochamus notatus, fungus possibly associated with, in pine logs in Minnesota, 43.

Monochamus sartor var. rosenmülleri, on conifers in Japan, 517.

Monochamus scutellatus, fungus possibly associated with, in pine logs in Minnesota, 43.

Monochloronaphthalene, 125. (See Naphthalenes, Chlorinated.)

Monocrepidius (see Conoderus). Monodoniomerus aereus, parasite of Lymantria dispar in Jugoslavia, 760.

Monodontomerus dentipes, parasite of Dendrolimus spectabilis Japan, 669.

Monohammus (see Monochamus). Monolepta australis, dusts against,

on plants in New South Wales, 755.

Monolepta gossypii, sp. n., on cotton in Nyasaland, 372.

Monolepta rosea, measures against, on Citrus, in Queensland, 268.

Monolexis, parasite of Minthea rugicollis in Malaya, 764.

Monomorium carbonarium ebeninum, predacious on Diatraea saccharalis in Porto Rico, 288, 289; fostering Sipha flava, 289.

Monomorium destructor, damaging electric cables, etc., in Netherlands Indies, 262.

gracillimum Monomorium mayri, damaging electric cables in Netherlands Indies, 262.

(Monophadnoides) Monophadnus rubi, measures against, on raspberry in Montana, 45.

Monosteira unicostata, on pear in Libya, 260; bionomics of, in Morocco, 340.

monostigma, Scirpophaga. Montana, Blapstinus substriatus in, 26-28; grasshoppers in, 43, 609, 680, 681; miscellaneous pests in, 44, 170, 581; beneficial insects in, 44.

montana, Ochrophara.

Montezuma speciossima, Platyedra gossypiella on, in Porto Rico, 43. monticola, Actia.

monticolae, Dendroctonus.

montrouzieri, Cryptolaemus. Montserrat, resting-stage of 83; Platyedra gossypiella in, summaries of plant quarantines in, 134, 491.

moravica, Mohelnaspis.

morbillosus, Cleonus.

mori, Bombyx.

Moringa, pests of, in India, 55, 405,

moringae, Noorda; Stictodiplosis. Mormidea poecila, on rice in Br. Guiana, 369.

Mormon Cricket (see Anabrus simplex).

Mormoniella vitripennis, factors concerned in host-finding by, 150.

Morning Glory (see Ipomoea).

Morocco, Aphids of, 48, 696; beet pests in, 416, 417, 526; cereal pests in, 340, 500, 640, 641; pests of cork oak in, 46; weevil on leguminous crops in, 641; locusts in, 180; Epilachna chrysomelina on melon in, 417; orchard pests in, 340, 341; beneficial insects in, 340, 417, 526; summaries of plant quarantines in, 134, 385, 692.

morogorensis, Allodontermes.

Moropactus lepidotus, gen. et sp. n., on cacao in Brazil, 405.

Moropactus perfidus, sp. n., on cacao in Brazil, 405.

morstatti, Xyleborus.

mosellana, Sitodiplosis (Clinodiplosis, Thecodiplosis).

Mosquito Larvae, tests of insecticides on, 617.

Mosaic Diseases, question of relation of Aleurodids to, in cassava, 240, 413, 455, 706; in Chinese cabbage, 577; in cucumber, 615; in leguminous plants, 12, 265; in narcissus, 288; in solanaceous plants, 759; in sugar-cane, 39, 40, 411, 577; relation of Aphids to, 12, 39, 40, 265, 411, 577, 615, 759.

mossambicus, Hodotermes.

Moth-proofing Materials, tests of, on Anthrenus vorax, 227.

Mountain Ash (see Sorbus aucuparia).

Mozambique (see Africa, Portuguese East). Mucuna huberi, Maruca testulalis

on, in Brazil, 642.

mughiana. Rhyacionia (Evetria) turionana.

Mulberry, Aonidiella aurantii on, in S. Africa, 455; dried-fruit moths breeding on fallen fruit of, in California, 30, 118, 324; Tettigoniid on, in China, 244; Tetranychus telarius on, in Formosa, 674; Pseudococcus nipae on, in Hawaii, 586; pests and disease of, in Japan, 208, 604; pests of. in Russian Union, 429, 530; Lymantria dispar not experimentally surviving on, 759.

multistriatus, Scolytus.

Mundulea, evaluation of samples of, 258.

content of, 183. Mundulea rotenone

Mundulea sericea (suberosa), insecticidal properties and constituents of, 34-36, 37, 361.

mundum, Bethelium.

muricatus, Sciaphilus S. (see asperatus).

murinus, Lacon.

murraea, Cassida.

Murraya, Coccid on, in S. Africa,

Musca domestica, infesting drying fruit in California, 228; tests of insecticides on, 25, 35, 567.

Muscina pabulorum, parasite of Lymantria dispar in Jugoslavia, 760.

Muscina stabulans, parasite of Lymantria dispar in Jugoslavia, 760; in unsound tomatos in U.S.A., 398.

muscosa, Lycophotia. musculosa, Oria.

musculus, Anthonomus.

Mushrooms, pests of, in Britain, 297, 712; pests of, in Holland, 655, 656; pests of, in U.S.A., 286, 328, 579; survey of pests and diseases of, 304.

musicus, Gastrimargus.

Mussel Scale, Citrus (see Lepidosaphes beckii).

Mustard, little attacked by Melolontha in Germany, 521.

Mustard Beetle (see Phaedon cochleariae).

Mycalesis gotama, on rice and bamboo in China, 674.

Mycodiplosis alternata, measures against, producing galls on Cornus in U.S.A., 546; insects associated with, 546.

mycophagus, Mégn., Caloglyphus (Isoglyphus, Tyroglyphus).

mycophagus (Mégn.) Oudm., Caloglyphus (see C. moniezi).

Mycophila speyeri, spray against, in mushroom houses in Holland, 655.

myeloenta, Apanteles.

Myelois ceratoniae, on carob in Cyprus, 106, 458; new parasite of, 106.

Myelophilus minor (on pines), in Estonia, 462; in Sweden, 634,

Myelophilus piniperda (on pines), in Estonia, 462; bionomics of, in Holland, 348, 349; in Sweden, 634, 635; measures against, 348. 634, 635.

Myiolia lucida (see Acidia).

Myiopardalis pardalina, in India, 242: on melon in Palestine. 77. Mylabris aperta, on Cajanus cajan in Tanganyika, 642.

Mylabris dicincta, destroying locust eggs in Nyasaland, 114; on Cajanus cajan in Tanganyika, 642.

Mylabris pertinax, destroying locust eggs in Nyasaland, 114.

Mylabris variabilis, destroying eggs of Dociostaurus maroccanus in Italy, 163, 593; bionomics of, 163.

Myllocerus discolor, on Gliricidia maculata in Ceylon, 358.

Myllocerus viridanus, on Erythrina in India, 149.

Myochrous armatus, on sweet potato in Barbados, 222.

Myrica rubra, Lebeda nobilis on, in China, 244.

myricae, Bemisia.

Myrmelachista ambigua ramulorum, measures against, in coffee plantations in Porto Rico, 222.

Mysore, Xylotrechus quadripes on coffee in, 55; miscellaneous pests in, 204, 361, 606; beneficial insects in, 204, 409; tests of insecticidal properties of various plants in, 204, 360; Trichogramma minutum introduced into Madras from, 765.

mytilaspidis, Aphytis (Aphelinus). Mytilaspis (see Lepidosaphes).

Mvzoides (see Mvzus).

Myzus circumflexus, effect of starvation on transmission of virus diseases by, 615.

Myzus ornatus, food-plants of in Britain, 297, 527, 589; in Bel-

gium and California, 589.

Myzus persicae, in S. Africa, 177; in British Isles, 31, 47, 342, 379, 527, 615, 709; in Canada, 144, 537; in Hawaii, 214; in India, 241, 765; in Japan, 265; in S. Rhodesia, 377; in Russian Union, 237; in U.S.A., 21, 286, 445, 564, 577; on potato, 144, 214, 709; relation of, to virus diseases of potato, 47, 379, 527, 564, 615; on other plants, 21, 48, 145, 177, 237, 241, 286, 537, 765; relation of, to virus diseases of other plants, 31, 68, 265, 342, 377, 445, 577, 615; detection of isolated plants by, 379; penetration of setae of,

225; parasites of, 537, 538; measures against, 177, 237, 241, 286, 564; susceptibility of, to nicotine fumigation, 21.

Myzus pseudosolani (see Macrosi-

phum solani).

N.

Nabis, predacious on Teleonemia lantanae in Australia, 87; species of, predacious on Eutettix tenellus in Utah, 128.

Nabis alternata, 128. Nabis capsiformis, 87.

Nabis ferus, 128.

Nabis rosipennis, 128.

Nacerda melanura, in a store in Connecticut, 726.

naevana, Rhopobota. nanella, Recurvaria.

nankingensis, Pimpla (Epiurus).

nanus, Rhynchites.

Naphthalene, mixture containing, against ants in timber, 219 dissolved in oil sprays against bark-beetles, 394; unsatisfactory against mites in beet seed, 160; as a repellent, 251, 346, 350, 400, 489; against pests in soil, 236, 327, 383, 393, 461, 512, 631, 645, 712, 747, 771; formula for spraying with, against Tipulo aino, 771; dusts containing, 350, 484; and carbon bisulphide emulsion, 747; and chloropicrin, 236; derivatives of, as fumigants, 648.

Naphthalene Sulphonic Acids. activators, wetters and spreaders containing, 237, 615, 704.

Naphthalenes, Chlorinated, bands treated with, against Cydia and Anthonomus, 613; and kerosene, crates dipped in, against Cydia, solution of, for treating timber against Lyctus, 190, 707.

Napomyza (see Phytomyza).

Naranga aenescens, bionomics and control of, on rice in Korea, 767; in Japan, 671; parasites of, 671, 768.

nararia, Natada.

Narcissus, Brithys crini on, in Formosa, 53; pests of, in U.S.A., 288, 393; mosaic disease of, 288. nasturtii, Contarinia.

Nasturtium, Aphids on, in U.S.A., 21.

nasuta, Provops.

Natada nararia, parasites of, on tea in Ceylon, 639.

Naupactus, in Mississippi, 580. Naupactus bondari, sp. n., on cacao

in Brazil, 106.

Naupactus leucoloma, on lucerne in New South Wales, 132; distribution of, in U.S.A., 131, 132, 580; bionomics and control of, 131, 132; digestive and reproductive systems of, 696.

navale, auct., Tribolium (see T.

castaneum).

Navomorpha spp., trees attacked by, in New Zealand, 266.

nawae, Holochlora.

nawai, Telenomus.

Nebraska, winter food-plant of Paratrioza cockerelli in, 578; tests of grasshopper baits in, 608. nebulosa, Cassida,

Necremnus leucarthros, parasite of Coleophora laricella in Austria,

660.

Necrobia rufipes, in stored cassava tubers in Madagascar, 455; associated with copra in Malaya, 31; in New South Wales, 451; predacious on other Coleoptera, 31, 451.

Nectarine, Macropsis trimaculata on, in N. America, 173; pests of, in New Zealand, 57, 266.

Nectarine (Dried), Plodia interpunctella in, in California, 30. Neem (see Melia azadirachta). neglecta, Lotis.

negundinis, Periphyllus.

Nematodes, infesting insects, 55, 123, 124, 142, 504, 623, 671, 746; types of association between insects and, 49, 86; new species of, 49, 504.

Nematospora, bugs associated with, on coffee and cotton in E. Africa,

178, 375, 376.

Nematus abietinus (see Pristiphora). Nematus abietum (see Pristiphora abietina).

Nematus dimmocki (see Pikonema). Nematus ruralis (see Pikonema). nemea, Zenillia (Phryxe).

Nemeritis, parasite of Cydia molesta in Italy, 154.

Nemeritis canescens, olfactory conditioning and host selection in, 1, 2; oogenesis in, 119.

Nemeritis palmaris, parasite of Tirathaba trichogramma in Fiji, 334.

Nemeritis tectonae, parasite of Xyleutes ceramicus in Burma, 339.

nemorata, Phyllotoma.

Nemorilla floralis, parasite of Sparganothis sulfureana in New Jersey, 563; parasite of Psanmotis hyalinalis in Jugoslavia, 760; parasitised by Theronia atalantae, 760.

Nemorilla floralis maculosa, parasite of Desmia funeralis in California, 286.

nemorum, Anthocoris; Phyllotreta. nenuphar, Conotrachelus.

Neoaplectana, parasite of Cydia nigricana in Quebec, 623.

Neoaplectana affinis, sp. n., bionomics of, in Denmark, 49, 50.

Neoaplectana bibionis, sp. n., bionomics of, in Denmark, **49, 50.** Neocatolaccus indicus, sp. n., para-

Neocatolaccus indicus, sp. n., parasite of Pempheres affinis in India, 89.

Neodiprion (see Diprion).

Neodiscoides, characters of, 212.

Neodryoczetes, in butter-nuts imported into Holland from Dutch Guiana, 529.

Neogastrallus librinocens, gen. et sp. n., infesting books in Florida, 329.

Neomyzaphis (see Rhopalosiphum). Neoproba varians (see Engytatus geniculatus).

Neoquassin, 120, 368.

neosalviae, Eriophyes.

Neosciara (see Sciara).

Neotermes (see Calotermes).

Neotetranychus buxi, on Buxus in Connecticut, 17.

nephantidis, Elasmus.

Nephantis serinopa (on coconut), in Ceylon, **359, 360;** in Cochin, **452;** utilisation of *Trichospilus pupivora* against, **359.**

Nephelium litchi (see Litchee).

Nephelium longana, Eriophyes on, in Formosa, 674.

Nephotettix bipunctatus (apicalis), cincticeps, in Japan, 54, 167, 439; transmitting dwarf disease of rice, 167, 439; parasites of, 439; caught in light-traps, 54, 439.

Nephrolepis, Coccids on, in Britain, 712.

Nephrotoma maculata, on carnations in Switzerland, 689.

Nerium, Asterolecanium pustulans restricted to, in Formosa, 263.

Nerium odorum, tests of insecticidal properties of, 360, 361.

nervosa, Depressaria; Hymenarcys.

Netherlands Indies, Cerambycids on clove trees in, 295; coconut pests in, 337, 499, 674; mealybugs on coffee etc. in, 182, 606, 607, 608; pests of leguminous plants in, 110, 261, 292; pests of malvaceous plants in, 106, 358, 765, 766; miscellaneous pests in, 205, 453, 457, 603; Coleoptera in stored products in, 311, 336, 434, 639; insects damaging books in, 706; ants damaging electric cables in, 262; new species of Coleoptera in, 106, 262, 292; new mite in ants' nests in, 95; Locusta migratoria manilensis in, 685; beneficial insects and biological control in, 89, 111, 337, 358, 688; summary of plant quarantines in, 385; introduction of beneficial insects into other countries from, 176, 204, 337, 639.

Nettle (see *Urtica dioica*).

neumanni, Locris.

Neurotol, bees not affected by dusting with, against forest pests, 69.

neustria, Malacosoma.

Nevis, resting-stage of *Platyedra* gossypiella in, 83; summaries of plant quarantines in, 491, 692.

New Brunswick, Cirphis unipuncta in, 441; forest pests in, 487, 539; new parasite of Malacosoma disstria in, 227; Myzus persicae on potato and Prunus in, 144.

New Caledonia, Tetrastichus giffardianus introduced into, 212.

New Guinea, Territory of, miscellaneous pests in, 435, 495; beneficial insects and biological control in, 435, 436.

New Hampshire, miscellaneous pests in, 20, 487; Nematode parasites of grasshoppers in, 124.

New Jersey, cranberry pests in, 508, 562, 563; Diorymerellus laevimargo on orchids in, 550; forest pests in, 394, 580; Coleoptera associated with Dutch elm disease in, 325; pests in nurseries in, 118; orchard pests in, 323, 385, 555, 737, 738; Pyrausta nubilalis in, 730; vegetable pests in, 28, 565; new Coccid in, 226; insects and virus diseases of plants in, 12, 385; beneficial insects and biological control in, 108, 323, 324, 325, 394, 555, 563.

New Mexico, Paratrioza cockerelli causing yellows of potato in, 281.

South Wales, Cosmopolites New sordidus on banana in, 210; insects on Citrus in, 56, 335, 502, 754, 755: miscellaneous pests in, 111, 209, 334, 404, 449, 582; orchard pests in, 56, 111, 267, 335, 448, 510, 608; vegetable pests in, 56, 209, 210, 334, 335, 755; vine pests in, 111, 335, 608; grasshoppers in, 440, 449, 450, 583, 584, 629, 676; Naupactus leucoloma in, 132; legislation respecting pests in, 267, 608; beneficial insects in, 111, 335, 450. 629 ; establishment of Encarsia formosa against Trialeurodes vaporariorum in, 449; introduction of Teleonemia lantanae against Latana in, 87, 212; beneficial insects introduced into other countries from, 702, 753.

New York, forests pests in, 80, 270, 329, 392, 487, 547, 548; Lepidoptera on maize in, 271, 564, 730; miscellaneous pests in, 136, 270, 287; orchard pests in, 25, 130, 269, 270, 288, 489, 555, 557–559, 561, 562, 574; pests of greenhouse and ornamental plants in, 273, 393, 490; vegetable pests in, 270, 564, 566, 731; new Coccids in, 226; new Thysanoptera in, 329, 490; insects taken on high buildings in, 215; beneficial insects and biological control in,

270, 273, 557, 558, 559.

New Zealand, pests of conifers in, 58, 266; pests of Eucalyptus in, 752-754; Carabid on maize in, 361; orchard pests in, 46, 56, 57, 58, 208, 247, 266; Phyllocoptes on tomato in, 45, 103; Leucodiaspis of, 303; beneficial insects and biological control in, 10, 57, 58, 208, 449, 582, 752, 753, 754; summary of plant quarantines in, 229.

Nezara chloris, bionomics and control of, on rice in Sierra Leone, 240.

Nezara viridula, in Australia, 246, 458; in Belgian Congo, 315; bionomics of, in Egypt, 457; on pecan and Citrus in Florida, 587, 727; in Java, 89; on cotton 315, 457; on papaya, 246; natural enemies and biological control of, 89, 457, 458, 727.

Nicaragua, summary of plant quarantines in, 134.

nicevillei, Glyptomorpha (Stenobracon). Nico-Zin, 79.

Nicofume Fumigating Powder, 20. nicolasi, Campylomma.

Nicotiana glutinosa, crucifer mosaic transmitted to, 578.

Nicotiana sylvestris, occurrence of l-nornicotine in, 183.

Nicotiana tabacum (see Tobacco).
nicotianae, Dicyphus (see Engytatus

tenuis). Nicotine, against Anasa tristis, 739; against Aphids, 20, 21, 32, 62, 177, 355, 387, 443, 523, 740, 741, 748, 762; against Capsids, 250, 534. 710: against Jassids, 554, 646, 652, 700, 701; and alcohol, etc., against Pseudococcus citri, 701; against cockroaches, 20; against Coleoptera, 265, 323, 421, **522, 566, 653, 691;** against Collembola, **656;** against Cydia spp., 62, 79, 100, 254, 272, 401, 442, 488, 544, 554, 555, 652, 653, 725, 734; against other Lepidoptera, 45, 172, 185, 270, 299, 323, 327, 421, 648, 664, 713, 723; against Diptera, 53, 62, 131, 313, 325, 328, 401, 467, 541, 711; ineffective against mites, 646, 693, 712; against plum sawfly, 614, 615; against thrips, 29, 129, 211, 291, 484, 711; injected into fruit trees against borers, 267; tests of toxicity and action of, on insects, 332, 439, 536; dusting with, 53, 129, 131, 177, 211, 327, 355, 484, 522, 534, 544, 565, 646, 691, 701, 710, 712, 723, 740, 741, 748; lime as carrier for, 646; fumigation with, 20, 21, 274, 313, 314, 711; duration of activity of sprays of, 421, 422; equipment for vaporising, 387; formulae containing 100, 172, 250, 272, 328, 355, 467, 534, 544, 615, 646, 656, 664, 700, 701, 711, 740, 762; and cubé, 740, 741; and derris, 172, 740, 741; and lead arsenate, 250; and lime-sulphur, 250, 291; and oils, 29, 45, 62, 185, 254, 270, 291, 368, 442, 555, 708, 711, 739; and soap, 32, 45, 62, 100, 291, 421, 422, 467, 536, 646, 700, 701, 710, 711, 713, 762; lethalate as a wetter for, 615; fixed compounds of (including nicotinebentonite, etc.), 79, 149, 272, 323, 368, 442, 448, 510, 554, 555, 556, 562, 566, 648, 652, 653, 723, 725, 732, 733, 734; determination of, **368, 456, 695**; standardisation of

insecticides containing, in Switzerland, 156; reviews of data on, 651; other insecticides compared with, 35, 62, 79, 255, 401, 442, 443, 522, 523, 544, 614, 652, 653, 723, 734.

Nicotine Laurate, in sprays against Sophronia humerella, 301.

Nicotine Oleate, fixed nicotine sprays prepared with, against Cydia and Chionaspis, 554, 555, 562, 733; semi-fixed spray of, 554; and oil emulsions, 554, 555; and Coposil, 554.

Nicotine Peat, preparaton of, as an insecticide, 24.

Nicotine Salicylate, toxicity of, to Aphids, 17.

Nicotine Soaps, spraying test with, against Cydia funebrana, 100.

Nicotine Sulphate, against Aphids, 17, 65, 210, 241, 260, 380, 399, 443, 704, 707; against Coccids, 65, 270, 361, 484, 549, 562; against other Rhynchota, 108, 260, 397, 437, 559, 690, 739; against Ceuthorrhynchus maculaalba, 659; against Cydia spp., 79, 149, 168, 442, 448, 479, 510, 554, 556, 652, 653, 732, 733; against other Lepidoptera, 9, 16, 138, 172, 301, 647, 713, 743; against Diptera, 131, 172, 335, 404, 527, 546, 636, 647; against mites, 46, 583; against thrips, 29, 88, 211, 355, 484, 508, 581, 626, 727; tests of toxicity of, to insects, 332, 390; dusting with, 211, 397, 399, 443, 508, 659; carriers for, 397, 399; greenhouse fumigation with, 46; formulae containing, 9, 29, 46, 65, 131, 138, 149, 168, 172, 210, 241, 260, 270, 335, 355, 380, 397, 399, 404, 442, 484, 508, 510, 546, 554, 556, 559, 562, 581, 647, 652, 732, 738, 743; and bentonite or bentonite sulphur, 79, 149, 448, 510, 554, 652, 653, 732, 733; and Bordeaux mixture, 131, 210; and Coposil, 554; and lead arsenate, 168, 210, 260, 442, 554, 555; and lime, 554; and lime-sulphur, 29, 131, 149, 727; and molasses, 131, 484, 546, 647; and nicotine oleate, 554, 556; and oils, 9, 16, 79, 131, 138, 149, 210, 260, 270, 335, 442, 448, 484, 508, 510, 546, 549, 554, 556, 559, 562, 647, 652, 708, 732, 733, 739, 743; and soap, 46, 65, 88, 260, 270, 355, 404, 549, 559, 704, 707, 733; other wetting and spreading agents for, 652, 732, 733; activators for, 238, 332; and sulphur. 149, 479, 732; and zinc sulphate. 138; analysis of insecticides for. 456; other insecticides compared with, 16, 17, 35, 79, 138, 149, 204, 225, 361, 554, 647, 652, 653, 690. 704, 732, 733.

Nicotine Tannate, uses of, as an insecticide, 648, 723.

Nicotine Thiocyanate, preparation and tests of, as an insecticide, 703, 704.

nidulans, Eupteromalus.

niger, Acromyrmex (Atta); Athous; Lasius.

Nigeria, cotton pests in, 666, 667; parasites of Trypetids in, 492; summary of plant quarantines in.

nigra, Ceratitis; Saissetia; Trichogramma pretiosum.

nigrescens, Mermis. nigribasalis, Pimpla (Itoplectis).

nigricana, Cydia (Laspeyresia).

nigricans, Hylesia.

nigricella, Coleophora (Eupista).

nigricephala, Encarsia.

nigricornis, Psila; Valanga.

nigricornis, auct., Phytomyza (see P. atricornis)

nigrifrons, Thamnotettix.

nigrimarginalis, Enicospilus. nigripes, Zeuxia.

nigriventris, Anagrus armatus.

nigrofasciatum, Lecanium. nigrofasciatus, Dysdercus.

nigrofuscata, Graptosaltria. nigropilosus, Thrips.

Nipa fruticans, Acanthopsyche spp. on, in Indo-China, 184; Promecotheca papuana on, in New Guinea Territory, 495.

nipae, Acanthopsyche; Pseudococcus.

Nipponaphis takahashii, new parasite of, on oak in Japan, 296.

nipponensis, Lophyroplectus.

nipponicum, Diprion.

Niptus hololeucus, 519; measures against, in Germany, 48.

nitens, Anaphoidea.

nitidalis, Diaphania. nitidiventris, Phytomyptera.

nitidula, Psilopa.

nitidulus, Anastoechus.

nitidus, Ephedrus.

nitobei, Chelonella (see Chelonus pectinophorae).

nivella, Scirpophaga.

nobilis, Cassida; Lebeda; Scutellera (see S. perplexa).

nocivus, Dicyphus (see Engytatus tenuis); Systates.

noctilio, Sirex.

noctiluca, Lampyris.

Noctua exoleta (see Calocampa).

nodiceps, Cryptognatha.

Nodonota puncticollis, bionomics and control of, in U.S.A., 287, 288.

Nodostoma bituberculatum, on roses

in Ceylon, 359.

Nomadacris septemfasciata, in Portuguese E. Africa, 676, 678, 682, 687: in S. Africa, 411, 676, 682; in S.W. Africa, 676; in Angola, 687; in Belgian Congo, 687; in Kenya, 679; in Nyasaland, 104, 114; in N. and S. Rhodesia, 114, 210, 383, 387; in Sudan, 112; in Tanganyika, 113, 642, 683; in Uganda, 112, 372, 679; outbreak centres of, 683, 684, 687; migrations of, 111, 112, 114, 678, 687; effects of temperature and humidity on, 104, 114, 147, 148; plants attacked by, 372, 411, 687; estimates of losses and expenditure due to, 676, 688; natural enemies of, 114, 682, 687; measures against, 410, 682.

Nonagria typhae (see Phragmatiphila).

nonvitta, Epibootania.

Noogoora Burr (see Xanthium pungens).

Noorda, measures against, on Moringa in India, 55.

Noorda blitealis, on Moringa in India, 55.

Noorda moringae, sp. n., on Moringa in S. India, 55, 405.

Nopoiulus, destroying carrots in Germany, 468.

nordmannianae, Chermes (Dreyfusia).

Norfolk Pine (see Araucaria excelsa). Nornicotine, laevo-, in Nicotiana sylvestris, 183.

norvegicus, Calocoris.

Norway, summary of plant quarantines in, 385.

Norway Pine (see *Pinus resinosa*). Nosema, in Cactoblastis cactorum in S. Africa, 202.

Nosema carpocapsae, sp. n., in Cydia pomonella in France, 464. notabilis, Aesiotes.

notatus, Bruchus (Bruchidius): Monochamus.

Nothobeus (see Coloborhombus). Nothris marginella (see Dichomeris).

Notolophus (see Orgyia).

Nova Scotia, Cirphis unipuncta in, 441, 540; Lepidoptera caught in light-traps in, 535; orchard pests in, 534, 535; Diprion polytomum on spruce in, 487.

noveboracensis, Ithycerus.

Novius (see Rodolia). noxius, Tyrophagus.

nubilalis, Pyrausta. nuciferus, Poecilips.

Nun Moth (see Lymantria monacha). nursei, Goryphus.

nüsslini, Chermes (Dreyfusia) (see C. nordmannianae).

Nutgrass Armyworm (see Laphygma

exempta).

new Coleoptera on Nyasaland, coffee in, 372, 405; cotton pests in, 113, 114, 177, 367; Gonipterus scutellatus on Eucalyptus in, 203; Aphis laburni transmitting rosette disease of ground-nuts in, 114; locusts in, 105, 114, 115; miscellaneous pests in, 114; beneficial insects in, 114, 204; summary of plant quarantine restrictions in. 229.

Nygmia flava (see Euproctis).

Nygmia phaeorrhoea, in Czecho-slovakia, 656, 657; in Germany, 660; in Holland, 183, 299, 347; in Russia, 432; in U.S.A., 560, 589, 723; on fruit trees, 299, 656, 660; on hawthorn, 183; on oaks, 183, 299, 347, 657, 660; on strawberry, 660; bionomics of, 347; parasites and biological control of, 589: other measures against, 183, 299, 347, 560; tests of duration of activity of nicotine preparations on, 421.

Nysius cymoides, food-plants and control of, in Cyprus, 108.

Nysius ericae, on potato and maize in Montana, 44.

O.

Oak (Quercus), pests of, in Czechoslovakia, 657; Tischeria complanella on, in Finland, 201; Pyrochroa coccinea on, in France, 316; pests of, in Germany, 50, 69, 344, 345, 420, 660, 664; Nygmia phaeorrhoea on, in Holland, 183, 299, 347; Cynipid on, in Hungary, 527; Phylloxera on, in Italy, 189; pests of, in Japan,

296, 311, 669; Tortricids on, in Ontario, 509, 510; pests of, in Poland, 256; new Coccids on, in Portugal, 315; pests of, in Rumania, 193, 721; pests of, in Russia, 75, 431, 432, 433; Tortrix postvittana on, in Tasmania, 8; pests of, in U.S.A., 80, 547, 550, 588; girdling to reduce starch content of timber of, against Lyctus, 635; fauna of litter of, 193; injured by calcium cyanide, 431; (timber) Coleopterous pests of, 190, 191, 635, 707; resistant to Hylotrupes and Codiosoma, 155, 354; Calotermes castaneus in, 220.

Oak, Cork (see Quercus ilex var.

suber).

Oak Leaf-roller (see Argyrotoxa semipurpurana).

Oak Processionary Caterpillar (see Thaumetopoea processionea).

Oats, Aphids on, in Argentina, 289, 290; pests of, in Britain, 382, 709; Cirphis unipuncta on, in Canada, 145, 441; Lema melanopa on, in Czechoslovakia, 657; pests of, in Germany, 51, 52, 251; attacked by Dociostaurus maroccanus in Italy, 162; pests of, in Morocco, 500, 501, 641; Penthaleus major on, in New South Wales, 582; pests of, in Russia, 422, 428; pests of, in U.S.A., 78, 140, 577.

Oats (Stored), pests of, and their control, 125, 414, 746; methyl bromide not impairing germina-

tion of. 390.

obesus, Alcides; Chlaenius. obfuscana, Epiblema.

oblineatus, Lygus pratensis. obliqua, Allograpta.

obliquana, Ctenopseustis.

Oblique-banded Leaf-roller Tortrix rosaceana).

obliquus, Listroderes. obliterata, Aphidecta.

oblongana, Harmologa. obovatus, Tenuipalpus.

obscura, Cytherea; Leucopis; Rhabdocnemis.

obscurata, Brachymeria. obscurator, Orgilus.

obscurus, Agriotes; Chrysomphalus; Hylastes (Hylastinus); Tenebrio.

obsita, Minthea.

obsoleta, Eleodes; Heliothis (Chloridea) (see H. armigera).

obtectus, Bruchus (Acanthoscelides, Bruchidius).

obtusa, Agromyza.

occidentalis, Copa; Pogonomyrmex; Tortrix (Cacoecia).

occiduus, Blissus.

occulta, Agrotis.

ocellana, Eucosma (Spilonota, Tmetocera).

ocellata, Anatis.

ocellatella, Phthorimaea.

ochracea, Oxyplax.

contraria, Ochrogaster enemies and control of, on Acacia in Australia, 449.

ochrogaster, Euxoa.

ochroleucus, Heptamelus.

Ochrophara montana, on bamboo in Burma, **339.**

Ochrosidia villosa, damaging lawns in U.S.A., 398, 726, 729; bionomics of, 729; measures against, 398, 729.

ochus, Telenomus.

Ocimum, experiment in infestation of, by insects and mites, 250.

Ocimum basilicum, repellent Deimestids, 250.

Ocimum sanctum, Pyrausta phoenicealis on, in Seychelles, 433.

Ocladius dianthi, sp. n., on carnations in S. Africa, 106.

Ocnerostoma piniariella (copiosella), on Pinus cembra in Austria, 664. ocreatus, Pachynematus (see Pikonema dimmocki).

octo, Amyna.

octocinctus, Stenaraeoides.

octosignatus, Brumus.

oculiperda, Thomasiniana.

Odites microbolista, sp. n., on coffee in Uganda, 151.

Odoiporus longicollis, on banana in Ceylon, 360.

Odontotermes (see Termes).

Odynerus seychellensis, predacious on Lepidoptera in Seychelles, 433. Oebia undalis (see Hellula).

Oecanthus, on tobacco in U.S.A, 393. Oechalia consocialis, predacious on Phalaenoides glycine in New South

Wales, 111.

Oecophylla smaragdina, question of value of, against Promecotheca papuana in New Guinea Territory, 435, 495.

Oedaleonotus enigma, in California, 19.

Oedaleus australis, distribution of, in Australia, 584.

Oemona hirta, bionomics and control of, in New Zealand, 266.

oestracea, Trichopsidea.

Ohio, cereal pests in, 129, 130, 505, 580, 729, 730; spiders and insects associated with maize in, 590; pests of forest and shade trees in, 399, 543, 580; probable vector of Dutch elm disease in, 325; thrips on gladiolus in, 398; orchard pests in, 61, 62, 399, 401, 735; Ancylis comptana on strawberry in, 399, 735; Lamellicorns damaging turf in, 398, 729; vegetable pests in, 13, 327, 399, 400, 651; Erythroneura comes on vines in, 323; beneficial insects and biological control in, 13, 369, 399, 649, 735.

Oil, Asphaltic, soil treatment with,

against Diaprepes, 403.

Oil, Castor, emulsions of, against Lepdiosaphes beckii, 82; in banding mixtures, 93, 202, 480; (sulphonated), as a spreader and adhesive for sprays, 274, 553, 750.

Oil, Cottonseed, emulsions of, against insects, **82, 245, 246;** factors affecting retention of

sprayed, 371.

Oil, Ground-nut, tests of emulsions containing, against insects, 167, 739.

Oil, Linseed, toxicity of emulsions of, to Lepidosaphes gloveri, 702; as an adhesive for lead arsenate, 151; unsatisfactory in baits for Cydia, 135.

Oil, Olive, surface tension and toxicity to *Lepidosaphes* of emulsions of, **702**; increasing toxicity of derris, **739**; in bait for locusts, **699**.

Oil, Pine, injected into timber against ants, 219; use of, against Cydia pomonella, 442; as an activator for pyrethrum, 550.

Oil, Soy-bean, as an adhesive for arsenical sprays, 125; forms of nicotine bentonite containing, 555, 652, 732, 733.

Oil, Sunflower Seed, calcium cyanide used with, as a soil fumigant, **431.** Oil, Tea-seed, increasing toxicity of

derris, 739.

Oil Emulsions, against Aphids, 16, 62, 210, 272, 290, 472, 707, 716, 738; against Coccids, 18, 25, 26, 59, 82, 125, 172, 185, 187, 220, 260, 270, 272, 273, 307, 323, 350, 456, 480, 481, 484, 501, 508, 549, 562, 575, 587, 637, 663, 690, 701, 702, 707, 712, 724, 728, 738, 747; against Jassids, 282, 558, 559; effect of, on natural enemies

of Typhlocyba pomaria, 558, 559; against Psylla mali, 353, 482 against other Rhynchota, 260, 269, 574, 690; against Coleoptera, 116, 308; against *Cydia* spp., 62, 74, 79, 149, 150, 168, 169, 174, 175, 254, 259, 272, 401, 442, 448, 482, 510, 544, 554, 555, 556, 588, 614, 652, 653, 733; against other Lepidoptera, 9, 16, 22, 45, 57, 108, 116, 138, 167, 185, 270, 299, 562, 574, 582, 647, 736, 743; against Diptera, 131, 298, 335, 537, 547; soil treatment with, against *Tipula aino*, 53; against mites, 16, 26, 46, 191, 297, 298, 334, 465, 466, 472, 488, 508, 648, 665, 690, 709, 712, 724, 738; against sawflies, 191, 708; and Paris green, methods of using, against termites, 187; against thrips, 29, 191, 192, 222, 291, 484, 711; tests of ovicidal action of, 16, 353; characteristics affecting insecticidal efficiency of, 701; and emulsion-suspensions, factors affecting retention and of, **280**, **370**, **371**; deposits method for determination of deposits of, on foliage, 227; types of oils for, 9, 16, 18, 25, 59, 62, 74, 79, 82, 85, 100, 108, 125, 131, 138, 149, 167, 168, 169, 172, 175, 185, 191, 210, 227, 254, 259, 260, 269, 270, 272, 297, 298, 307, 308, 323, 334, 335, 350, 355, 371, 394, 401, 442, 448, 465, 466, 472, 480, 481, 482, 484, 488, 496, 501, 508, 510, 546, 554, 555, 556, 558, 559, 562, 574, 582, 614, 637, 647, 648, 652, 653, 663, 665, 701, 702, 708, 712, 716, 733, 736, 737, 738, 739, 743, 750; formulae for, 18, 29, 46, 59, 82, 85, 125, 131, 168, 185, 187, 260, 270, 272, 273, 298, 308, 334, 335, 350, 466, 480, 481, 484, 501, 537, 546, 574, 637, 701, 702, 708; formulae containing, 16, 25, 74, 131, 138, 149, 167, 16, 23, 14, 151, 138, 145, 161, 169, 172, 174, 175, 191, 210, 259, 260, 270, 272, 335, 394, 442, 466, 481, 484, 510, 537, 554, 555, 556, 559, 562, 614, 647, 652, 653, 690, 708, 728, 738, 738, 738, 738, 739, 743; and anabasine, 739; and Bordeaux mixture, 210, 298, 537, 728; and other copper compounds, 371, 554; and cubé, 708; and cresol, 16, 738; and dinitro-cresol, 62, 549; dinitro-cyclohexylphenol, 25, 270, 272, 562; and derris, 191, 192,

510, 574, 690, 708, 739; and fluorine compounds, 62, 174, 401, 488; and lead arsenate, 46, 169, 210, 254, 259, 270, 318, 402, 442, 488, 554, 555, 588, 614, 652, 708, 750; inverted spray mixtures of arsenates or cryolite and, 174, 175, 488, 544, 652; and lime, 554, 653; and lime-sulphur, 272, 708; and nicotine, 9, 16, 29, 45, 62, 79, 131, 138, 149, 150, 185, 210, 254, 260, 270, 272, 291, 335, 442, 448, 484, 508, 510, 546, 549, 554, 555, 556, 559, 562, 647, 652, 708, 711, 733, 739, 743; and paraffin wax emulsion, 18; as a carrier for paradichlorobenzene against borers, 125, 271, 272; and pyrethrum, 53, 74, 282, 508, 736; and Selocide, 690; and sulphur, 26, 222; and tar distillates, 25, 272, 298, 343, 466, 482, 562, 738; and thiocyanates, 16, 172, 272; and zinc sulphate, 273, 298, 355, 472, 481, 482, 484, 556, 614, 653, 663, 701, 708, 711, 712, 733, 737, 738; preventing scorching of tomato by Bordeaux mixture in mixed sprays, 210; effect of, on removal of spray residues from fruit, 175, 318, 402, 614, 750; standardisation of, in Switzerland, 156; general papers on, 496; tar distillates compared with, 16, 62, 270, 297, 466, 482, 562, 663; other insecticides compared with, 16, 62, 79, 108, 138, 149, 150, 191, 272, 401, 456, 537, 647, 724, 736, 738.

Oil Palm (Elaeis guineënsis), Oryctes rhinoceros on, in Malaya, 434. Oils, in baits, 270, 398; use of,

against Coccid in cranberry bogs, 172; treatment of maize ears with, against *Heliothis*, 745; warehouses sprayed with, against rice pests, 218; atomised sprays of insecticides in, 134, 368, 400, 404, 464; aeroplanes applying sprays of, 19, 134, 748; dusts impregnated with, 125, 357, 426, 478, 544, 653, 734; and betanaphthol, bands treated with, against Cyida pomonella, 108, 168, 479, 480, 507; and naphthalene against bark-beetles, 394;

and orthodichlorobenzene against termites, 748.

Oils, Sulphonated (see Ultrawet and Oil, Castor).

oishii, Caliroa (Eriocampoides).

Oklahoma, Anthonomus grandis in, 274; Blissus leucopterus in, 63, 144; Chrysobothris femorata in, 136; grasshoppers in, 138, 139, 246, 398, 580; beneficial insects in, 137, 139.

oleae, Dacus; Filippia; Liothrips (Phloeothrips); Parlatoria; Sais-

setia.

oleellus, Prays.

Oleic Acid, types of, in baits for Cydia, 135; "soap-stock" containing, 246; wetting and spreading properties of mixtures of sodium hydroxide with, 321; effect of, on inversion of spray mixtures, 174. (See Ammonium, Triethanolamine Sodium and Oleates.)

Oleine, soap for derris spray prepared from, 615.

oleiperda, Hylesinus. oleisuga, Clinodiplosis.

oleivorus, Phyllocoptruta (Eriophyes, Phyllocoptes).

oleracea, Haltica; Polia.

oleraceum, Eurydema.

Olethreutes abietana (see Argyroploce).

Olethreutes schistaceana Eucosma).

Olfactometers, 1, 137.

Oliarus, on rice in Tanganyika, 113. Oligonychus ulmi, auct. (see Paratetranychus pilosus).

Oligotrophus bergenstammi

Apiomyia).

Olive, pests of, in Cyprus, 108; Dacus oleae on, in Greece, 71 pests of, in Italy, 161, 261, 350; pests of, in Libya, 260, 261; D. oleae on, in Malta, 469; Saissetia oleae on, in Peru, 81; pests of, in Spain, 699, 700; book on pests of, in U.S.A., 722.

Olive Fruit-fly (see Dacus oleae). Olive Oil (see Oil, Olive).

Olive Scale (see Saissetia oleae).

olivina, Euphyllura.

olynthus, Telenomus.

Omophlus, relation of, to type of soil in Russia, 72, 356.

Omophlus lepturoides, on rubber-producing plants in Ukraine, 72. Omophlus proteus, on rubber-produc-

ing plants in Ukraine, 72.

Omphale varipes, parasite of Contarinia medicaginis in Sweden, 232.

Oncideres, on avocado in Peru, 82. Oncideres cingulatus, on pecan in Florida, 587.

Oncideres dejeani, on Casuarina in Brazil, 145.

Oncideres poecila, on avocado in Peru, 82.

Oncopeltus fasciatus, tests of insecticides on, in U.S.A., 739.

Oncometopia lateralis, on cotton in Texas, 278.

Oncometopia undata, on cotton in Texas, 278.

Oncopera rufobrunnea, introduced toad of little value against, in Australia, 449.

Oncophanes atriceps, parasite of Sparganothis sulfureana in New Jersey, 563.

Onebala hibisci, bionomics of, on Hibiscus sabdariffa in Nether-

lands Indies, 358.

Onion, Thrips tabaci on, in Bermuda, 626; Hylemyia antiqua on, in Denmark and Britain, 49, 298; Lilioceris merdigera on, in Finland, 201; pests of, in Germany, 39; T. tabaci on, in Hawaii, 143, 752; T. tabaci on, in Japan, 143; H. antiqua on, in Ontario, 536; pests of, in Tanganyika, 113, 642; pests of, in U.S.A., 285, 326, 399; effect on Euxoa segetum of feeding on, 424; Maladera holosericea not experimentally feeding on, 194; insecticides causing injury to, 298.

Onion Fly (see Hylemyia antiqua).

onobrychis, Macrosiphum.

Ontario, Diprion spp. on conifers in, 487, 538; Pyrausta nubilalis on maize in, 486, 536; pests of peach in, 509, 535, 734; vegetable pests in, 536, 540, 541; Tyroglyphus farinae infesting cheese in, 539; beneficial insects and biological control in, 509, 510, 535, 537, 538.

onusta, Chariessa pilosa.

Ooencyrtus pyrillae, parasite of Pyrilla in India, 243.

Ootetrastichus, parasite of Megamelus proserpina in Philippines, 213.

Ootheca bennigseni, on Sesamum in, 642.

opaca, Eleodes.

opacum, Inostemma.

opacus, Microcryptus (Pezoporus). opalescens, Aegeria (Synanthedon).

Opatrum sabulosum, in Russia, 72, 356, 429, 432; in forest nurseries, 432; on rubber-producing plants, 72; relation of, to type of soil, 72, 356, 429.

operculella, Phthorimaea (Gnorimos-

chema).

Operophtera brumata, in Czechoslovakia, 657; bionomics and races of, in Germany, 52, 420, 716; on oak in Poland, 256; on fruit trees, 52, 420, 657; banding against, 52, 256, 420; tests of insecticides on, 297, 616; metabolism in, 440.

Ophideres (see Othreis).

Ophiomyia (see Agromyza).

Ophion, parasite of sugar-cane borers in Madagascar, 67.

Ophonus pubescens (see Harpalus rufipes).

Opilo domesticus, bionomics of, in Germany, 662.

Opius, parasite of Trypetids in Africa, 492; parasite of Oscinis theae in India, 149.

Opius fijiensis, parasite of Dacus in Fiji, **59.**

Opius fletcheri, parasite of Dacus spp. in Ceylon, **359**; proposed introduction of, into Hawaii, **359**. Opius fullawayi (see Diachasma).

Opius giffardi, parasite of Ceratitis spp. in Africa, 492.

Opius humilis, parasite of Ceratitis spp. in Kenya, 492; parasitism of C. capitata by, in Hawaii, 84.

Opius longicaudatus (see Biosteres). Opius perproximus, parasite of Trypetids in Africa, 492; introduced into Hawaii, 492.

Opius perproximus modestor, parasite of Ceratitis nigra in Kenya, 492.

Opius rhagoleticolus, parasite of Rhagoletis cerasi in Switzerland, 156.

Opius striativentris, parasite of Phytomyza ilicicola in Maryland, 541.

Opius tryoni (see Diachasma).

Opogona chlorophanes, on cassava in Tanganyika, 112.

oppositus, Leptoglossus.

Opuntia, breeding of Cactoblastis cactorum against, in S. Africa, 202; new Eriophyid on, in California, 650; thrips on, in Japan, 673.

Orange, Coccid on, in S. Africa, 377; mites possibly transmitting disease of, in Argentina, 581;

Ceratitis capitata on, in Bermuda. 626; thrips on, in Caucasus. 354; fruit-piercing moths associated with fungi on, in Dominica, 751; Dasynus antennatus on, in India, 55; C. capitata on, in Libya, 261; pests of, in Palestine, 406, 407, 636; Cossus cossus on, in Sicily, 93; pests of, in U.S.A., 508, 650; (fruit) uses of, in baits, 139, 699.

Orange, Mandarin or Tangerine, termite attacking, in California, 219; susceptibility of, to Ceratitis capitata in Palestine, 407: Cossus cossus on, in Sicily, 93; Aspidiotus perniciosus on, in Transcaucasia, 305; effects of

oil emulsions on, 355.

Orange, Satsuma, mite on, Alabama, 690.

Orange Bug, Bronze (see Rhoecocoris sulciventris).

Orchid Weevil (see Diorymerellus laevimargo).

orchidii, Anaphothrips.

Orchidophilus aterrimus, occurring and intercepted on orchids in Hawaii, 692.

Orchids, new Lamiid on, in Netherlands Indies, 262; other pests of, 550, 673, 692; methyl bromide as a fumigant for, 389.

Orcus chalybeus, failure to establish, against Eriococcus coriaceus in

New Zealand, 753.

ordinata, Sybra. Oregma lanigera (on sugar-cane), in China, 243; in Formosa, 263, 769; in Loochoo Islands, 437; value of Coccinellid predacious on, and its utilisation, 769; varieties of sugar-cane in relation to, 263.

Oregon, Epochra canadensis 60; miscellaneous pests in, 291, 398; pests of peas in, 291, 327, 653; new Aphid on spruce in, 369; beneficial insects and biological control in, 291, 398, 589, 590, 654; termite in timber imported into Japan from, 55.

Oregon Pine (see Pseudotsuga taxi-

folia).

Oreta carnea, new parasite of, on Uncaria gambir in Sumatra, 688. Orgilus longiceps, liberation of,

against Cydia molesta in Connecticut, 724.

Orgilus obscurator, establishment of, against Rhyacionia buoliana in U.S.A., 745.

Orgyia anartoides, bionomics and control of, in New South Wales, 56.

Orgyia antiqua, fungus infesting, in Italy, 152; on cranberry in Washington, 172; tests of ovicides on, 297.

Orgyia postica, on castor in Ceylon, 359; new parasite of, in Formosa,

119.

Oria musculosa, factors affecting infestation of cereals by, Russia, 428.

Oriental Peach Moth (see Cydia

molesta).

Oriental Spruce (see Picea crientalis). orientalis, Anomala; Anychus: Aonidiella (Aspidiotus); Aserica (Serica); Crioceris; Dorylus;

Orius insidiosus, predacious on Bregmatothrips iridis in U.S.A.,

273.

ornatipennis, Melcha.

ornatum, Eurydema.

ornatus, Glycyphagus; Myzus.

Ornix (see Parornix).

Orphiloides ovivorus, bionomics of, in Japan, 514; in Formosa, 514. Orphinus aethiops, in tobacco seed in Java, 434.

Orphulella speciosa, destroying blue-

grass in Iowa, 23.

Orthacris filiformis, on Citrus in Ceylon, **359.**

Orthezia insignis, tests of insecticides. on, in Tanganyika, 35, 36.

praelonga, Anthomyzid Orthezia attacking, in Brazil, 319; on Citrus etc. in Grenada, 752.

Ortho Nicotine, 733.

Orthodichlorobenzene, injected into timber against ants, 219; and oils, soil treatment with, against termites, 748; and paradichlorobenzene, against wireworms, 630; effect of, on beet, 631.

orthogonia, Porosagrotis (Agrotis). Orthol K, 554, 653, 733, 736.

Orthoptera, book on biology of, 629. Orthorrhinus cylindrirostris, nomics of, in New South Wales, 335.

Orthotomicus (see Ips).

Orthotylus marginalis, oviposition of, on currant in Germany, 234.

Orvus W. A. Paste, 395.

Oryctes rhinoceros, on coconut in Ceylon and Cochin, 360, 434, 452; relation of compost to, on oil palms in Malaya, 434; on Pritchardia in Samoa, 702; compost-pit temperatures affecting control of, by Metarrhizium anisopliae, 434.

oryzae, Calandra (Sitophilus); Lema; Podothrips (Melampodo-

thrips).

Oryzaephilus surinamensis, in copra and stored rice in Malaya, 31; in cereal products in U.S.A., 322, 369, 389; parasite of, 369; susceptibility of, to cold, 322; tests of fumigants against, 389, 741; a suitable insect for laboratory testing of insecticides, 610.

Oryzaephilus surinamenisis mercator,

610.

Oscinella frit, on Lolium perenne in Britain, 382; growth physiology affecting resistance of oats to, in Germany, 251; food-plants of, in Russian Union, 427, 617; methods of cultivation in relation to, 427.

Oscinis theae, parasites of, on tea in

India, **149.**

oslari, Hemerocampa. ostreaeformis, Aspidiotus.

Othreis fullonia (fullonica), measures against, damaging tomatos in Madras, 409; damaging papayas in Queensland, 246.

Othreis materna, measures against, damaging tomatos in Madras,

409.

Otiorrhynchus clavipes, on roses in Britain, 297.

Otiorrhynchus hirticornis food-plants of, in Germany, 344.

Otiorrhynchus ligneus, on chrysanthemums in Britain, 297.

Otiorrhynchus ligustici, bionomics and control of, in Germany, 520; importance of, in New York, 136.

Otiorrhynchus ovatus, in U.S.A., 15, 45, 270, 392; bionomics of, on nursery conifers, 392; on strawberry, 45, 270; measures against, 45, 270, 393.

Otiorrhynchus rugosostriatus, on strawberry in Britain, 383; Carabids predacious on, 384.

Otiorrhynchus singularis, in fruittree bands in Britain, 689.

Otiorrhynchus sulcatus, food-plants of, in Britain, 350, 383; bionomics of, on strawberry in Tasmania, 209; on cranberry in Washington, 172; Carabids predacious on, 384; measures against, 172, 209.

Otiorrhynchus tenebricosus, auct. (see O. clavipes).

otis, Zizera (Zizina). oudemansi, Eberhardia.

Ougenia dalburgioides, tests of insecticidal properties of, in Mysore, 360, 361.

ovata, Brachymeria.

ovatus, Aleuroglyphus (Tyroglyphus); Otiorrhynchus (Brachyrrhinus).

ovivorus, Orphiloides. oweni, Cylindroiulus.

Oxalis violacea, Tetranychina on, in Formosa, 674.

Oxya, warmth accelerating development of hibernating, 54.

Oxya chinensis, measures against, on rice in China, 498.

Oxya vicina, on Juncus in Japan, 672.

Oxycarenus hyalinipennis (on cotton), in Belgian Congo, 315; in Sudan, 512.

Oxycarenus rufiventris, on cotton in Sudan, 512.

Oxycephala longissima (longipennis) (see Brontispa).

Oxyplax ochracea, bionomics of, in Ceylon, 639.

Oxytelus sculptus, measures against, on cucumber in Holland, 299.

Oxythyrea funesta, on flowers and vegetables in Switzerland, 689.

Ozocerite, not preventing scorching of apple bark by oil and betanapthol, 480.

P.

pabulinus, Lygus.

pabulorum, Muscina. Pachydissus, on Xylia dolabriformis

in Burma, 339.

Pachynematus alaskensis (see Pikonema).

Pachynematus ocreatus (see Pikonema dimmocki).

Pachynematus piceae (see Pikonema alaskensis).

Pachyneuron aphidis, parasite of Brevicoryne brassicae in Ukraine, 303.

Pachyneuron siphonophorae, parasite of Aphids in Porto Rico, 39.

Pachyrhina maculata (see Nephrotoma).

Pachytychius strumarius, bionomics of, on peas etc. in Morocco, 641. pacificus, Aenasius; Germalus;

Tetranychus.
packardi, Melanoplus.

padellus, Hyponomeuta. padi, Anuraphis. paenerivalis, Angitia.

pagana, Haltica.

Pagoda Tree (see Sophora japonica). Palaearctic Region, key to species of Inostemma in, 526.

Palau Islands (ee Japanese Man-

dated Islands).

Paleacrita vernata, on apple and elm in Indiana, 126.

Pales pavida, parasite of Polia

oleracea in Italy, 528.

Palestine, pests of Citrus in, 77, 379, 405, 406, 514, 536, 537, 538; orchard pests in, 500, 514; Prodenia litura in, 638; Schis-tocerca gregaria in, 486; Empoasca benedettoi and disease of potato and egg-plant in, 77; Icerya purchasi on Citrus imported in Cyprus from, 108.

pallens, Geocoris.

palliatus, Hylastes (Hylurgops); Tanymecus.

pallidifrons, Erythroneura.

pallidipennis, Anomala (Phyllopertha).

pallidipes, Macrocentrus abdomin-

pallidus, Chlorodryinus; ades; Tarsonemus. pallipes, Diabrotica. Creonti-

palmaris, Nemeritis.

palmarum, Rhynchophorus.

Palms, Homaledra sabalella on, in Florida, 22; Pseudococcus brevipes on, in Formosa, 263; Acanthopsyche spp. on, in Indo-China, 184; Hispids on, in Japanese Mandated Islands, 336; new Capsid on, in Malaya, 198; Promecotheca papuana on, in New Guinea Territory, 435, 495; pests of, in Samoa, 702; fumigation of, with methyl bromide, 388. (See Coconut, Date and Oil Palm.)

Palomena prasina (on cereals), bionomics of, in Germany, 51, 52; in Switzerland, 91.

Palomena viridissima, on cereals and potato in Switzerland, 92.

Palpifer sexnotatus var. ronin, on Colocasia in Formosa, 517.

palpigera, Brachyacma. paludosa, Tipula.

Panama, Aenasius spp. parasitic on mealybugs in, 212; new weevil 405; summary of plant quarantines in, 134.

pancratii, Brithys.

Pandemis (see Tortrix). pandora, Coloradia.

panicea, Sitodrepa.

Panicum crusgalli, Pyrausta nubilalis on, in Poland, 622.

Panicum miliaceum, Pyrausta nubilalis on, in Poland, 622, 623.

Panicum prionitis, unfavourable to Schistocerca paranensis in Argentina, 147.

Panicum setigerum, Spodoptera mauritia on, in India, 241.

panicus, Thrips.

Panolis flammea (Pine Noctuid), measures against, in Germany, 235; factors affecting distribution and abundance of, in Poland, 4; ant attacking, 235; bacterial and fungus diseases of, 235, 420; samples for forecasting abundance of, **196**.

Pantomorus godmani (see Asynonychus).

Pantomorus inimicus, sp. n., on apple in Brazil, 612.

Pantophthalmus gigas, in forests in Brazil, 502.

Pantophthalmus pictus, bionomics of, on Casuarina in Brazil, 145.

Pantophthalmus vittatus, in forests in Brazil, 502.

papaveris, Aphis.

Papaya, whitefly on, in Florida, 727; pests of, in Queensland, 246; Dacus cucurbitae on, in Philippines and Hawaii, 181, 182, 325; pests of, in W. Indies, 171, 288, 491; new Jassid associated with bunchy-top of, 288.

Papaya Whitefly (see Trialeurodes

variabilis).

papayae, Empoasca.

Papilio, on Citrus in Jamaica, 490. Papilio aegeus, bionomics and control of, on Citrus etc. in New South Wales, 56.

Papilio anactus, bionomics and control of, on Citrus etc. in New South Wales, 56.

Papilio demodocus, bionomics and control of, on Citrus in Madagascar, 167.

Papilio glaucus f. turnus, Trichogramma parasitic on, in U.S.A., 744.

Papilio machaon, on carrot and Cryptotaenia japonica in Japan, 673.

Papua, summary of plant quarantines in, 692.

papuana, Promecotheca.

Parabillaea rhynchophorae, gen. et sp. n., parasite of Rhynchophorus palmarum in Argentina, 689.

of. Paradichlorobenzene, against borers, 125, 260, 262, 271, 272, 295, 591; against ants in timber, 219; against pests of stored grain, 240, 409; against pests in soil, 236, 393, 403, 436, 461, 526, 630; regulations for treatment with, against Popillia japonica in U.S.A., 384; uses of, against termites, 113, 223, 449; against woodlice in mushroom houses, 328; as a repellent, 400, other fumigants mixed 721; other fumigants mixed with, 236, 630; solvents for, 125, 219, 262, 271, 272, 295, 449, 591; causing injury to plants, 550, 631. paradoxa, Cedria.

Paraisin, active principle of Melia

azedarach, 180.

Paralipsa gularis (see Aphomia). Paranagrus perforator, parasite of Megamelus proserpina in Philip-

pines, 213.

paranensis, Schistocerca. Paraneotermes, subgen. n. for Calotermes simplicicornis, 219.

Paraplesius ugandanus, sp. n., on tea in Uganda, 612.

Parasa lepida, parasites of, on tea in Ceylon, 639; on coconut in Cochin, 452.

Parasites, physiological causes of immunity from, 287.

Parasthenaridea arecae, gen. et sp. n., on Areca catechu in Malaya, 198.

Paratenodera sinensis, Dermestid parasitic on eggs of, in Japan, 514.

Paratetranychus, measures against, on satsuma orange in Alabama,

Paratetranychus citri, measures against, on Citrus in U.S.A., 131, 283, 286, 508; method population studies of, 283.

Paratetranychus pilosus (on fruit trees), in Britain, 191, 297, 466, 709; in Germany, 100; bionomics of, in Holland, 464, 465; in Japan, 53, 604; in New Zealand, 46; in U.S.A., 15, 16, 26, 488, 646, 724, 738, 748; predacious enemies of, and treatments destroying them, 15, 16, 100, 465, 724; measures against, 15, 16, 26, 46, 191, 297, 465, 466, 488, 646, 709, 724, 738; laboratory tests of ovicides against, 297; Metatetranychus ulmi considered correct name for, 465.

stenoperitrematus, Paratetranvchus sp. n., food-plants of, in Central Asia, 530.

Paratheresia (see Theresia).

Paratrioza cockerelli, causing Psyllid yellows on potato in U.S.A., 281, 578, 643; physiology of Psyllid yellows caused by, 281; winter food-plant of, 578; measures against, 579, 643.

pardalina, Locustana; Myiopardalis.

Parectopa robiniella, on Robinia pseudacacia in Virginia, 549. parenthesis, Hippodamia.

Paria canella, measures against, on Corylus in Connecticut, **547**. pariana, Anthophila (Hemerophila).

Paris Green, in baits, 40, 139, 211, 282, 441, 540, 566, 574, 698, 705, 727; dusting with, 142, 211, 222, 226, 277, 279, 433, 638, 652; carriers for, 211, 222, 226, 277, 652; spraying with, 65, 99, 274, 285, 291, 352, 399, 431, 433, 478, 484, 489, 552, 553, 643, 652, 664; adhesives and spreaders for, 285, 399, 431, 433, 552, 553, 652 and Bordeaux mixture, 99, 285, 522, 664; and calcium arsenate, 142, 279; and hydrol, 65; and lime, 291, 431, 433, 484; and sugar, 274, 399, 489; maize seed treated with, against Carabid, 362; uses of, against termites, 113, 187, 338, 512; tests of toxicity of, to *Hippodamia convergens*, 390; causing injury to plants, 274, 338, 399, 484, 489, 552, 553; estimation of arsenic content of, 772; composition and physical properties of commercial samples of, 183, 282; homologues of, 30, 284; Soviet green similar to, 479; insecticides compared with, 30, 279, 552.

Parkia africana, not containing rotenone, 182.

Parlatoria blanchardi (Date Scale), eradication of, in California, 18; in Sudan, 513.

Parlatoria oleae, measures against, on apple in Cyprus, 260.

Parlatoria pergandei camelliae, new parasite of, in Haiti, 41.

Parlatoria zizyphus (on Citrus), in Formosa, 263; measures against, in Indo-China, 185.

Parnara borbonica, on rice in Madagascar, 118.

Parnara guttata, bionomics and control of, on rice in China, 244. 498.

Parnara mathias (on rice), Ceylon, 358; bionomics of, in China, 244.

Parnara pouteri, on rice in Madagascar, 118. paroditis, Pyroderces.

Paropsis dilatata, on Eucalyptus in New Zealand and Australia, 753; parasites and biological control of, 753.

Parornix geminatella, measures against, in orchards in Michigan, 743.

Paroxyna tessellata, bionomics and control of, in Ukraine, 619.

Parsley, Listronotus latiusculus on, in U.S.A., 565.

Parsnip, Depressaria heracleana on, in Britain, 296.

parvipennis, Dasyscapus.

parvula, Epitrix.

parvulus, Longitarsus. parvus, Microcerotermes.

Paspalum urvillei, Solubea pugnax on, in Louisiana, 218.

Passiflora, Dacus cucurbitae on, in Philippines, 181.

passiflorae, Dacus (Chaetodacus).

Patanga succincta, in N. Borneo, 112; bionomics of, in India, 629. paucispinosa, Frankliniella.

paulistus, Aenasius.

Paulownia tomentosa, Phassus signifer on, in Formosa, 517.

pauxillus, Rhynchites.

pavida, Pales.

Pavonia kraussiana, Dysdercus on, in Uganda, 373.

pavonia, Eudia (Saturnia).

Pea, Heliothis armigera on, in S. Africa, 364; pests of, in Australia, 582, 754; thrips on, in Britain, 527; little damaged by Bruchids in Burma, 7; pests of, in Canada, 60, 293, 623; Tychius quinquepunctatus on, in Czechoslovakia, 657; pests of, in Japan, 208, 265, 605, 669; Pachytychius strumarius on, in Morocco, 641; pests of, in Russia, 75, 355, 428, 432; pests of, in U.S.A., 19, 21, 26, 65, 128, 129, 270, 285, 291, 327, 387, 443, 489, 545, 651, 653, 740; Myzus persicae transmitting mosaic disease of, 265; susceptibility of varieties of, to Cydia nigricana, 428.

Pea (Stored), pests of, and their control, 75, 644, 654; practically immune from grain mites, 96; methyl bromide not impairing germination of, 390.

Pea Aphis (see Macrosiphum ono-

brychis).

Pea Bruchid (see Bruchus pisorum). Pea Mite (see Penthaleus major). Pea Moth (see Cydia nigricana).

Peach, pests of, in Australia, 335, 448; Dacus ferrugineus dorsalis on, in Bonin Islands, 672; pests of, in Canada, 173, 510, 535, 734: Tettigoniid on, in China, 244; Recurvaria nanella on, in Cyprus, 108; pests of, in France, 343, 416; Anarsia lineatella on, in Germany, 718; pests of, in India, 602, 765; pests of, in Japan, 53, 54; pests of, in Italy, 163, 189, 594, 714, 758; Charagia virescens on, in New Zealand, 266; Coccid on, in Porto Rico, 329; Cydia molesta on, in Uruguay, 582; pests of, in Russian Union, 305, 530; pests of, in Switzerland, 163, 715; pests of, in U.S.A., 16, 17, 18, 19, 20, 26, 125, 126, 127, 173, 270, 271, 287, 288, 323, 385, 395, 399, 443, 444, 506, 544, 545, 551, 560, 587, 644, 647, 653, 724, 725, 726, 734, 735, 748; Macropsis trimaculata and virus diseases of, 173, 385; types of injury to, by Capsids and Pentatomids, 127, 444; Dacus cucurbitae experimentally developing in, 668; Eutettix disciguttus and Lymantria dispar not surviving on, 208, 759; insecticides causing injury to, 16, 26, 125, 285, 323; zinc sulphate as a corrective for arsenical injury to, 587.

Peach (Dried Fruit), pests of, in California, 30, 228; not attacked by Aphomia gularis in California,

Peach Aphis (see Myzus persicae). Peach Borer (see Aegeria exitiosa). Peach Leafhopper (see Macropsis trimaculata).

Peach Moth, Oriental (see Cydia

molesta).

Peach Moth, Yellow (see Dichocrocis punctiferalis).

Peach Scale, White (see Aulacaspis pentagona).

Peach Twig Borer (see Anarsia lineatella).

Peakesia, in W. Australia, 584.

Peanuts (see Ground-nuts).

Pear, Aonidiella aurantii on, in S. Africa, 455; Lachnus salignus on, in Britain, 613; pests of, in Br. Columbia, 59, 171; Coccids on, in Chile, 221; Stephanitis pyrion, in Cyprus, 260; pests of, in Czechoslovakia, 620, 657, 658; pests of, in France, 192, 259, 416, 525; pests of, in Germany, 52, 660, 761, 762; pests of, in India, 340, 765; pests of, in Italy, 594, 636, 758; pests of, in Japan, 53, 54, 604, 671; Lymantria dispar on, in Jugoslavia, 759; pests of, in Korea, 605, 671; Monosteira unicostata on, in Libya, 260; Zeuzera pyrina on, in Malta, 469; M. unicostata on, in Morocco, 340; Cydia pomonella on, in Palestine, 500; C. pomonella on, in Poland, 99; pests of, in Russian Union, 477, 481, 530; Aglaope infausta on, in Spain, 697; pests of, in U.S.A., 19, 127, 130, 288, 748; Cydia spp. on, in Victoria, 168, 169; Dacus cucurbitae experimentally developing on, 668; Eutettix disciguttus not surviving on, 208; insecticides causing injury to, 97, 285.

Pear (Fruit), injured by HCN fumigation, 307; problem of spray residues on, 169, 259, 525; (dried), not attacked by Aphomia gularis in California, 573.

Pear Blister Mite (see Eriophyes pyri). Pear Blossom Weevil (see Anthono-

mus piri).

Pear Midge (see Contarinia pyrivora). Pear Sawfly (see Caliroa limacina). Pear Thrips (see Taeniothrips inconsequens).

Peat, preparation of nicotine peat from types of, 24.

Pecan, pests of, in U.S.A., 18, 136, 138, 323, 324, 587, 690, 722, 747; zinc sulphate preventing injury to, by lead arsenate, 138.

Pecan Nut Case-bearer (see Acrobasis caryae).

Pecan Weevil (see Curculio caryae). pecki, Sesamia.

pectinata, Puliciphora.

Pectinophora gossypiella (see Platy-

pectinophorae, Chelonus.

pectoralis, Amorphoidea; Dacus (Tridacus).

pedata, Glypta.

pedestris, Cleonus (Leucosomus); Riptortus.

Pediculoides ventricosus, predacious on Platyedra gossypiella in Tanganyika, 112.

pedro, Proteides mercurius.

Pegohylemvia (see Hylemvia). bicolor, Pegomvia apparently restricted to Rumex in France, 314.

apparently Pegomyia hyoscyami, restricted to Solanaceae in France, 314.

Pegomyia hyoscyami var. betae, bionomics of, in France and

Germany, 314.

Pelargonium roseum, tests of oil of, against Aphids and mites, 249, 250; experiment in artificial infestation of, by insects and mites, 250.

pellucida, Camnula; Diaphnidia.

pellucidus, Barypithes.

pembertoni, Microbracon.

Pempheres affinis (on cotton), in Burma, 409; in India, 89, 409; new parasites of, 89.

pempheriphila, Euderus.

Pemphigus betae, acquiring virus of

beet curly-top, 445.

Pemphigus populitransversus, in U.S.A., 19, 649; probably on lettuce, 19; technique for rearing Coccinellids on frozen, 649, 650. pendula, Lonchaea (Carpolonchaea). Pendulinus antennatus (see Dasynus).

penetrans, Macroglenes.

Pencillium, associated with Corioxenos antestiae in Antestia in E. Africa, 179; infesting Citrus fruits attacked by Ceratitis in Palestine, 406.

Peniophora gigantea, insects associated with, in pine logs in Minnesota, 43.

pennipes, Trichopoda.

Pennisetum, Aphis maidis on, in Porto Rico, 40.

Pennisetum typhoideum, Peregrinus maidis and disease in, in Madras, 532; Cecidomyiid on, in Sudan, 513; attraction of, for Schistocerca gregaria, 628.

Pennsylvania, pests of greenhouse and ornamental plants in, 488, 547; pests of leguminous plants in, 287, 580; orchard pests in, 26, 135, 287, 488; Popillia japonica and Aserica castanea in, 324, 325, 552; beneficial insects and biological control in, 324, 325, 552, 580.

pennsylvanicus, Camponotus hercu-

Pentachloro - dioxy - triphenal methane-sulphonic acid, against carpet beetles, 227.

pentagona, Aulacaspis.

Pentodon idiota, relation of, to types of soil in Russia, 72, 356; to on rubber-producing plants, 72.

Penthaleus major, food-plants and control of, in Australia, 85, 582, 583; bionomics and distribution of, 582.

Pentilia castanea, introduced into Santo Domingo against Aspidiotus destructor, 10.

peponis, Plusia.

Pepper Weevil (see Anthonomus

eugenii).

Peppergrass (see Lepidium nitidum). Peppers (see Capsicum).

perdix, Rhinaria.

Coccinellid Peregrinus maidis, favoured by honeydew of, in Florida, 727; relation of, virus disease of Sorghum, etc., in India, 55, 532.

peregrinus, Trionymus. perfidus, Moropactus. perforator, Paranagrus.

perforatus, Eucalymnatus (Lecanium).

pergandei, Parlatoria.

pergandiella, Encarsia.

Perilampus, hyperparasite of Cydia pomonella in U.S.A., 30.

Perilla frutescens var. crispa, Pyrausta phoenicealis on, Japan, 771.

Perilloides (Perillus) bioculatus (predacious on Coleoptera), effect of hosts on development of, in Ohio, 13, 14; rearing of, imported into France, 595.

periparetus, Telenomus.

Periphyllus lyropictus, giant cells in larvae of, parasitised by Aphidius vibis, 464.

Periphyllus negundinis, penetration of setae of, 225.

Periphyllus testudinatus, ants associated with, in Germany, 195.

Periplaneta americana, experiments with nicotine and, 225, 439.

Perisierola angulata, liberation of, against Cydia molesta in Connecticut, 16.

Perisierola cellularis var. punctaticeps, parasite of Platyedra gossypiella in Mexico, 275.

Perisierola emigrata, parasite of Argyroploce illepida in Hawaii, 497.

Peritelus hirticornis (see Otiorrhynchus).

Peritetranychus glabrisetus, gen. et sp. n., on watermelon in Central Asia, 530.

Peritetranychus tuberculatus, sp. n., on mulberry in Central Asia, 530.

Perkinsiella saccharicida (on sugarcane), in Hawaii and Mauritius, 454; in Natal, 411; bionomics and biological control of, 454. perla, Chrysopa.

perniciosi, Prospaltella.

pernicosus, Aspidiotus (Aonidiella). pernyi, Antheraea.

Peronea minuta, on cranberry in New Jersey, 563.

Peronospora tabacina, 744.

perplexa, Scutellara. perplexus, Pilophorus.

perproximus, Opius. perpusilla, Pyrilla.

Persea gratissima (see Avocado). perseae, Meroprion (Prionopus).

Persia (Iran), Platyedra vilella on cotton in, 249; losses and expenditure due to locusts in, 685; summaries of plant quarantines in, 134, 692, 751.

Anuraphis (see persicae, schwartzi); Myzus (Myzoides).

Persimmon, insects on, in Caucasus, 305, 354; Dacus ferrugineus dorsalis not developing on, in Formosa, 263: Lymantria dispar on, in Japan, 604; book on pests of, in U.S.A., 722.

persona, Xyleutes.

personatus, Chrysomphalus (Aspidiotus).

perspicillator, Barylypa.

persuasoria, Rhyssa. pertinax, Mylabris.

perturbans, Entedon.

Peru, cotton pests in, 533, 627; miscellaneous pests in, 81, 362; beneficial insects and biological control in, 81, 627; rotenoneyielding plants in, 319; parasites

into U.S.A. from, 390, 391. Petrol, in mixtures against ants in telephone poles, 219; in mixtures for treating bands against Cydia

of Diatraea saccharalis introduced

pomonella, 108, 507. Petroleum Ether, as a solvent, 227,

412.

Pezomachus (see Gelis). Pezoporus (see Microcryptus). Phaedon cochleariae (on crucifers), measures against, in Britain, 382; effect of fertilisers on infestation by, in Latvia, 257.

Phaenobremia, predacious on Brevicoryne brassicae in Britain, 464.

Phaenocarpa livida, parasite of Geomyza tripunctata in Britain, 382.

Phaeogenes, parasite of Cydia molesta in Italy, **154.**

Phaeogenes haeussleri, liberation of, against Cydia molesta in Connecticut, 16, 724.

phaeorrhoea, Nygmia.

Phalaenoides glycine, natural enemies and control of, on vines in New South Wales, 111.

phalaenopsidis, Diaxenes.

Phalaris, cereal pests on, 290, 427. Phalera bucephala, on Tilia in Finland, 201.

phalerata, Campsomeris.

Phanerotoma philippinensis, parasite of Maruca testulalis in Java, 111.

Phanerotoma planifrons, parasite of Cydia glycinivorella in Manchuria, 452; imported into U.S.A. against Etiella zinckenella, 329.

Phanerotoma tibialis, parasite of Cydia pomonella in U.S.A., 30, 323; utilisation of, 323.

Phanomeris phyllotomae, establishment of, against Phyllotoma nemorata in U.S.A., 745.

Phanurus (see Telenomus).

phaseoli, Agromyza; Bruchus; Trifidaphis.

Phaseolus, Aphis laburni on, in Japan, 53.

Phaseolus lunatus (see Lima Beans). Phaseolus radiatus, Aserica japonica

on, in Japan, 605. *Phaseolus vulgaris* (see Beans).

Phassus, migrating from Buettneria to Gmelina arborea in Burma, 339.

Phassus signifer (formosanus), measures against, in forests in Formosa, 517; synonymy of, 517.

Phaulacridium gemini, attacking tobacco in Queensland, 494.

Phaulacridium vittatum, bionomics and distribution of, in Australia, 584.

Pheidole megacephala (fostering Aphids and Coccids), measures against, in Bermuda, 626; in Natal, 411.

Pheidole punctulata, measures against, fostering Pseudococcus kenyae in Kenya, **592.**

Pheletes, soil fumigation against, in

Washington, 326.

Pheletes californicus, crops attacked by, in U.S.A., 326, 327, 569, 747; flooding against, 569.

Pheletes canus, on onions and potatoes in U.S.A., 326, 327.

Phellinus cryptarum, Xestobium associated with, in oak timber in Britain, 707.

Phenacoccus, new parasite of, in Brazil, 212; in France, 248.

Phenacoccus gossypii, practice favouring infestation of chrysanthemum by, in U.S.A., 396.

Phenacoccus iceryoides, on mango in Ceylon, 359.

Phenacoccus saratogensis, sp. n., on Hystrix patula in New York, 226. Phenacoccus solani, on Portulaca

oleracea in Hawaii, 213.

Phenothiazine (see Thiodiphenylamine).

Philaenus spp., on narcissus in U.S.A., 288.

philippinensis, Phanerotoma.

Philippines, Artona catoxantha on coconut in, 675; Dacus cucurbitae in, 181; Megamelus proserpina on taro in, 212; beneficial insects in, 213, 675; birds destroying locusts in, 675; Orchidophilus aterrimus intercepted in Hawaii from, 692.

Philodicus femoralis, predacious on Alisssonotum impressicolle in

Burma, 294.

Phlepsius irroratus, on narcissus in U.S.A., 288.

Phlepsius strobi, acquiring virus of beet curly-top in California, 445. Phleum pratense, Meromyza ameri-

cana on, in Kansas, 122; Cirphis unipuncta on, in Nova Scotia, 441. Phloeothrips oleae (see Liothrips).

Phlosothora maticulas (ass. Putter)

Phlogophora meticulosa (see Brotolomia).

phloxiphaga, Heliothis.

Phlyctaenia rubigalis, practice favouring infestation of chrysanthemum by, in U.S.A., 396.

Phlyctaenodes bifidalis (see Loxo-

stege).
phoenicealis, Pyrausta.

Phoenix, Homaledra sabalella on, in Florida, 22.

Phoenix dactylifera (see Date Palm).

Phoracantha recurva, on Eucalyptus in New Zealand and Australia, 754.

Phorbia brassicae (Cabbage Fly), measures against, in Massachusetts, 647; in Ukraine, 159.

Phorbia cilicrura, parasitising locust eggs in Argentina, 147.

Phorbia genitalis, date of sowing wheat in relation to, in Russia, 427.

Phorbia regina, susceptibility of, to injections of nicotine, 332.

Phorocera erecta, parasite of Tortrix rosaceana in Utah, 141.

Phorodon humuli (Hop Aphis), predacious enemies and control of, in Germany, 523.

phorodontis, Aphidius.

Phosphorus, distribution and damage caused by species of, 469.

Phosphorus gabonator, bionomics of, in Br. Cameroons, 469.

Photoelectric Cell, use of, in titration of nicotine, 695.

Phragmatiphila typhae var. fraterna, measures against, on Typha latifolia in Czechoslovakia, 345.

Phragmites, Oscinella frit breeding on, in Azerbaijan, 617; Acridid on, in Formosa, 769.

Phrynosoma, destroying ants in Utah, 397.

Phryxe (see Zenillia).

Phthia picta, bionomics and control of, on tomatos in Texas, 397.

Phthorimaea, measures against, on tomato in Hawaii, 752.

Phthorimaea atriplicella, on beet in Switzerland, 689.

Phthorimaea heliopa, measures against, on tobacco in Mysore, 606.

Phthorimaea lycopersicella, bionomics and control of, in U.S.A., 217, 328, 580, 747.

Phthorimaea ocellatella, on beet in Morocco, 416.

Phthorimaea operculella, in Brazil, 145, 320; in Hawaii, 214, 752; in Malta, 469; in U.S.A., 393; on potato, 145, 214, 320, 469, 752; on tobacco, 145, 320, 393.

Phthorimaea plaesiosema, on tomato in New South Wales, 496.

Phyllaphis fagi, on beech in Britain, 296.

Phyllocnistis citrella (on Citrus), measures against, in Indo-China, 185; bionomics of, in Japanese Empire, 438. Phyllocoptes advens, sp. n., in California, 650.

Phyllocoptes ligustri, sp. n., on privet in California 650

in California, **650.**Phyllocoptes lycopersici, sp. n., **103**;
on tomato in Australia and New

Zealand, 45, 103, 334; measures against, 46, 335.

Phyllocoptes oleivorus (see Phyllocoptruta).

Phyllocopies schlechtendali, measures against, in orchards in Washington, 488.

Phyllocoptes vitis, winter measures against, on vines, **521**.

Phyllocoptruta, gen. nov., 650.

Phyllocoptruta (Phyllocoptes) oleivorus (on Citrus), in Formosa, 674; in U.S.A., 650, 690; measures against, 674, 690.

phyllocroma, Chrysopa.

Phyllognathus dionysius, on sugarcane in Bombay, 261.

Phyllopertha (see Anomala).

Phyllophaga (see Lachnosterna). phyllopus, Leptoglossus.

phylloscia, Eucosma.

Phyllotoma nemorata (on birch), in Austria, 215; in U.S.A., 215, 745; parasites and biological control of, 215, 745.

phyllotomae, Phanomeris.

Phyllotreta atra, on crucifers in Poland, 255.

Phyllotreta cruciferae, on crucifers in Poland, 255.

Phyllotreta nemorum, on crucifers in Poland, 255.

Phyllotreta undulata, on crucifers in Poland, 255.

Phyllotreta vittata, on crucifers in Japan, 605, 769; bionomics of, 769.

Phylloxera, Chrysopa perla predacious on, on oak in Italy, 189.

Phylloxera vitifoliae (vastatrix) (on vines), distribution of and legislation against, in Australia, 608; distribution of forms of, in Germany, 601; resistance of varieties of vines to, in Japan, 560; in Switzerland, 299; review of data on, 608.

Phylus coryli (with var. avellanae), eggs of, on hazel in Germany, 234.

Physcus bifasciatus, sp. n., in Japan, 296.

Physocephala, parasitic on honey bees in U.S.A., 133.

Phytalus (see Lachnosterna).

Phytodecta fornicata, bionomics of, in Jugoslavia, 760; tests of pyrethrum dust on, 197.

Phytodietus crassitarsis, parasite of Cydia molesta in Italy, 154.

phytolymus, Maleuterpes (see M. spinipes).

Phytometra (see Plusia).

Phytomonas pruni, zinc sulphate ineffective against, on peach, 588. Phytomyptera nitidiventris, parasite of Clysia ambiguella in France,

Phytomyza atricornis (nigricornis, auct.), measures against, on cineraria in Australia, 404; on chrysanthemums in Britain, 349; on crucifers and peas in Japan, 605; bionomics of, 404, 605; characters distinguishing Agromyza solani and, 710.

Phytomyza ilicicola, bionomics and control of, on Ilex spp. in U.S.A.,

541, 546.

Phytomyza lateralis, measures against, on chicory in Belgium, 313.

Phytonomus (see Hypera).

Phytonomus posticus (see Hypera variabilis).

Phytophaga destructor (see Mayetiola).

Picea (see Spruce).

Picea abies, pests of, in Czecho-slovakia, 657.

Picea canadensis, Chermes ishiharai on, in Japan, 438.

Picea engelmanni, Rhopalosiphum abietinum on, in Austria, 715. Picea glauca, Diprion polytomum

on, in Canada, 487.

Picea glehni, pests of, in Japan and Sakhalin, 438, 452.

Picea jezoensis, pests of, in Japan and Sakhalin, 438, 452, 516.

Picea mariana, Diprion polytomum on, in Canada, 487.

Picea omorika, Otiorrhynchus ovatus on, in nurseries in New York, 392. Picea orientalis, relation of Chermes

spp. to, in Germany, 3, 661.

Picea rubra, Diprion polytomum on, in Canada, 487.

Picea sitchensis, Barypithes pellucidus on, in nurseries in Britain, 625; Chermes japonicus on, in Japan, 438.

piceae, Chermes (Adelges, Dreyfusia); Pachynematus (see Pikonema alaskensis).

piceaella, Recurvaria.

piceana, Tortrix (Cacoecia).

piceicola, Cinara.

piceus, Alphitobius (see A. laevi-

gatus); Attagenus. pichtae, Cinara (Lachnus).

Picrasmin, constituent of quassin, 694.

Picroscytoides cerasiops (parasite of Lixus spp.), bionomics of, in Morocco, 417.

picta, Melanophila; Phthia.

pictus, Cirrospilus (Entedon); Pantophthalmus.

picus, Halyomorpha. piercei, Microthrips. pieria, Comocritis.

pierie, Martine pieria, Comocritis.

Pieria brassicae (on crucifers), in Britain, 342, 632; in Czechoslovakia, 656; in Italy, 712, 713; bionomics of, in Poland, 720; in Ukraine, 159; composition of food-plant affecting development of, 632; parasites of, 342; giant cells in larvae of, parasitised by Apanteles glomeratus, 464; measures against, 713, 720.

Pieris rapae (on crucifers), in Britain, 342; in Egypt, 456; in Hawaii, 214; in Japan, 605; in U.S.A., 13, 285, 399, 651; natural enemies of, 214, 342, 456, 605; warmth accelerating development of hibernating, 54; pH and buffer value of blood of, 386, 387; toxicity of arsenicals and fluorine compounds to, 388; measures against, 285, 399, 651.

Piesma quadratum (on beet), trapstrips against, in Germany, 419; crinkle disease transmitted by, 419, 496.

Piestopleura thomsoni, hyperparasite of Sitodiplosis mosellana in Sweden, 231.

Pigeon Pea (see Cajanus cajan). piger, Cleonus; Microbracon.

Pigs, destroying Naupactus leucoloma in Florida, 132.

Pikonema, gen. nov., 293.

Pikonema alaskensis, on spruce in Canada and U.S.A., 293; synonym of, 293.

Pikonema dimmocki, on spruce in Canada and U.S.A., 293; synonym of, 293.

Pikonema ruralis, in Oregon, 293. Pilophorus spp., eggs of, on currant

in Germany, 234.
pilosa, Anoxia; Chariessa; Icerya.
pilosella, Campsomeris.
pilosellus, Brachypeplus.

pilosulus, Hyposoter.

pilosus, Paratetranychus.

Pimpla albicincta, parasite of Clania minuscula in China, 245.

Pimpla alternans, parasite of Cydia molesta in Italy, 154.

Pimpla atrocoxalis, parasite of Tortrix rosaceana in Utah, 141. Pimpla attaci, parasite of Dendroli-

mus spectabilis in Japan, 669.

Pimpla disparis, hosts of, in Korea and Japan, 264, 669.

Pimpla epicnapterus, parasite of Dendrolimus spectabilis in Japan, 669.

Pimpla examinator, parasite of Lymantria dispar in Germany,

Pimpla foliae, sp. n., parasite of Phyllotoma nemorata in Austria, 215; introduction of, into U.S.A., 215.

Pimpla glycinivorellae, parasite of Cydia strobilella in Japan, 516.

Pimpla indagatrix, parasite Thyridopteryx ephemeraeformis in Illinois, 173.

Pimbla inquisitor, parasite Lymantria dispar in Jugoslavia, 759.

Pimpla nankingensis, parasite of Clania minuscula in China, 245.

Pimpla nigribasalis, sp. n., parasite of Dendrolimus spectabilis China, 119.

Pimpla pluto, parasite of Dendrolimus spectabilis in Japan, 669.

Pimpla roborator, attempted introduction of strains of, into Texas, Mexico and Porto Rico against Platyedra gossypiella, 275, 329; parasite of Pyrausta nubilalis in Europe, **275.**

Pimpla sagax, parasite of Cydia molesta in Italy, 154.

Pimpla sanguinipes, parasite of Desmia funeralis in California, 286.

pinastri, Hylobius.

Pine (Pinus), pests of, in Australia, 8, 449; Melolontha hippocastani on, in Belgium, 193; pests of, in Czechoslovakia, 657; Thaumetopoea pinivora on, in Denmark, 518; Chrysobothris chrysostigma on, in Finland, 518; pests of, in Germany, 189, 235, 312, 419; pests of, in Holland, 183, 348, Diprion sertifer on, in 621; Hungary, 197, 198; pests of, in Japan, 295, 296, 669; pests of, in Korea, 767; Hylobius abietis on, in Latvia, 90, 722; suitability of species of, for Dendrolimus spectabilis in Manchuria, 672; Sirex noctilio on, in New Zealand, 58; pests of, in Poland, 4, 523; pests of, in Russian Union, 158, 429, 430, 431, 432; pests of, in Sweden, 197, 198, 634; pests of, in U.S.A., 270, 326, 329, 394, 399, 441, 548, 580, 650, 723, 745; insecticides causing foliage injury to, 270; effects on, of soil treatment with lead arsenate, 441; (timber) Coleopterous pests of, 91, 195, 354, 726; nutritive value of, for Hylotrupes bajulus,

Pine, Cembran (see Pinus cembra). Pine, Jeffrey (see Pinus jeffreyi). Pine, Lodgepole (see Pinus contorta). Pine, Norway or Red (see Pinus resinosa).

Pine, Scots (see Pinus sylvestris). Pine, Sugar (see *Pinus lambertiana*). Pine, Weymouth (see Pinus strobus). Pine Geometrid (see Bupalus piniarius).

Pine Needle Scale (see Chionaspis pinifoliae).

Pine Noctuid (see Panolis flammea). Pine Oil (see Oil, Pine).

Pine Processionary Caterpillar (see Thaumetopoea pinivora)

Pine Shoot Moth (see Rhyacionia buoliana).

Pine Weevil (see Hylobius abietis). Pineapple, pests of, in Formosa, 263, 674; pests and disease of, in Hawaii, 143, 213, 570, 692; Pseudococcus brevipes on, in Porto Rico. 329.

Pineapple Mealybug (see Pseudococcus brevipes).

pineti, Solenobia. Pineus (see Chermes).

pinguis, Agromyza (Ophiomyia); Leucopholis (Lepidiota).

pini, Chermes (Pineus); Diprion (Lophyrus); Ips; Pissodes.

piniariella, Ocnerostoma. piniarius, Bupalus.

pinicola, Fumaria (Fumea).

pinifoliae, Chionaspis.

Myelophilus piniperda, (Blastophagus).

piniphilus, Gnophothrips; Pissodes.

pinivora, Thaumetopoea. pinnaeformis, auct., Lepidosaphes

(see L. beckii). aspidistrae, Pinnaspis measures against, on ferns in Britain, 712. Pinnaspis buxi, on coconut in

Seychelles, 433.

Pinus (see Pine).

Pinus cembra, pests of, in Austria, 663, 664; Chermes cembrae on, in Germany, 762.

Pinus contorta, pests of, in U.S.A., 394, 580.

Pinus himekomatsu, new Aphid on, in Japan, 452.

Pinus jeffreyi, pests of, in U.S.A., 226, 394, 441.

Pinus lambertiana, Dendroctonus monticolae on, in California, 394.

Pinus mugo, Rhyacionia turionana var. mughiana on, in Austria, 663.

Pinus patula, Fumaria pinicola on, in Transvaal, 68.

Pinus ponderosa, pests of, in U.S.A., 369, 394, 441, 580.

Pinus resinosa, pests of, in U.S.A., 43, 386, 547, 549; insects associated with fungi in, 43, 386; insecticides causing injury to, 549.

Pinus strobus, Barypithes pellucidus on, in Britain, 625; Chermes strobi on, in Germany, 189.

Pinus sylvestris, pests of, in Czechoslovakia, 657; bark-beetles on, in Estonia, 462; Diprion frutetorum on, in Ontario, 538; Lamellicorn damaging, in Ukraine, 158; pests of, in U.S.A., 18, 549.

Piophila casei, bionomics of, in meat factories, 38.

Pipiza, predacious on Chermes nordmannianae in Germany, 661.

Pipunculus, parasite of Nephotettix in Japan, 439; parasite of Eutettix tenellus in Utah, 127.

Pirene, parasite of Sitodiplosis mosellana in Japan, 670.

Pirene chalybea, bionomics of, in Sweden, 231.

piri, Anthonomus.

piricola, Epidiaspis (see E. leperii); Toxoptera.

pisi, Macrosiphum (Illinoia) (see M. onobrychis); Polia (Mamestra).

pisivorus, Kakothrips.

Pisonia subcordata, Orthezia praelonga on, in Grenada, 752.

pisorum, Bruchus.

Pissodes pini, measures against, on pine in Sweden, 634, 635.

Pissodes piniphilus (on pine), in Estonia, 462; factors favouring outbreaks of, in Sweden, 634.

Pistachio, Hylesinus vestitus on, in Sicily, 119.

Pithecolobium lobatum, Xystrocera festiva on, in Java, 261.

Pityogenes (see Ips).

Pityogenes micrographus (see Pityophthorus).

Pityophthorus micrographus, measures against, on spruce in Sweden, 634.

plaesiosema, Phthorimaea.

Placsius javanus, introduced into Porto Rico from Fiji, 212.

plagiata, Anaitis; Tiracola.

Plagiognathus arbustorum, on currant in Germany, 234.

Plagiolepis longipes (see Anoplolepis).

plagiophleps, Pteroma.

Plane, Stomacoccus platani on, in California, 18.

planifrons, Phanerotoma.

planirostris, Rhynchites (see R. nanus).

Planispa, gen. nov., 336.

Planispa castaneipennis, sp. n., on coconut and other palms in Saipan, 336.

Planispa chalybeipennis, on coconut and other palms in Caroline Islands, **336.**

Plant Diseases, symposium on the relations of insects to, **385**, **386**. (See Bacteria, Fungi and Virus Diseases.)

Plant Pest Legislation, in S. Africa, 66, 691; in Australia, 267, 608; historical review of, in Canada, 23; in Egypt, 385, 682, 692; in Germany, 91, 229, 719, 751; in Br. Guiana, 134, 491; in S. Rhodesia, 134, 366; in U.S.A., 133, 134, 228, 229, 384, 385, 722, 727, 750, 751; in Zanzibar, 111; summaries of, in other countries, 107, 120, 134, 201, 229, 385, 491, 691, 692, 751.

Plant Protection, text-book on, 703; bibliography of literature of, in 1936, 248.

plantaginis, Aphis.

Plantago spp. (Plantains), Dociostaurus maroccanus feeding on, in Italy, 162; relation of pests to, in U.S.A., 323, 565.

Plants, bibliography on pests of tropical, **560**.

platani, Stomacoccus.

platensis, Aphidius.

Platyedra gossypiella (Pink Cotton Bollworm), in Brazil, 504, 689, 704; in China, 498; in Belgian Congo, 315; in Cyprus, 108; in Formosa, 309; in India, 408,

532; in Java, 766; in Mexico, 274, 275; not occurring in Nigeria, 667; in Sudan, 511; in Tanganyika, 112, 642; in Uganda, 83, 376; in U.S.A., 216, 274, 275, 328; legislation against, in U.S.A., 133, 229; in W. Indies, 42, 66, 83, 222, 275, 329, 445, 446; miscellaneous food-plants of, 43, 83, 446, 511, 690, 766; carriage of, by wind, 42, 216; resting stage in larvae of, 42, 83, 376, 446, 511, 690; other bionomics of, 532, 766; natural enemies and biological control of, 112, 274, 275, 329, 704: other measures against. 66, 133, 229, 274, 328, 408, 446, 504, 766; characters of, 249; bibliography of literature on, 247.

Platyedra malvella, bionomics of, on hollyhock in Britain, 33; on cotton, etc., in Uzbekistan, 249;

characters of, 249.

Platyedra vilella, bionomics of, on cotton in Persia, 249; on wild malvaceous plants in Uzbekistan, 249; characters of, 249.

platyedrae, Elasmus.

Platygaster, parasite of Sitodiplosis mosellana in Japan, 670; bionomics of, parasitising Contarinia tritici in Sweden, 231.

Platygaster herricki (parasite of Mayetiola destructor), bionomics

of, in U.S.A., 121.

Platygaster hiemalis (minutus), parasite of Mayetiola destructor in Morocco, 340.

Platygaster pleuron, imported into U.S.A., against Mayetiola des-structor, 329.

Platygaster tuberosula (parasite of wheat gall-midges), bionomics of, in Sweden, 231.

platyhypenae, Euplectrus.

Platymetopius acutus, on narcissus in U.S.A., 288.

Platymetopius frontalis, on narcissus in U.S.A., 288.

platynotae, Microbracon.

Platyphytoptus, gen. nov., species of, on pines in California,

Platypleura kaempferi, measures against, on pear in Japan, 604.

Platypus, incidence of species of, on poisoned and girdled Styrax in Malaya, 453.

Platypus cupulatus, on Styrax in Malaya, 453.

Platypus cylindrus, in green oak timber in Britain, 707; on Acer pseudoplatanus in France, 316.

Platypus gerstaeckeri, on Podocarpus and Endospermum in Fiji, 702.

Platypus lepidus, on Styrax in Malaya, 453.

Platypus solidus, Malaya, **453.** on Styrax

Platypus suffodiens (with var. malayensis), on tapped jelutong trees in Malaya, 110.

plebejus, Heteronychus.

plejadellus, Chilo.

Plesiocoris rugicollis, bionomics of. on apple in Germany, 234, 250; measures against, 250.

Plesispa reichei, synonym of, 499.

pleuron, Platygaster.

Pleurotropis, parasite of Cassida spp. in Italy, **528**; parasite of Mayetiola destructor in Morocco,

Pleurotropis benefica, parasite of Cephus pygmaeus in Britain, 102. Pleurotropis epigonus (see P. metal-

Pleurotropis lithocolletidis, parasite of Phytomyza ilicicola in Maryland, 541.

Pleurotropis metallica, parasite of Agromyza simplex in Britain, 33.

Plochionus amandus, predacious on Homaledra sabalella in Florida, 22.

Plodia, tests of methyl bromide on, in stored grain in U.S.A., 389.

Plodia interpunctella, in Germany, 460: in butter nuts from Dutch Guiana, 529; in Holland, 529, 599; in U.S.A., 29, 322, 389; in flour warehouses, 322; in dried fruit, etc., 29, 389, 460, 599; bionomics of, 29, 460; effect of low temperatures on, 322, 599; schizogregarine infesting, 598; methyl bromide against, 389; characters of eggs of, 196.

Plum, pests of, in Britain, 191, 297, 612, 614, 708, 709; Macropsis trimaculata on, in Canada, 173; new termite on, in Canary Islands, 413; pests of, in Czechoslovakia, 657, 659; Hyponomeuta padellus on, in Cyprus, 259, 260; pests of, in Germany, 50, 52, 718; Hyalopterus arundinis on, in Finland, 201; Paratetranychus pilosus on, in Holland, 465, 466; Hoplocampa on, in Italy, 5; Lymantria dispar on,

in Jugoslavia, 759; Monosteira unicostata on, in Morocco, 340; Tortricids on, in New Zealand, 57, 58; Hoplocampa on, in Palestine, 514; pests of, in Russian Union, 305, 306, 481, 530; Aglaope infausta on, in Spain, 697; Cydia funebrana on, in Switzerland, 99; pests of, in U.S.A., 19, 62, 173, 287, 560; Eutettix disciguttus not surviving on, 208; insecticides causing injury to, 100, 466, 708; fumigation of fruit of, against Aspidiotus perniciosus, 307.

Plum, Beach (see *Prunus maritima*). Plum, Japanese (see *Prunus*

salicina).

Plum Aphis, Mealy (see Hyalopterus arundinis).

Plum Curculio (see Conotrachelus nenuphar).

Plum Leafhopper (see Macropsis trimaculata).

Plum Sawfly (see Hoplocampa flava). Plum Scale (see Lecanium corni). Plumbago zeylanica, insecticidal properties of, 361.

plurispinosa, Megaselia.

Plusia brassicae, measures against, on cabbage in U.S.A., 13, 285, 399,651; tests of insecticides on, 388.

Plusia chalcites, on potato in Hawaii, 214; on beans and potato in New South Wales, 335. Plusia festata, on Juncus in Japan,

672.

Plusia gamma, in Russian Union,
351 352 354 475 Coccinellide

351, 352, 354, 475; Coccinellids predacious on, on cotton, 354; effects of temperature and humidity on, 351, 352, 475.

Plusia peponis, on cucurbits in Ceylon, 358.

plusiaeformis, Stretchia.

Plutella maculipennis (cruciferarum) (on crucifers), in Australia, 449; in India, 409; in New Zealand, 449; in Tanganyika, 113; in U.S.A., 13, 28, 399, 651; possible migration of, 28; Podisus maculiventris reared on, 13; parasites and biological control of, 13, 449; other measures against, 399, 409, 651.

pluto, Pimpla.

Pnyxia scabiei, Actinomyces scabies associated with, on potato in Ontario, 540.

Poa spp., Geomyza tripunctata on, in Britain, 382; grasshoppers

damaging, in Iowa, 23; Pyrausta nubilalis experimentally fed on, in Russia, 475.

Podagrica puncticollis, on cotton in Sudan, 513.

podana, Tortrix (Cacoecia).

Podapolipus berlesei, attacking Schistocerca paranensis in Argentina, **147**.

Podisus maculiventris (predacious on Coleoptera and Lepidoptera) introduced into France, 595; bionomics and rearing of, in U.S.A., 13, 595, 596.

Podocarpus, Coccid on, in Japan, 604.

Podocarpus vitiensis, pests of, in Fiji, 702.

Podonta daghestanica, soil types in relation to, on rubber-producing plants in Ukraine, 72.

Podothrips, scope and subgenera of, 457.

Podothrips lucasseni, on sugar-cane in Java, 457; synonym of, 457.

Podothrips (Melampodothrips) oryzae, subgen. et sp. n., on rice in Siam, 457.

poecila, Mormidea; Oncideres.

Poecilips, characters and synonymy of, **528.**

Poecilips nuciferus, sp. n., in butternuts imported into Holland from Dutch Guiana, 529.

Poecilonota variolosa, on poplar and ash in Russian Union, 429.

Pogonomyrmex occidentalis, association of Tenebrionids with, in U.S.A., 292.

Poland, cereal pests in, 518, 622; pests of flax and hemp in, 255, 622, 719; forest pests in, 4, 154, 256, 523; pests of hops in, 622, 719; orchard pests in, 98, 99; tobacco pests in, 72; pests of vegetable and root crops in, 99, 255, 256, 720; (Tatra Mts.), Coccids of, 772; Halticids of, 255; beneficial insects in, 98, 256, 623; summary of plant quarantines in, 385; properties and standardisation of tar distillates in, 97, 98, 99, 300.

Polia oleracea, on carnation in Britain, 349; parasites of, on beet in Italy, 528; Tachina civilis ovipositing on, 303.

Polia pisi, Tachina civilis ovipositing on, 303.

poligraphus, Polygraphus; Colias hyale.

Polistes macaensis, predacious on Lepidoptera in Seychelles, 433. politus, Chilocorus.

Pollard (see Bran).

Pollination, relation of insects to. 184, 196; pest control in relation to. 400.

Polyamia inimica, on narcissus in U.S.A., 288.

Polychrosis botrana (Vine Moth), utilisation of bacteria against, in France, 619; in Germany, 4, 51, 154, 419, 521, 661; in Jugoslavia, 720; in Switzerland, 299; list of food-plants of, 4; bionomics of, 661, 720; laboratory breeding of, 4, 721; measures against, 51, 154, 299, 419, 521; tests of duration of activity of contact insecticides against, 421.

Polychrosis viteana, measures against, on vines in U.S.A., 323. Polyembryony, biological aspects

of, 248.

Polygonum Pyrausta aviculare, nubilalis reared on, 475.

Polygraphus poligraphus, on Pinus sylvestris in Estonia, **462.** Polyhedral Disease, Lepidoptera

infected with, 70, 205, 234.

polynesialis, Angitia.

Polyphylla fullo, in forests in Russia, 158, 429, 431; soil conditions in relation to, 429, 430; soil fumigation against, 431.

Polypodium vulgare, Heptamelus ochroleucus on, in Britain, 152.

Polystictus versicolor, Xestobium rufovillosum associated with, in timber in Britain, 707.

polytomum, Diprion. pomaria, Typhlocyba.

Pomegranate, Aphis gossypii on, in Korea, 309; thrips on, in Sierra Leone, 240; book on pests of, in U.S.A., 722.

pometaria, Alsophila.

pomi, Aphis.

pomonella, Cydia (Carpocapsa, Laspeyresia); Rhagoletis.

pomorum, Anthonomus.

Pomphopoea sayi, on lupins in Connecticut, 17.

Pongamia glabra, tests of insecticide prepared from, 361.

Popillia japonica, list of food-plants of adults of, in Japan, 54; in U.S.A., 14, 23, 108, 216, 324, 441, 443, 444, 543, 545, 551, 552, 723, 726, 727, 751; legislation against, 384, 727, 751; damaging turf, 727; on shade and fruit trees,

441, 443, 543, 551; on potato, 727; carried by wind, 216; parasites and biological control of, 108, 324, 545, 552, 723, 726; artificially infected with milky disease, 324; other measures against, 23, 324, 384, 441, 443, 444, 543, 551, 727, 751; review of investigations on control of, 727.

popilliavora, Tiphia.

Poplar (Populus), Stilpnotia salicis on, in Austria, 589; Dicranura vinula on, in Britain, 612; Lamellicorns associated with, in France, 230; pests of, in Germany, 345, 461, 717; S. salicis on, in Hungary, 589; Pemphigini on, in Italy, 696; Doraphis populi on, in Japan, 673; Monosteira unicostata on. in Morocco, 340; Lithocolletis populifoliella on, in Rumania, 193; pests of, in Russian Union, 429, 432; pests of, in U.S.A., 15, 17, 23, 45, 225, 589; Lepidoptera mining leaves of, 193, 440; experiment with Lymantria dispar on, **759.**

Poplar, Lombardy (see Populus nigra italica).

Poppy, pests of, in Czechoslovakia, 657, 659.

Population Studies, of insects, 56, 72, 196, 283, 311, 366, 367, 379, 439, 611, 626, 630, 639, 677, 678.

populi, Melasoma.

populifoliella, Lithocolletis. populitransversus, Pemphigus.

populnea, Saperda.

Populus (see Poplar). Populus nigra italica, drought affecting damage to, by Synaphaeta guexi in California, 650.

Porosagrotis orthogonia (on cereals), in Canada, 170, 751; in U.S.A., 44, 126; bionomics and control of, 751.

porosus, Trialeurodes.

Porricondyla, associated with galls on Cornus in U.S.A., 546.

portentosus, Brachytrypes.

Porthesia hololeuca (see Acyphas leucomelas).

Porthesia virguncula, bionomics of, on Hibiscus sabdariffa in Netherlands Indies, 358.

Porthetria dispar (see Lymantria). Portia Tree (see Thespesia populnea).

Porto Rico, coffee pests in, 222, 223; cotton pests in, 41, 42, 275, 329; miscellaneous pests in, 223, 329, 643: sugar-cane pests in, 39, 40, 43, 223, 288, 289; insecticidal plants and their pests in, 224; Anastrepha spp. in, 325; new Jassids in, Scapteriscus vicinus in, 288 ; 40: insects and plant diseases in, 39, 40, 288; beneficial insects and biological control in, 39, 41, 43, 212, 222, 223, 275, 288, 289, 329, 330; Bufo marinus destroying insects in, 40, 317; development of economic entomology in, 116; Pyrophorus luminosus introduced into Mauritius from, 317.

Portugal, Aelia rostrata on wheat in, 697; Coccids of, 119, 315.

Loxostege bifidalis migrating to

cotton from, in Brazil, 333.

Portulaca oleracea, Phenacoccus solani, on, in Hawaii, 213;

Aphis gossypii on, in Korea,

309; list of insects feeding on, 217.

Post-oak Locust (see Dendrotettix quercus).

postfasciatus, Euscepes.

posticus, Orgyia (Notolophus); Phytonomus (see Hypera variabilis). postvittana, Tortrix.

Potassium Ammonium Selenosulphide, against mites, 131.

Potassium Chromate, for treating timber against *Lyctus*, **190**.

Potassium Cyanide, for generating hydrocyanic acid gas (q.v.), 53, 702, 703.

Potassium Hydroxide, as an activator for nicotine sulphate, **332.**

Potassium Permanganate, used with formalin for fumigation, 539, 579; and quick-lime, spraying with, against mites, 72.

Potassium Phosphate, increasing toxicity of lead arsenate to Popillia japonica, 324.

Potassium Selenosulphide, against mites, 648.

Potassium Thiocarbonate, of little value against chicory flies, 313.

Potato, pests of, in Argentina, 147, 290; Leptinotarsa decemlineata on, in Belgium, 368, 719; pests of, in Brazil, 145, 290, 320; pests of, in British Isles, 47, 379, 527, 615, 709, 763; pests of, in Canada, 144, 170, 540, 541; L. decemlineata on, in France, 5, 368, 412, 594, 595, 597; pests of, in Germany, 68, 252, 347, 368, 418, 461, 520, 521, 717, 719; pests of, in Hawaii, 213, 752;

pests of, in India, 361, 409; Eutettix disciguttus on, in Japan, 208; L. decemlineata on, in Luxemburg, 368; Phthorimaea operculella on, in Malta, 469; Épicaerus cognatus on, in Mexico, 11; Plusia chalcites on, in New South Wales, 335; Empoasca benedettoi on, in Palestine, 77; apparently not injured by E. libyca in Spain, 700; pests of, in Russia, 159, 424; pests of, in Switzerland, 70, 92; pests of, in U.S.A., 13, 14, 44, 128, 140, 217, 281, 285, 327, 399, 400, 564, 565, 578, 643, 648, 731, 751; Epicauta spp. on, in Uruguay, 290; Blapstinus substriatus experimentally fed on, 27; conference on control of L. decemlineata on, 467; as a trap crop for Melolontha, 521; symptoms caused by Paratrioza cockerelli on, 281, 578; insects and virus diseases of, 47, 69, 77, 379, 495, 527, 564, 615; insecticides causing injury to, 252, 648; (tubers) legislation dealing with fumigation of, against Popillia japonica in U.S.A., 751: vacuum fumigation of, against L. decemlineata, 597; Epicaerus intercepted cognatus in, Louisiana, 11.

Potato Beetle, Colorado (see Leptinotarsa decembineata).

Potato Leafhopper (see *Empoasca fabae*).

Potato Scab-gnat (see Pnyxia scabiei).

Potato Weevil (see Epicaerus cognatus).

Potentilla, Capitophorus fragariae on, in Britain, 104, 613; pests on, in U.S.A., 288, 291, 736; Pyrausta nubilalis reared on, 475.

poutieri, Parnara. praelonga, Orthezia.

praemorsus, Leptostylus.

praetiosa, Bryobia.

prasina, Chrysopsa; Palomena.

prasinus, Thamnotettix.

pratensis, Lygus.

Praxibulus laminatus, in Australia, 584.

Prays oleellus (on olive), measures against, in Italy, 350; in Spain, 700.

prelli, Chermes (Drevfusia).

Prepodes vittatus, toad destroying, on Citrus in Jamaica, 490. pretiosum, Trichogramma.

Prickly Lettuce (see Lactuca scari- Promecotheca papuana (antiqua),

Prickly Pear (see Opuntia).

prima, Pseudhomalopoda.

Prince Edward Island, Diprion polytomum on spruce in, 487. Prionomerus gudimiri, sp. n., on

avocado in Brazil, 502.

Prionomerus sexgibbosus, sp. n., on avocado in Brazil, 502.

Prionopus perseae (see Meroprion). Prionus laticollis, on Cornus florida

in New York, 547.

Pristiphora, on willow in Japan, 771. Pristiphora abietina (on spruce), in Czechoslovakia, 657; in Germany, 189.

(Pristomerus) agilis, Pristomeridia parasite of Cydia pomonella in

U.S.A., 30.

Pristomerus vulnerator, parasite of Cydia molesta in Italy, 154.

Pritchardia pacifica, pests of, in Samoa, 702.

Privet, Aonidiella aurantii on, in S. Africa, 455; new Eriophyid

on, in California, 650; Ctenopseustis obliquana on, in New Zealand, 58.

Probits, for expressing toxicity of fumigants, **571**, **633**

procerus, Saccharosydne. processionea, Thaumetopoea.

Prochiloneurus bolivari, parasite of Pseudococcus citri in Spain, 701.

Prociphilus fraxinifolii, penetration of setae of, 225.

Prociphilus tessellatus, alimentary canal of, 624.

Procometis trochala, on sugar-cane in Bombay, 261.

Prodenia eridania (see Xylomyges). Prodenia litura, on cotton in Belgian Congo, 315; relation between Thrips tabaci and, on cotton in Egypt, 458; Telenomus nawaii introduced from Fiji into Egypt against, 212; in Formosa, 207; on tobacco and castor in Indo-China, 185, 205; on cassava in Madagascar, 455; in Palestine, 638; bionomics of, 205, 207, 638; measures against, 185, 205, 207, 638.

producta, Euproctis.

Profeltiella soya, on soy beans in Japan, 264; systematic position of, 264.

prolongata, Xiphydria.

Promecopsis canavaliae, sp. n., on Canavalia and Erythrina in Brazil, 106.

measures against, on coconut and other palms in New Guinea Territory, 435, 495; question of value of Oecophylla smaragdina against, 435, 495.

Promecotheca reichei, on coconut in Fiji, 211; on Pritchardia pacifica in Samoa, 702.

Pronomeuta sarcopis, on Hydnocarpus wightiana in Ceylon, 359. Prontaspis citri, on Citrus in Trini-

dad, 10.

pronubana, Tortrix.

Prorops nasuta, failure to establish. against Stephanoderes hampei in Netherlands Indies, 337.

proscarabaeus, Meloë. proserpina, Megamelus.

Prospaltella bemisiae, sp. n., parasite of Bemisia myricae in Japan, 296.

Prospaltella berlesei, imported into Porto Rico, against Aulacaspis pentagona, 329.

Prospaltella citri, sp. n., parasite of Dialeurodes citri in Japan, 296.

Prospaltella elongella, sp. n., parasite of Lepidosaphes gloveri in Louisiana, 41.

Prospaltella perniciosi, utilisation of, against Aspidiotus perniciosus in Idaho, 65, 747.

Prospaltella pulchella, sp. n., imported into Haiti from Delaware with apple twigs, 41.

Prospaltella thoracaphis, sp. n., parasite of Nipponaphis takahashii in Japan, 296.

Proteides mercurius pedro, on Derris elliptica in Porto Rico, 224.

proteus, Omophlus.

Protoparce, on tobacco in U.S.A., 133, 393; cubé of little value against, 133.

quinquemaculata, Protoparce tobacco in Connecticut, 726.

Protoparce sexta, on tobacco U.Ŝ.A., **393, 726.**

Protoparce sexta jamaicensis, parasites of, on field and harvested tobacco in Jamaica, 490.

Protostrophus dianthi, sp. n., on carnation in S. Africa, 612.

proximus, Ips (Orthotomicus).

Prune, pests of, in U.S.A., 29, 62, 291.

Prunes (Dried), pests in, in California, 30, 573.

pruni, F., Hyalopterus arundinis).

pruni, auct., Brachycaudus Anuraphis padi).

pruniella, auct., Argyresthia (see A. ephippella).

prunifoliae, Rhopalosiphum.

prunivora, Enarmonia (Laspeyresia). Prunus, Anomala spp. on, in Korea, 767; species of, serving as winter food-plants of Myzus persicae in New Brunswick, 145; relation of Macropsis trimaculata to species of, 173. (See Cherry, Plum, Sloe, etc.)

Prunus laurocerasus, Aspidiotus perniciosus experimentally devel-

oping on, 305.

Prunus maritima, Cryptocephalus incertus on, in Massachusetts, 646.

Prunus padus, Rhagoletis cerasi not observed on, in Switzerland, 156. Prunus salicina, 173; heat treat-

Prunus salicina, 173; heat treatment of fruits of, against Dacus dorsalis in Formosa, 207.

pryeri, Clania.

Psalidium maxillosum (Black Beet Weevil), in Russia, 94, 432; on tree seedlings, 432; new parasite of, 94.

psallidis, Erythmelus.

Psallus ambiguus, oviposition of, on apple and current in Germany, 234.

Psallus seriatus (Cotton Fleahopper), in U.S.A., 136, 227, 276, 277, 328; factors affecting infestation by, 276; parasites of, 227, 328; measures against, 136, 277.

Psammotis hyalinalis, parasites of,

in Jugoslavia, 760.

Pseudanogmus stironotus, parasite of Cydia pomonella in Victoria, 169.

Pseudaonidia, new parasite of, on Ficus in Japan, 296.

Pseudaonidia articulatus (see Selenaspidus).

Pseudaonidia duplex, new parasite of, on Citrus in Japan, 296.

Pseudaphycus utilis (parasite of Pseudococcus nipae), in Mexico, 586; establishment of, in Hawaii, 586

Pseudhomalopoda prima, parasite of Coccids in W. Indies, 41.

Pseudiastata brasiliensis, sp. n., attacking Pseudococcus brevipes in Brazil, 319.

pseudoaonidiae, Aspidiotiphagus. pseudobrassicae, Rhopalosiphum. pseudocitriculus, Pseudococcus. pseudococcina, Hambletonia. Pseudococcobius terryi (parasite of Pseudococcus spp.), in Hawaii, 137; establishment of, in U.S.A., 137, 138; bionomics of, 138.

Pseudococcus, key to species of, in South-eastern Asia, 607; new parasites of, in Brazil and Colombia, 212; on apple and cherry in Br. Columbia, 59; on vines in California, 748; on pear in Japan, 604; measures against, 59, 748.

Pseudococcus adonidum, measures against, on coffee and kapok in Indo-China, 185, 187; sex ratio and status of males of, 105, 106.

Pseudococcus boninsis, establishment of Pseudococcobius terryi against, on sugar-cane in U.S.A., 137, 138.

Pseudococcus brevipes (Pineapple Mealybug), in Brazil, 319; in Colombia, 212; in Formosa, 263; in Hawaii, 212, 214, 570; in Porto Rico and U.S.A., 329; causing wilt and spotting of pineapple, 570; other foodplants of, 214, 263, 319; bionomics of forms of, 570; natural enemies and biological control of, 212, 214, 319, 329.

Pseudococcus citri, in S. Africa, 202: in Netherlands Indies, 182, 606, 607; sooty fungus associated with, in Italy, 758; Lepdiosaphes beckii recorded as, in Palestine, 638; Trypetid feeding on honey-dew of, in Sicily, 92; in Spain, 701; in U.S.A., 138; on Citrus, 202; on coffee, 182, 606, 607; on vines, 701, 757; other food-plants of, 92, 138, 182, 598; sex ratio and status of males of forms of, 105, 106, 598; other bionomics of, 606, 607; ants associated with, 202, 606, 758; natural enemies and biological control of 607, 701; experimentally parasitised by Pseudococcobius terryi, 138; explanation of immunity of, from Coccophagus gurneyi, 287; measures against, 182, 758; P. lilacinus considered a synonym of, 608.

Pseudococcus coffeae, 607.

Pseudococcus crotonsis, P. tayabanus considered correct name for, 608. (See P. lilacinus.)

Pseudococcus cryptus, on Citrus and coffee in Brazil, 502, 607; morphology of, 502; P. deceptor possibly identical with, 607.

Pseudococcus cuspidatae, sp. n., bionomics of, on Taxus, apple, etc. in U.S.A., 226.

Pseudococcus deceptor, sp. n., on coffee etc. in Java, 607.

Pseudococcus defluiteri, sp. n., on coffee in Java, 607.

Pseudococcus gahani, sex ratio and status of males of, 105.

Pseudococcus kenyae, ant associated with, on coffee etc. in Kenya, 592; not occurring in Tanganyika, 592.

Pseudococcus kraunhiae, on Citrus in California, 20: P. cuspidatae

recorded as, 226.

Pseudococcus lepelleyi, sp. n., on Anona and Lansium in Java, 607. Pseudococcus lilacinus, in India,

592; P. kenyae compared with, 592; question of attribution of name, **608.**

Pseudococcus longispinus (see P.

adonidum).

Pseudococcus maritimus, new parasite of, imported into California from Brazil, 212; on gladiolus in Iowa and Hawaii, 65, 213; sex ratio and status of males of, glands of, wax 105; 184; measures against, 65.

Pseudococcus nipae, on coconut in Cayman Islands, 268; foodplants of, in Mexico and Hawaii, 586; natural enemies and bio-

logical control of, 586.

Pseudococcus pseudocitriculus, n., on Tephrosia in Java, 607.

Pseudococcus pseudofilamentosus, sp. n., food-plants of, in Java, 607.

Pseudococcus sacchari (see Triony-

Pseudococcus tayabanus, considered correct name for P. crotonis, 608. (See P. lilacinus.)

Pseudococcus virgatus (see Ferrisiana).

Pseudococcus vitis, on vines in Malta, 469.

Pseudococcus walkeri, description of

female of, 560. Pseudoepameibaphis, species of, on Artemisia in U.S.A., 772.

pseudofilamentosus, Pseudococcus. Pseudograsserie, new type of, in

Euxoa segetum in France, 164. Pseudoleucopis benefica, introduced New against into Zealand Eriococcus coriaceus, 753.

pseudomagnoliarum, Coccus.

(see Pseudosarcophaga monachae Agria).

Pseudoscorpions, in bee-hives, 664. pseudosolani, Myzus (see Macrosiphum solani)

pseudospretella, Borkhausenia (Acompsia, Hofmannophila).

Pseudotsuga taxifolia, Hemerocampa pseudotsugata on, in Br. Columbia, 293; Dermestes vulpinus timber of, in New South Wales, 451; Navomorpha lineata on, in New Zealand, 266; pests of, in U.S.A., 441, 581; effect of soil treatment with lead arsenate on.

pseudotsugata, Hemerocampa.

psi, Acronicta.

Psidium guayava (see Guava).

Psila nigricornis, on chrysanthemums in Britain, 349.

Psila rosae, measures against, on carrot in U.S.A., 489, 647; injury by Listroderes latiusculus attributed to, 565.

Psilopa nitidula compta, associated with Lixus junci in Morocco, 417.

Psylla mali (on apple), in Czechoslovakia, 659; in Russia, 353, 482; measures against, 659; action of oil emulsions on eggs of, 353.

Psylla pyricola, on pear in Italy,

758.

Psyllid Yellows, of potato, physiology and symptoms of, caused by Paratrioza cockerelli in U.S.A., 281, 578.

Psylliodes attenuata, bionomics of. on hops in Poland, 719; on hemp in Ukraine, 477; measures against, 477, 719.

Psylliodes punctulata, measures against, on beet in Utah, 743; breeding on Cheirinia repanda,

psylloides, Gibbium.

Pterochlorus salignus (see Lachnus). Pterocomma smithiae, technique for rearing Coccinellids on frozen, in Ohio, 649.

Pteroma plagiophleps, on tamarind in Madras, 499.

Pteromalus matsuyadorii, parasite of Dendrolimus spectabilis Japan, 669.

Pteromalus puparum, parasite of Pieris rapae in Japan, 605.

Pteromalus semiclavatus, parasite of Coleophora laricella in Austria,

Pteroptrix maskelli, parasite Rhinocola eucalypti in Zealand, 754.

Ptinus tectus, in dried Hyoscyamus and Datura in Germany, 164. Ptochorvetis simbleuta, bionomics and control of, on tea in Assam, 601. pubescens, Ophonus (see Harpalus rufipes) pudens, Sphingomorpha. pudibunda, Dasychira. pudicus, Carpocoris. puera, Hyblaea. pugnax, Solubea. pulchella, Prospaltella. pulchellus, Hemiteles. pulcher, Dysdercus orientalis. pulchrum, Lecanium L. rufulum). Puliciphora bectinata, sp. n., parasite of Schistocerca paranensis in Argentina, 689. pullata, Zenillia (Clemelis). Pulvinaria cellulosa, on Citrus in Cevlon, 359. Pulvinaria floccifera, on holly in France, 156; food-plants of, in Switzerland, 156, 689. Pulvinaria iceryi (gasteralpha), on sugar-cane in China, 243. Pulvinaria vitis, on maple in Indiana, 126. pumilionis, Chlorops (see C. taeniopus). pumilus, Microdus (Bassus). Pumpkin, scorched by essential oils, punctata, Ceratitis; Hypera. punctaticeps, Perisierola cellularis. punctatifrons, Dacus. punctatum, Anobium. punctatus, Aenasius. puncticollis, Nodonota; Podagrica. punctiferalis, Dichocrocis. punctiger, Isostasius. punctillum, Stethorus. punctiventris, Cleonus (Bothynoderes). punctoria, Angitia (Dioctes, Inareolata). punctulata, Pheidole; Psylliodes. punctulatus, Elasmus. punctum, Stethorus. Punica granatum (see Pomegranate). Punjab, cotton pests in, 180, 532. puparum, Pteromalus. pupivora, Trichospilus. purchasi, Icerya. Purslane (see Portulaca oleracea). pusana, Coniopteryx; Pyrilla. pusilla, Sciara (Lycoria). pusilla, auct., Austroicetes (see A. cruciata).

pusillus, Lophocateres.

pustulans, Asterolecanium. Puto yuccae, new parasite of, in California, 212. putrescentiae, Tyrophagus (Tyroglyphus). Pycnoscelus surinamensis, measures against, on rose in New Hampshire, 20. pygmaeus, Cephus; Scolytus. Pyralis farinalis, characters of eggs of, 196. pyralis, Sacadodes. Pyrameis (see Vanessa). Pyrausta diniasalis, parasite of, in Manchuria, 296. Pyrausta nubilalis (European Corn Borer), in Canada, 329, 486, 536; in Formosa, 309; in Italy, 528, 712, 713; in Manchuria, 296; in Poland, 622; in Spain, 695; in Russian Union, 159, 257, 303, 475, 476; in U.S.A., 14, 126, 216, 285, 326, 329, 367, 505, 545, 580, 590, 648, 723, 729, 730; legislation against, in U.S.A., 750; varieties and time of planting of maize in relation to, 14, 326, 505, 536, 729; on cotton, 309; on dahlia, 723; on hemp, 159, 475, 476, 622; types of damage to hops and millet by, 622; on potato, 14, 285, 731; on sorghum, 528, 750; bionomics and physiology of, 54, 216, 257, 440, 475, 476, 486, 622, 713; sampling of populations of, 367; parasites and biological control of, 14, 275, 296, 326, 329, 623; Tachina ovipositing on, civilis 303: measures against, 285, 623, 648, 713, 714, 723, 750. Pyrausta phoenicealis, on Perilla in Japan, 771; predacious enemies of, on Ocimum sanctum in Seychelles, 433. Pyrethrins, contents of, in pyrethrum dusts and sprays, 17, 125, 129, 279, 347, 348, 382, 397, 509, 534, 550, 559, 563, 645, 646, 725, 740; effect of hydrogenation of, 120, 567; methods for determining, 330, 331, 600, 695; partial synthesis of, 368.

Pyrethrolone, 120, 368, 694.

(including

against Aphids, 464, 523, 604,

740; against Capsids, 279, 376,

534; against Jassids, 134, 282, 508, 534, 559, 609, 646; against

other Rhynchota, 10, 180, 397,

404, 533, 725, 726, 755; against

extracts).

Pyrethrum

Coleoptera, 17, 194, 197, 265, 268, 328, 382, 400, 418, 421, 522, 550, 566, 572, 645, 646, 648, 691, **752, 755, 762, 763;** against Diptera, **466, 541, 655;** against Lepidoptera, 22, 74, 172, 279, Lepidoptera, 22, 74, 172, 273, 312, 327, 328, 347, 421, 422, 533, 582, 647, 664, 735, 736, 752, 768; against Lepismids, 37; against mites, 583, 646, 648, 712; against sawflies, 348; against thrips, 129, 508, 553; applied to soil against insects, 40, 53; mixed with stored wheat against pests, 125; effect of, on beneficial insects, 180, 559; tests of toxicity of, to insects, 17, 35, 109, 312, 347, 399, 421, 422, 508, 563, 567, 572; harmless to sheep, 282; dusting with, 10, 17, 35, 53, 129, 172, 180, 197, 268, 279, 312, 327, 328, 347, 348, 376, 397, 404, 418, 421, 466, 508, 522, 534, 550, 563, 565, 566, 583, 609, 645, 646, 647, 655, 691, 712, 725, 726, 735, 736, 752, 755, 763; carriers for, 17, 129, 172, 197, 279, 397, 524, 550, 609, 645, 647, 726, 736 534, 550, 609, 645, 647, 725, 726, 736, 752, 755; impregnated and activated dusts of, 509, 550, 725; adhesiveness of dusts of, 197; formulae for, in sprays, 35, 74, 134, 382, 534, 559, 604, 664, 725, 736, 755; preparation of extracts and sprays of, 412; comparison of water suspensions and extracts of, 648; tests of duration of activity of, 421, 422; applied from aeroplanes, 134, 508; and Bordeaux Mixture, 648; and calcium arsenate, 559; and cubé, 740; and derris, 268, 421, 422, 522, 523, 553, 740, 762; and lime, 559; unsatisfactory with lime-sulphur and iron sulphate, 534; and oil emulsions, 74, 508, 736; atomised sprays of oils containing, 134, 400, 404, 464, 582; and soap, 559, 604, 646, 762, 768; Ultrawet as an emulsifier for, 725; chemistry, constituents and evaluation of, 120, 330, 331, 368, 651, 694, 695, 772; proprietary forms of, 550, 725, 735, 752; standardisation of insecticides containing, in Switzerland, 156; other insecticides compared with, 17, 35, 279, 312, 328, 347, 348, 522, 523, 645, 735, 736.

pyri, Eriophyes; Stephanitis. pyricola, Hoplocampa; Psylla. Pyridine, of little value against chicory flies, 313; soil treatment with, against Euscepes batatae, 447; and soap, spraying with, against Contarinia nasturtii, 467. Pyrilla (on sugar-cane), in Ceylon, 8; in India, 8, 241, 243, 261; natural enemies of, 242, 243; identity of species of, 8. Pyrilla aberrans, 8.

Pyrilla aberrans, 8.
Pyrilla perpusilla, 8.
Pyrilla pusana, 8.

pyrillae, Ooencyrtus; Tetrastichus.

pyrina, Zeuzera. pyrivora, Contarinia.

Pyrochroa coccinea, on sycamore, beech and oak in France, 316. Pyrocide, Dry, 550, 725, 735, 736, 752; preparation of, 550, 725.

Pyroderces paroditis, on coconut in Pacific Islands, 497.

Pyroderces rileyi, on maize in S. Carolina, 643.

Pyroderces simplex, on cotton in Formosa, 309.

Pyrophorus bellamyi (predacious on noxious insects), bionomics of, in Guatemala, 493; introduced into Hawaii, 493; unlikely to reduce value of Campsomeris marginella modesta, 493.

Pyrophorus luminosus (attacking Lachnosterna smithi), establishment of, in Barbados, 222; introduced into Mauritius, 316, 317; Bufo marinus destroying, in Porto Rico, 317.

Q.

quadraria, Thalassodes. quadratum, Piesma. quadridens, Ips (Pityogenes). quadrimaculatus, Bruchus (see B. maculatus). quadrinotata, Aphrophora; bela (Arbela). quadrinotatus, Thamnotettix (Cicadula). quadripes, Xylotrechus. quadripustulata, Winthemia. quadripustulatus, Exochomus. Quarantine, pests intercepted in: in Germany, 101; in U.S.A., 24, 116; in Vancouver, 483. Quassia, spraying with, against plum sawfly, 5, 191, 615; proprietary spray containing, against Anthonomus pomorum, harmless to bees, 5; preparation

of sprays of, 191.

Quassin, preparation and constituents of, 120, 368, 651, 694.

quatuordecimpunctata, Crioceris. Quebec, apple pests in, 541; Bregmatothrips iridis possibly in, 273; outbreak of Cirphis unipuncta in, 145; Cydia nigricana on peas in, 60, 623; Diprion polytomum on spruce in, 77, 487, 538; parasites and biological control of noxious insects in, 77,

Queensland, apple pests in, 149, 495; Thysanoptera on banana

in, 211, 494; pests of Citrus in, 268, 494, 754; new Coccid on Eremocitrus in, 226; miscellaneous pests in, 494; pests of papaya in, 246; pests damaging pasture grasses in, 267, 449, 494; sugarcane pests in, 56, 151; pests of timber in, 449, 450; grasshoppers in, 449, 494, 583, 676, 686; beneficial insects in, 151, 450, 494, 502; establishment of Bufo marinus against Lepidoderma in, 56; epizootic in Lepidiota caudata in, 267; biological control of weeds in, 87, 450; restrictions on movement of vines from, against Phylloxera,

quercifolia, Gastropacha.

608.

Quercus ilex var. suber, pests of, in Morocco, 46.

quercus, Dendrotetrix.

Quillaia saponaria, use of bark of, in sprays against Coccids, 221.

Quince, leaf-cutting bees damaging, in Germany, 522; Lymantria dispar on, in Jugoslavia, 759; Cydia pomonella on, in Palestine, 500; Eriosoma lanigerum cn, in Réunion, 706; pests of, in U.S.A., 399, 647.

quinquemaculata, Protoparce. quinquepunctatus, Tychius. quinquesignata, Hippodamia.

R.

Radiation, methods of measuring, for biological purposes, 772. radicis, Hylobius.

vadiolus, Apion.

Radish, Tetranychus telarius on, in Formosa, 674; pests of, in Japan, 605, 769, 770; Hylemyia flavopicta on, in Sakhalin, 770. Ragi (see Eleusine coracana). Ragwort (see Senecio jacobaea).

Rainfall, relations of locusts and grasshoppers to, 140, 169, 198, 199, 372, 426, 580, 583, 584, 628, 629, 666, 678, 681, 683, 755, 756, 769; relations of other insects to, 4, 27, 47, 92, 104, 151, 157, 158, 174, 223, 279, 283, 333, 374, 375, 376, 377, 406, 418, 494, 532, 640, 661, 710; resistance of spot deposits of sprays to, 522.

Raisin Moth (see Ephestia figuli-

lella).
Raisins, in baits for weevils, 172.

270, 393, 566. (See Fruit, Dried). Rajputana, locusts in, 628, 683. ramulorum, Myrmelachista ambigua.

rapae, Pieris (Ascia). rapax, Aspidiotus.

Rape, pests of, in Czechoslovakia, 657, 718; Meligethes aeneus on, in Holland, 299; pests of, in Hungary, 197; susceptibility of varieties of, to Eurydema ornatum, 718.

rapo, Tetrastichus (Geniocerus). Rarotonga, Tetrastichus giffardianus

shipped to, from Fiji, 212.

Raspberry, Lycaenopsis argiolus on, in Britain, 296; pests of, in U.S.A., 45, 136, 288, 328; thio-diphenylamine causing injury to, 328.

Raspberry Fruit-worm (see *Byturus* unicolor).

raui, Asprothrips.

Ray Grass, French (see Arrhena-therum).

reaumurana, Cydia splendana.

recurva, Phoracantha.

Recurvaria nanella, measures against, on peach and almond in Cyprus, 108.

Recurvaria piceaella, measures against, on spruce in New York,

270.

Red Arrow, 735, 736.

Red-backed Cutworm (see Euxoa ochrogaster).

Red-legged Earth Mite (see *Halo-tydeus destructor*).

Red-shouldered Leaf Beetle (see Monolepta rosea).

vedenianus, Microtermes.

redikorzevi, Eberhardia (Cosmoglyphus).

reducta, Leptocneria (Lymantria).

regina, Phormia.

reichei, Plesispa; Promecotheca. reicherti, Brontispa (see B. longissima).

reitteri, Rhadinoceraea.

vemulus, Trichacis. remus. Telenomus. renalis, Antrocephalus. renipustulatus, Chilocorus. repetita, Anasa.

repleta, Drosophila.

Resin, in preparations for banding. 202, 480, 495, 554; uses of, in sprays, 148, 361, 542, 704, 755.

Resin Residue, as an adhesive and spreader for sprays, 443, 542, 543, 553, 547, 548, 551, 552; reducing toxicity of derris, 553. resinella, Rhyacionia (Evetria).

Respiration, in insects, bibliography

of papers on, 48.

reticulana, Tortrix (Cacoecia). reticulata, Analeptes (Diastocera).

Reticulitermes flavipes, in France, 314; in Germany, 193; in U.S.A., 285, 748; attacking turnips, 285; in building timbers, 193, 285, 314, 748; measures against, 748.

Reticulitermes hesperus, soil fumigation with methyl bromide against,

in U.S.A., 389.

Reticulitermes lucifugus, in building timber in France, 314.

Retithrips aegyptiacus, new parasite of, in Tanganyika, 610.

Retithrips syriacus, new parasite of, in Gold Coast, 610.

Réunion, miscellaneous pests in, 706; Chionaspis tegalensis on

sugar-cane in, 501.

Reviews: -- Andersen (K. T.), The Grain Weevil, 519; Appel (O.) Ed. (in Sorauer, P.), Plant Protection, 703; Brown (H. B.), Cotton . . . , 2nd Edn., 722; Chopard (L.), La biologie des Orthoptères, 629; de la Torre-Bueno (J. R.), A Glossary of Entomology, 11; Glover (P. M.), Lac Cultivation in India, 311; Greimer (K.) & Michael (H.), Handbook of the practical Disinfector, 347; Imms (A. D.), Recent Advances in Entomology, 2nd Edn., 47; Keen (E. P.), Insect Enemies of Western Forests, 400; Manolache (C. I.), Fauna of the Litter of Larch and Oak in Rumania, 193; Obarski (J.), The Enemies of Tobacco Plants and their Control, 72; Quayle (H. J.), Insects of Citrus and other Subtropical Fruits, 722; Schmitschek (E.), Key to . . . Coleoptera injurious in Forestry, 312; Smith (K. M.), A Text-book of Plant

Virus Diseases, **68**; Snedecor (G. W.), Statistical Methods, **688**; Stepantzev (I. N.), Kosobutzkii (M. I.) & Lyubishchev (A. A.), The Interpretation of Data of entomological and phytopathological Records, 94; Weidner (H.), Keys to Store Pests and Household Vermin in Central Europe, 196; Zacher (F.), Maintenance and Breeding of Pests of Stored Products, 468; A Guide to the Insects of Dalbergia sissoo, 531.

Rhabditis hambletoni, sp. n., relation of, to Gasterocercodes brasiliensis in Brazil, 504.

Rhabdocnemis obscura, on sugarcane in Queensland, 151.

Rhaconotus scirpophagae, parasite of Scirpophaga in India, 243, 602; bionomics of, 602.

Rhadinoceraea reitteri, bionomics and control of, on Iris in Italy,

598.

Rhadinosomus lacordairei, measures against, on strawberry and Eucalyptus in Tasmania, 209.

rhagoleticolus, Opius.

Rhagoletis cerasi (on cherry), in Czechoslovakia, 657; in France, 526; in Germany, 156, 346; in Italy, 712, 713; in Spain, 697; problem of food-plants of, in Switzerland, 156, 300; bionomics of, 156, 300, 346, 697; measures against, 300, 347, 526, 697, 713.

Rhagoletis cingulata, measures against, on cherry in New York,

489.

Rhagoletis fausta, on cherry in U.S.A., 44, 489; measures against, 489.

Rhagoletis pomonella, in Quebec, 541; in U.S.A., 16, 269, 333, 401, 559, 561, 647, 725; bionomics of, 725; experiments with different foods and types of cages for rearing, 561; measures against, on apple, 16, 269, 333, 401, 541, 559, 725; (blueberry race), measures against, in Maine,

Rhagoletis ribicola, bionomics and control of, in U.S.A., 60, 61.

rhamni, Aphis.

Rhamnus frangula, not attacked by Rhagoletis cerasi in Switzerland, 156.

Rhaphidopalpa femoralis, on cucurbits in Japan, 605.

Rhaphidopalpa foveicollis, on melon in Algeria and India, 371, 765; rotenone dusts against, 371.

Rhembastus coffeae, sp. n., on coffee in Nyasaland, 372.

Rhicnopeltella eucalypti, in Australia, 754; bionomics of, on Eucalyptus globulus in New Zealand, 753.

Rhinaria perdix, on strawberry in Tasmania, 208.

rhinoceros, Oryctes.

Rhinocola eucalypti, parasite of, on Eucalyptus globulus in New Zealand, 754.

Rhinotermes lamanianus, soil fumigation against, damaging trees in

Tanganyika, 113.

Rhipiphorothrips cruentatus, bionomics and control of, on vines

in India, 87, 88.

Rhizobius lophanthae (see Lindorus). Rhizobius ventralis (predacious on Coccids), in New South Wales, 753; establishment of, in New Zealand, 753.

rhizoglyphoides, Eberhardia (Cosmo-

glyphus). Rhizoglyphus echinopus, on Allium

in Far East, 674.

Rhizoglyphus hyacinthi in Connecti-

Rhizoglyphus hyacinthi, in Connecticut, 246.

Rhizopertha dominica, in stored cassava tubers in Madagascar, 455; in stored ragi in Mysore, 204; in stored cereals in U.S.A., 116, 218; measures against, 116, 204, 218, 455; test of HCN on, 741.

Rhizopus nigricans, Nezara associated with, causing boll-rots in Egypt, 457.

vhizotrogoides, Hoplochelus.

Rhizotrogus aestivus, damaging lawns in France, 229; bionomics of, in forests in Russia, 429, 431. Rhizotrogus solstitialis (see Amphi-

mallus).

Rhodanates (see Thiocyanates).

Rhode Island, new thrips on pines in, 329.

Rhodesia, Northern, coffee pests in, 500; pests of cotton, etc., in, 177, 366, 500; Nomadacris septem-fasciata in, 114, 683, 687; summary of plant quarantines in, 134.

Rhodesia, Southern, cotton pests in, 177, 366; pests of field and stored maize in, 203, 366, 608; Myzus persicae apparently transmitting new virus disease of

tobacco in, 377; Nomadacris septemfasciata in, 410; new weevils in, 106; beneficial insects in, 89, 610; summary of plant quarantines in, 134.

rhododendri, Stephanitis (Lepto-

byrsa).

Rhododendron, pests of, in Britain, 350; Coccids on, in U.S.A., 226, 547. (See Azalea.)

rhodoproctalis, Scirpophaga.

Rhoecocoris sulciventris, bionomics and control of, on Citrus in Australia, 754.

Rhogas, parasite of Limacodids in Ceylon, 639; parasite of Anomis

luridula in Peru, 627.

Rhogas unicolor, parasite of Stilpnotia salicis in Europe, 576; attempted establishment of, in U.S.A., 576; bionomics of, 576, 577.

Rhopalosiphum, measures against, on cotton in S. Carolina, 645.

Rhopalosiphum abietinum, on Picea engelmanni in Austria, 715.

Rhopalosiphum lahorensis (see R. rufomaculatum).

Rhopalosiphum maidis (see Aphis). Rhopalosiphum prunifoliae, Aphis maidis recorded as, on cereals in Argentina, 31, 289; measures against, on apple in New Jersey, 738.

Rhopalosiphum pseudobrassicae, bionomics and control of, on crucifers in India, 241; Coccinellids reared on frozen, in Ohio, 649.

Rhopalosiphum rufomaculatum, on chrysanthemum in Britain, 297.

Rhopobota naevana (vacciniana), on cranberry in U.S.A., 172, 563, 646; bionomics and control of, 172.

Rhubarb, Empoasca fabae on, in Delaware, 443; Maladera holosericea on, in Germany, 194.

Rhus javanica, Aphids producing tannin galls on, in Far East, 264, 265.

rhusicola, Kaburagia.

Rhyacia ypsilon (see Agrotis).

Rhyacionia buoliana (on pines), in Czechoslovakia, 657; in Germany, 312; in Sweden, 634; in U.S.A., 329, 547, 745; parasites and biological control of, 329, 745; other measures against, 547, 634.

Rhyacionia resinella, on pines in

Germany, 189.

Rhyacionia turionana (on pines), in Czechoslovakia, 657; in Germany, 312.

Rhyacionia turionana var. mughiana. bionomics of, on pines in Austria,

663.

Rhynchites, in Russia, 478: measures against, on fruit trees, 297, 478, 621.

Rhynchites aequatus, on apple in Britain, 297.

Rhynchites auratus, food-plants of, in Russia, 432.

Rhynchites coeruleus, on apple in Britain, **297.**

Rhynchites nanus, bionomics of, on

willows in Germany, 664. Rhynchites pauxillus, bionomics of, on apple and pear in Czechoslovakia, 620.

Rhynchites planirostris (see R

Rhynchites tomentosus, on willow in Germany, 665.

rhynchophorae, Parabillaea.

Rhynchophorus ferrugineus (on coconut), in Ceylon, 360; in Cochin, 452.

palmarum, Rhynchophorus parasite of, in Argentina, 689.

Rhynocoris fuscipes, predacious on Laphygma exigua in India, 55. Rhynocoris violentus, predacious on

Nomadacris septemfasciata Nyasaland, 114.

Rhyssa persuasoria, establishment of, against Sirex noctilio in New Zealand, 58.

vibicola, Rhagoletis.

ribis, Aphidius. Rice, Parnara mathias on, in Ceylon, 358; pests of, in China, 244, 245, 498, 674; Prodenia litura on, in Formosa, 207; pests of, in Br. Guiana, 369; pests of, in India, 55, 240, 360; pests of, in Japan, 53, 167, 265, 309, 439, 669, 670, 671, 672, 770, 771; Naranga aenescens on, in Korea, 767; Sesamia inferens on, in Loochoo Islands, 264; pests of, in Louisiana, 218; Parnara spp. on, in Madagascar, 118; Lema tristis on, in Manchuria, 767; new thrips on, in Siam, 457 pests of, in Sierra Leone, 240; pests of, in Tanganyika, 113; insects and diseases of, 167, 386, 439; effects of infestation by certain insects on, 218, 671.

Rice (Stored), pests of, and their control, 31, 95, 158, 218, 240,

311, 336, 584, 639, 770; question of susceptibility of polished, to Calandra oryzae, 158, 584; in baits for crickets, 528; (flour), as an emulsifier for oils, 185, 187.

Rice Borer (see Chilo simplex). Rice Weevil (see Calandra oryzae). Ricinus communis (see Castor).

rickseckeri, Carpophilus.

rilevi, Eriosoma: Pvroderces.

rimac, Ipobracon.

Ripersia sacchari, on sugar-cane in Îndia, **55.**

Riptortus pedestris, on Citrus in Ceylon, 359.

Robinia, Lecanium corni on, in Germany, 251; pests of, Russian Union, 429, 432.

Robinia pseudacacia, leaf-cutting bees damaging, in Germany, 522; pests of, in U.S.A., 549; Lymantria dispar not surviving on, 759.

robiniella, Parectopa.

roborator, Pimpla (Exeristes).

robusta, Frankliniella (see Kakothrips pisivorus); Hypsipyla. Rocamboles, Rhizoglyphus echinopus

on, in Far East, 674.

rodionovi, Caloglyphus.

Rodolia cardinalis (predacious on Icerya purchasi), in Bahamas, 491; utilisation of, in Libya, 260; establishment of, in Porto Rico, 223.

Rodolia chermesina, predacious on Icerva sevchellarum in Sevchelles.

433.

Rodolia guerini, predacious Icerva pilosa in India, 243.

Rohdea japonica, Brithys crini on, in Formosa, **53.**

Rondania cucullata, parasite of Cleonus mendicus in Italy, 527.

Rondania dimidiata, R. cucullata recorded as, 527.

ronnai, Melaloncha.

rosaceana, Tortrix (Cacoecia). rosae, Macrosiphum; Psila.

rosana, Tortrix (Cacoenia).

Rose, Aonidiella aurantii on, in S. Africa, 455; new thrips on, in Australia, 318; pests of, in Britain, 104, 297, 711, 712; pests of, in Ceylon, 359; pests of, in Germany, 522, 717; Paratetranychus pilosus on, in Holland, 465; Eutettix disciguttus on, in Japan, 208; Lymantria dispar on, in Jugoslavia, 759; Megachile seychellensis on, in Seychelles, 433; pests of, in U.S.A., 20, 136, 141,

287, 291, 727; insecticides causing injury to, 711, 712.

Rose Leaf Beetle (see Nodonota puncticollis).

vosea, Monolepta. voseipennis, Nabis.

rosenmülleri, Monochamus sartor. Rosette Disease, of ground-nuts, transmitted by Aphis laburni in Africa, 114, 531, 592; of lilies,

Africa, 114, 531, 592; of filles, transmitted by A. gossypii in Bermuda, 626.

roseus, Anuvaphis (Aphis).

vostrata, Aelia.

Rotenone, dusting with, 16, 100, 172, 291, 327, 328, 371, 508, 691, 714, 725; carriers for, 172, 327, 371, 725; not increasing efficiency of oil-impregnated dusts against Cydia molesta, 544: spraying with, 17, 291, 327, 328, 541, 648, 714, 752; and Bordeaux Mixture, 648; and copper ammonium silicate, 752; and sulphur, 648; contents of, in derris (q.v.), 15, 17, 34, 35, 120, 128, 183, 224, 258, 274, 299, 318, 333, 347, 348, 361, 371, 372, 382, 383, 397, 399, 443, 543, 547, 550, 553, 563, 614, 645, 646, 651, 652, 675, 708, 735, 736, 739, 740, 741, 743, 766; comparison of toxicity to insects of other constituents of derris and, 616, 632; contents of, in cubé, 17, 120, 128, 129, 132, 279, 319, 333, 371, 372, 396, 399, 489, 543, 547, 553, 566, 645, 646, 647, 651, 708, 735, 736, 740, 741, 743; Mundulea containing, 34, 35, 36, 37, 183, 258, 361; other plants containing, 29, 224, 319, 749; classification and distribution of plants yielding, 67, 318, 319; evaluation of plants containing, 29, 37, 120, 183, 258, 331, 616, 631, 739; chemistry and properties of, 67, 120, 368; translocation of, in bean plants, 332.

Rotiboellia exaltata, Nezara chloris on, in Sierra Leone, 240.

votunda, Contheyla.

rotundatum, Gymnosoma.

rotundatus, Stethorus.

rotundicolle, Diploschema.

Rowan (see Sorbus aucuparia). rowani, Telenomus (Phanurus).

Rubber, Leucopholis pinguis on, in Cevlon, 358; pests of, in Indo-China, 186, 187; Oryctes rhinoceros breeding in decaying stumps of, in Malaya, 434. Rubber (Manufactured), attacked by Gibbium psylloides in Germany, 519; ants boring in, in Netherlands Indies, 262.

rubens, Ceroplastes.

rubi, Anthonomus; Monophadnus (Monophadnoides).

rubiella, Incurvaria (Lampronia).

rubigalis, Phlyctaenia.

rubiginosa, Cassida.

rubivora, Ceratitis.

rubricosta, Euproctis (see E. producta).

rubripes, Brachymeria (Chalcis); Coptocercus.

rubrocinctus, Selenothrips.

rubus, Batocera.

rudisana (rudis), Acropolitis.

rufa, Formica; Solenopsis geminata.

ruficollis, Dysdercus.

ruficrus, Apanteles.

rufifemur, Eulimneria.

rufimanus, Bruchus.

rufipenne, Callidium.

rufipes, Angitia; Crepidodera (Derocrepis); Harpalus; Hy-

(Derocrepis); Harpaius; Hy lastes (Hylurgopinus); Necrobia. rufiscutellaris, Glypta.

rufiventris, Oxycarenus. rufobrunnea, Oncopera.

rufocuprea, Anomala. rufomaculatum, Rhopalosiphum.

rufovillosum, Xestobium.

rufulum, Lecanium. rugicollis, Minthea; Plesiocoris.

rugosa, Lachnosterna (Phyllophaga). rugosostriatus, Otiorrhynchus.

rugulosus, Scolytus. ruidum, Ectatomma.

in, 134.

Rumania, fauna of forest litter in, 193; orchard pests in, 119,

192; vine pests in, 71, 721; leaf-mines caused by insects in, 192; Locusta migratoria in, 610; beneficial insects in, 71, 192; summary of plant quarantines

Rumex, Pegomyia bicolor apparently restricted to, in France, 314.

Rumex patientia, Listronotus latiusculus on, in New Jersey, 565.

rumicis, Acronicta; Aphis.

runneri, Frankliniella.

ruralis, Pikonema (Nematus); Voria.

Russia (U.S.S.R.), beet pests in, 94, 157, 235-237, 356, 423, 424,

470, 472, 473; pests of cereals in, 160, 257, 356, 422, 424, 425, 427, 428, 475, 476, 617; Citrus pests in, 354, 481; cotton pests 74, 158, 304, 422, 428–433, 665 hemp pests in, 159, 470, 475, 476, 477; pests of leguminous crops in, 75, 76, 257, 307, 355, 356, 425, 428, 432, 470, 471, 473, 474; miscellaneous pests in, 159, 429; orchard pests in, 74, 159, 302, 305-307, 352, 353, 354, 356, 432, 470, 477, 478, 479, 480, 481, 482, 483; pests of rubber-producing plants in, 72, 106, 618, 619; soil fauna of arable land in, 200, 368; mites in stored grain etc. in, 95–97, 159, 422; Codiosoma spadix in building timbers in, 354; new Coccids in, 48, 560; keys to Coccids on cultivated plants in, 304; Euxoa segetum in, 73, 160, 236, 422, 423, 424, 432, 472, 473; Heliothrips haemorrhoidalis in, 354, 358; Lamellicorns in, 158, 356, 429, 430, 431, 432; Loxostege sticticalis in, 94, 157, 257, 303, 354, 424, 472, 474; locusts in, 246, 357, 426; Plusia gamma in, 351, 354, 475; studies on wireworms in, 200, 424, 425, 429, 430, 432; beneficial insects and biological control in, 74, 94, 235, 236, 302, 303, 305, 308, 353, 354, 356; insects pollinating plants in, 184. (See Siberia and Asia, Central.)

Russian Far East, Ceuthorrhynchus sareptanus in, 351.

Russian Thistle (see Salsola).

rustica, Dexia.

Rye, pests of, in Czechoslovakia, 659; Cephus pygmaeus on, in Estonia, 462; pests of, in Germany, 51, 52, 717; Meromyza americana on, in Kansas, 122; Lema tristis on, in Manchuria, 767; Chlorops taeniopus on, in Poland, 518; Pentatomids on, in Switzerland, 91; type of injury to, by Pentatomids, 51.

Rye (Stored), new mite in, in Russian Union, 95.

Rye Grass (see Lolium spp.).

Rytigynia schumani, Dirphya usambica on, in Tanganyika, 113.

S.

Sabal spp., Homaledra sabalella on, in Florida, 22.

sabalella, Homaledra.

sabinianae, Platyphytoptus.

sabulosum, Opatrum.

Sacadodes pyralis, bionomics and control of, on cotton in Colombia, 403, 404.

saccharalis, Diatraea.

sacchari, Aphis; Lepidosaphes; Ripersia; Trionymus (Pseudococcus).

saccharicida, Perkinsiella.

saccharina, Lepisma; Tomaspis.

saccharivora, Ŝaccharosydne. saccharivorus, Ischnodemus.

Saccharosydne procerus, on rice in China, 498.

Saccharosydne saccharivora (on sugar-cane), in Florida, 326; in Porto Rico, 39.

Saccharum arundinaceum, Alissonotum impressicolle associated with, in Burma, 294.

Sacchiphantes (see Chermes). Safrol, in baits for Cydia, 135.

sagax, Pimpla.

Sagittaria variabilis, Listronotus latiusculus not found on, in New Jersey, **565**.

St. Kitts, summaries of plant quarantines in, 491, 692.

St. Lucia, Diaprepes spp. on Citrus etc. in, 402, 405, 448; miscellaneous pests in, 121; beneficial insects and biological control in, 121, 403; summaries of plant quarantines in, 134, 385, 491.

St. Vincent, thrips causing spotting of bananas in, 175; resting stage in *Platyedra gossypiella* in, 83; summaries of plant quarantines in, 134, 229, 491.

Saipan (see Japanese Mandated

Islands). Saissetia co

Saissetia coffeae (hemisphaerica), on ferns in Britain, 712; on tea in India, 149; on coffee in Indo-China, 185; on cassava in Madagascar, 454; measures against, 185, 712.

Saissetia nigra, parasite and control of, on cotton etc. in India, 55.

Saissetia oleae, in California, 574, 575; in Italy, 350; on Terminalia catappa in Haiti, 41; in Peru, 81; in Spain, 700; on Citrus, 81, 574, 575; on olives, 81, 350, 700; parasites and biological control of, 41, 81;

other measures against, 350, 574, 575.

Sakhalin, *Hylemyia flavopicta* on crucifers in, **770**; forest pests in, **452**, **516**; Pemphigini of, **496**.

salaminia, Eumaenas.

saliceti, Melasoma.

salicis, Chionaspis; Stilpnotia.

salignus, Lachnus (Pterochlorus).

Salix (see Willow).

Salpingogaster, parasite of Tomaspis in Brazil and W. Indies, 370.

Salsette Island, new Bruchid on Erythrina in, 292.

Salsola, relation of pests to, in U.S.A., 27, 323, 681.

Salt, in baits, **28, 224**; spraying with, against *Mycophila speyeri*, **656**; and lime-sulphur (see California Mixture).

saltator, Halticus.

saltuarius, Cryphalus.

Salvador, summaries of plant quarantines in, 134, 692.

Salvia, new Eriophyid on, in California, 173, 650.

Salvia sclarea, action of oil of, on insects and mites, 249, 250.

sambuci, Aphis.

Samia cecropia, parasite of, in Illinois, 330.

Samoa, pests of *Pritchardia pacifica* in, **702**; introduction of beneficial insects into, from Fiji, **212**. samsonovi, Eurytoma.

San José Scale (see Aspidiotus perniciosus).

sanborni, Macrosiphum.

Sand, seeds stored under, against pests, 8, 11, 204.

Sandwich Methods, for testing toxicity of insecticides, 135, 572.

sanguinea, Cycloneda.

sanguineus, Helopeltis.

sanguinipes, Pimpla (Ephialtes). sanguinolenta, Aceratagallia.

Santo Domingo (see Dominican Republic).

Saperda candida, measures against, on apple in U.S.A., 125, 269, 399; on Cotoneaster in Connecticut, 547.

Saperda carcharias, on ash and poplar in Russian Union, 429.

Saperda populnea, on ash in Russian Union. 429.

Saperda tridentata (see Eutetrapha). saperdae, Capitonius.

Saponin, as a wetting agent, 34, 36, 45, 206, 739.

sapporensis, Xylonomus. sapromyzarum, Histiostoma.

sarasini, Brontispa.

saratogensis, Phenacoccus.

Sarcophaga, parasite of locust in Angola, 688; parasite of Prodenia litura in Formosa, 207.

Sarcophaga atlanis, parasite of Dendrotettix quercus in U.S.A., **588.** Sarcophaga caridei, parasite of locust

in Argentina, 147.

Sarcophaga harpax, parasite of Lymantria dispar in Jugoslavia, 760.

Sarcophaga schützei, parasite of Lymantria dispar in Jugoslavia, 760.

Sarcophaga uliginosa, parasite of Lymantria dispar in Jugoslavia, 760.

sarcopis, Pronomeuta.

Sardine Oil, 702.

sareptanus, Ceuthorrhynchus.

sartor, Monochamus.

Sasakia charonda, on Celtis sinensis in Japan, 673.

Saskatchewan, grasshopper surveys in, 170, 440; miscellaneous pests in, 170.

Satin Moth (see Stilpnotia salicis). Satureia hortensis, experiment in infestation of, by insects and mites, 250.

Saturnia pavonia (see Eudia). satyriniformis, Melittia.

savastani, Capparimyia.
Sawdust, in baits: for cutworms,
540; for strawberry weevil,
270; for Orthoptera, 139, 326,

608, 686, 698; and Paris green, soil treatment with, against termites, 512.

Sawfly, Birch Leaf-mining (see Phyllotoma nemorata).

Sawfly, Black Wheat-stem (see Trachelus tabidus).

Sawfly, Dock (see Ametastegia gla-

brata). Sawfly, Pear (see Caliroa limacina). Sawfly, Pine (see Diprion pini and

D. sertifer).
Sawfly, Plum (see Hoplocampa

flava). Sawfly, Spruce (see Diprion polyto-

mum).
Sawfly, Wheat-stem (see Cephus

cinctus). saxeseni, Xyleborinus (see Xyleborus

xylographus). sayi, Chlorochroa; Pomphopoea.

scabiei, Pnyxia.

scabricollis, Lixus.

scabrisetus, Tetranychus (Eotetranychus).

scalaris, Stauroderus.

Scale, Brown Apricot (see Lecanium armeniacum).

Scale, Cherry (see Aspidiotus forbesi). Scale, Citrus Mussel or Purple (see Lepidosaphes beckii).

Scale, Citrus Red (see Aonidiella aurantii).

Scale, Date (see Parlatoria blan-chardi).

Scale, European Elm (see Gossyparia spuria). Scale, Florida Red (see Chrysom-

Scale, Florida Red (see Chrysom-phalus ficus).

Scale, Mango Shield (see Coccus mangiferae).

Scale, Obscure (see Chrysomphalus obscurus).

Scale, Olive (see Saissetia oleae). Scale, Pine Needle (see Chionaspis pinifoliae).

Scale, Plum (see Lecanium corni). Scale, San José (see Aspidiotus perniciosus).

Scale, White Peach (see Aulacaspis pentagona).

Scale, White Scurfy (see Chionaspis

furfura). Scale Insects, of N. Africa, 559; of Central Asia, 301; of Brazil, 367, 502; of Ceylon, 184; of France, 248, 303; of Italy, 119, 496; records of, in Japanese Mandated Islands, 48, 624; of Madeira, 560; of Portugal, 119; of Tatra Mountains, 772; of Ukraine, 560; keys to injurious species of, in Russian Union, 304; ants associated with, 202, 204, 411, 435, 588, 592, 606, 626, 642, 758; natural enemies of, 10, 18, 41, 43, 55, 65, 66, 81, 82, 101, 112, 114, 176, 204, 212, 214, 223, 243, 260, 268, 269, 296, 305, 312, 315, 319, 329, 337, 378, 410, 433, 456, 491, 501, 586, 587, 607, 611, 637, 641, 648, 659, 692, 715, 747, 753; classification and new species of, 48, 66, 184, 226, 248, 302, 303, 367, 469, 470, 496, 502, 559, 560, 607, 608, 696.

Scaphoideus aegyptiacus, on cotton in Sudan, **513**.

Scapteriscus vicinus, in Brazil, 223; in W. Indies, 40, 223; foodplants of, 40; natural enemies and biological control of, 40, 223; other measures against, 40.

scarabaeoides, Phloeotribus. Scarites, predacious on Heteronychus

licas in Natal, 411.

Scatopse fuscipes, on edible fungus in Japan and Korea, 438.

Scelio fulgidus, parasite of Chortoicetes terminifera in Australia, 450, 494; bionomics of, 450.

Scelio howardi (parasite of Nomadacris septemfasciata), in Angola, 688; in Nyasaland, 114.

Schedorhinotermes (see Rhinotermes). schistaceana, Eucosma (Olethreutes). Schistocerca americana, baits for, in Mississippi, 574.

Schistocerca cancellata, in Argentina, 147, 689; S. paranensis com-

pared with, 689.

Schistocerca gregaria, in N. Africa, **111**, **112**, **239**, **260**, **410**, 687; in S. W. Asia, 111, 112, 198, 199, 486, 683; in India, 199, 628, 678, 683, 686; in Mauretania, 755, 756; in Sudan, 6, 111, 198; outbreak centres of, 6, 198, 683, 687, 755, 756; colour variation and phases of, 6, 239, 410, 590, **628, 677, 678**; effects of temperature and humidity on, 147, 148. 410, 590, 628, 678, 756; watercontent of adults of, 184; migrations and breeding of, 111, 112, 628, 678, 683; other bionomics of, 628, 687; correlation of outbreaks of, with sun spots, 486: fungus and bacterial diseases of, 184, 591; mite attacking, 629; anatomy and histology of, 440, 624.

Schistocerca paranensis, 684; bionomics of, in Argentina, 146, 655, 689; measures against, damaging cotton etc. in U.S.A., 686; S. cancellata compared with, 689. Schistocerca vaga, in California, 19.

Schistoceros bimaculatus, boring in lead sheathing of cables in Greece, 232.

Schistoceros hamatus, in timber in Britain, 191; attacking twigs in U.S.A., 191.

Schizanthus, Heliothis phloxiphaga on, in Vancouver, 293.

Schizolachnus, producing honey-dew in Central Europe, **459**.

Schizomyia gennadii (see Eumarchalia).

Schizoneura (see Eriosoma).

Schizotrachelus cameratus, on tapped jelutong trees in Malaya, 109.

schlechtendali, Phyllocoptes.

Schlechtendalia chinensis (see Melaphis).

Schleichera trijuga (lac host-tree), pests of, in Ceylon, 359.

schoenobii, Shirakra; Tetrastichus. Schoenobius bipunctifer (incertellus), in China, 89, 109, 498; in India, 55; in Java, 89; measures against, on rice, 55, 498; parasites of, 89, 109; list of natural enemies of, 109.

schützei, Sarcophaga (Thyrsocnema).

schwartzi, Anuraphis.

Sciaphilus asperatus (muricatus), on strawberry in Britain, 383; Carabids predacious on, 384.

Sciara (on mushrooms), in U.S.A., 328; measures against, 328, 655. Sciara fenestralis, on mushrooms in Holland, 655.

Sciara pusilla, on mushrooms in

Holland, 655.

Scirpophaga (on sugar-cane), hand collection against, in China, 243; parasites of, in India, 243, 602.

Scirpophaga auriflua (see S. nivella). Scirpophaga intacta, new parasite of, in Java, 89.

Scirpophaga monostigma, on sugarcane in India, 241, 408, 499; susceptibility of varieties of sugarcane to, 499.

Scirpophaga nivella (on sugar-cane), in China, 243; in Formosa, 263; in India, 408, 499; in Loochoo Islands, 437; susceptibility of varieties of sugar-cane to, 263, 499; parasites of, 408.

Scirpophaga rhodoproctalis, parasites of, on sugar-cane in Madras, 408.

scirpophagae, Rhaconotus.

Scirtothrips dorsalis, measures against, on Capsicum in India, 55. Scirtothrips longipennis, measures against, on ornamental plants in Vancouver, 483.

Scirtothrips signipennis, bionomics and control of, on banana in Queensland, 211.

Sclerodermus, parasite of Minthea rugicollis in Malaya, 764.

Sclerodermus domesticus, parasite of Stromatium fulvum in Egypt, **592.** Scolia manilae (see Campsomeris

marginella modesta).

Scolothrips, predacious on Paratetranychus pilosus in Connecticut, 724; destroyed by tar oil sprays, 724.

Scolothrips sexmaculatus (predacious on Tetranychus telarius), in Bessarabia, 71; imported into Britain from California, 712.

Scolytids, of Korea, 207, 247; biological notes on Malayan species of, 295; classification and

new species of, **207**, **247**, **452**, **528**, **696**. (See Bark-beetles.)

scolvtivorus, Coeloides.

Scolytus intricatus, intercepted in Vancouver in crates from Britain, 483.

Scolytus multistriatus (on elms), in Czechoslovakia, 658; in Holland, 348, 656; in Russia, 74; in U.S.A., 325, 546, 551, 576; not found in New Hampshire, 20; feeding habits of, 551; associated with Dutch elm disease, 325, 547, 551, 656, 658; measures against, 348, 546, 551.

Scolytus pygmaeus, associated with Dutch elm disease in Czecho-

slovakia, 658.

Scolytus rugulosus, on peach in Indiana, 126.

Scolytus scolytus (on elms), in Czechoslovakia, 657; in Holland, 348, 656; associated with Dutch elm disease, 656, 657; measures against, 348.

Scopaeothrips unicolor, on Opuntia in Japan, 673.

scorbutica, Anasa.

Scorzonera latifolia, value of resin from, in banding adhesives, 480. Scorzonera tau-saghyz, pests of, in

Ukraine, **72, 106.**

scorzonerae, Xerophilaphis. Scotinophara coarctata, new parasite

of, in Malaya, 89.

Scotinophara lurida, on rice in China, 498.

Scots Pine (see Pinus sylvestris).

scrophulariae, Anthrenus.

sculderiana, Epiblema. sculptus, Oxytelus.

scutellata, Sturmia.

scutellator, Meteorus.

scutellatus, Gonipterus; Monochamus.

Scutellera perplexa (nobilis), on Citrus in Ceylon, 359.

Scutellista cyanea, parasite of Saissetia nigra in India, 55; imported into Peru from California against S. oleae, 81.

Scutigerella immaculata, in Argentina, 625; bionomics and control of, in U.S.A., 280, 488, 625.

Scymnillus aterrimus, predacious on Aspidiotus perniciosus in Idaho, 747.

Scymnus, predacious on Tetranychus telarius in Formosa, 674; predacious on Chermes cembrae in Germany, 763; introduced into

Mauritius from Java against Aspidiotus destructor, 204.

Scymnus minimus (see Stethorus punctillum).

Scyphophorus acupunctatus (Sisal Weevil), in Tanganvika, 113.

seabrai, Asterolecanium.

Seal-oil Soap (see Soap, Seal-oil). secalis, Trachae.

Seekay Waxes, 613. (See Napthalenes, Chlorinated.)

egetum, Anisoplia; Euxoa (Agrotis).

segmentata, Vanduzea.

Seius, predacious on Paratetranychus pilosus in Connecticut, 724. Selatosomus (see Corymbites).

selebensis, Brontispa (see B. longissima).

selenana, Ancylis.

Selenaspidus articulatus, on banana in Jamaica, 491; on Citrus in Peru, **81.**

Selenium, use of compounds containing, against pests, 65, 66, 131, 648; absorption of, by plants, 66, 131.

Selenothrips rubrocinctus, on mango in Jamaica, 490; on cacao in Trinidad, 10.

Selocide, 131.

Semanotus japonicus, bionomics of, on conifers in Japan, 515, 516. Semasia diniana (see Enarmonia).

semblidis, Trichogramma.

Semesan (see Hydroxymercurichlorophenol).

Semiadalia undecimnotata, and utilisation of, in nomics Central Asia, 238.

semiclavatus, Pteromalus.

semilaeve, Calosoma.

seminis, Cecidomyia (Itonida).

Semiotellus, parasite of Mayetiola destructor in Morocco, 340.

semipurpurana, Argyrotoxa. semistriatus, Microphanurus.

semivelutina, Hexamitodera. seneciella, Hylemyia (Pegohylemyia). Senecio jacobaea, insects introduced into Australia from England to

destroy, 450. seniculum, Apion.

senilis, Ceromasia.

Senotainia vigilans, parasite of Chlorochroa sayi in Montana, 44. septemfasciata, Nomadacris.

septempunctata, Chrysopa; Coccinella.

septendecim, Magicicada.

Serbian Spruce (see Picea omorika).

Serenoa serrulata, Homaledra sabalella on, in Florida, 22.

seriatus, Psallus; Stephanoderes. Serica castanea (see Hyposerica). Serica orientalis (see Aserica). sericata, Lucilia.

sericea, Blastothrix.

Sericothrips variabilis, on cotton in U.S.A., 277, 278, 644; measures against, 644.

Serinetha augur (see Leptocoris).

serinopa, Nephantis. serius, Eretmocerus.

serricorne, Lasioderma.

sertifer, Diprion (Lophyrus, Neodiprion).

Sesamia, parasite of, in Madras, 408.

Sesamia calamistis, on sugar-cane in Natal, 411.

Sesamia cretica, on sorghum and maize in Italy, 528, 712, 713:

measures against, 713.

Sesamia inferens (on sugar-cane), in China, 243; in Formosa, 263, 673; in Loochoo Islands, 264, 437; susceptibility of varieties of sugar-cane to, 264; on banana, 263; on cereals, 264; warmth accelerating development of hibernating, 54; measures against, 243, 673.

Sesamia pecki, sp. n., on Sorghum

in Br. Somaliland, 405.

Sesamia vuteria (on graminaceous plants), in Madagascar, 67; in Morocco, 500; in Spain, 695; bionomics of, 67, 500.

Sesamum, Aserica japonica on, in Japan, 605; Aphthona bimaculata on, in Nyasaland, 114; Ootheca bennigseni on, in Tanganyika, **642.**

Sesamum orientale, Antigastra catalaunalis on, in Cyprus, 108.

Sesbania, Coccids on, in Nyasaland, 114.

Setaria, Meromyza americana on, in Kansas, 122.

Setaria italica, Lema tristis on, in Manchuria, 767.

setariae, Carolinaia (Hysteroneura). setarius, Calomycterus.

Seventeen-year Cicada (see Magici-

cada septendecim). Sexava, on coconut in New Guinea Territory, 435, 495; parasites and biological control of, 435,

436; banding adhesives against, 495.

sexdens, Atta. sexdentatum, Sinoxylon. sexgibbosus, Prionomerus.

sexmaculatus, Scolothrips.

sexnotatus, Macrosteles (Cicada, Cicadula): Palpifer.

sexta, Protoparce.

seychellarum, Icerya.

seychellensis, Megachile; Odynerus.

Seychelles, coconut pests in, 433, 611: new Noctuid on mango in, 405; miscellaneous pests in, 433; beneficial insects and biological control in, 433, 611; summary of plant quarantines in, 134.

Shark Oil, 702.

sheldoni, Eriophyes.

Shelter Bag Moth (see Ochrogaster contraria).

shinjii, Cinara.

Shirakia schoenobii (dorsalis), parasite of Schoenobius bipunctifer in China, 109.

Shisham (see Dalbergia sissoo).

Shorea robusta, Brenthia dendronympha on, in India, 68.

Shorts (see Bran).

Sialis lutaria, parasite of, in Britain, 102.

Siam, new parasite of Attacus atlas in, 89; new thrips on rice in, 457: may-fly nymphs damaging timber under water in, 149.

Siberia, Ceuthorrhynchus sareptanus on flax in, 6, 351, 475, 601; Loxostege sticticalis in, locusts and grasshoppers in, 353, 357, 426; wireworms in, 425.

Siberian Grasshopper (see Aeropus sibiricus).

sibirica, Crioceris quatuordecimpunc-

sibiricus, Aeropus; Chermes (see C. cembrae).

Sicily, pests of caper in, 92; Hybalus cornifrons on cereals in, 757; miscellaneous pests in, 70, 93, 119.

Sida, Earias fabia on, in Java, 766. Sida paniculata, Dysdercus mimus on, in Peru, 533.

Sierra Leone, miscellaneous pests in, 118, 239; parasites of Trypetids in, 492.

signatus, Anthonomus.

signifer, Phassus.

signipennis, Scirtothrips.

Signiphora conjugalis, parasite of Pseudococcus citri in Spain, 701.

Signiphora dipterophaga (see Thy-

Silica, particle size affecting insecticidal action of, 252.

Silicic Acid, dusting with, against pests of stored cereals and peas, 644.

Silk, list of Lepidoptera and spiders producing, 68; insects damaging, 514, 515, 568, 588.

Silkworm (see Bombyx mori).

Silvanus gemellatus, methyl bromide against, in stored cereals in U.S.A., 746.

silvatica, Masicera.

silvaticus, Tetrastichus. Silverfish (see Lepisma saccharina and Thermobia domestica).

silvestrii, Hyperaspis; Micrococcus. simbleuta, Ptochoryctis.

similis, Micrococcus.

simmondsi, Brontispa (see B. longissima).

simonyi, Lygus.

simplex, Agromyza (Melanagromyza); Anabrus: Chilo: Hemithea costipunctata; Pyroderces; Taeniothrips.

simplicicornis, Calotermes (Kalotermes, Paraneotermes).

simplicipes, Glypta.

Sind, Schistocerca gregaria in, 628, 683.

sinensis, Paratenodera. singularis, Otiorrhynchus.

sinicus, Adoretus.

Sinodendron cylindricum, on cherry in Britain, 612.

Sinoxylon, sodium fluosilicate ineffective against, in timber in Sudan, **514.**

Sinoxylon conigerum, in stored cassava tubers in Madagascar, 455.

Sinoxylon sexdentatum, boring in lead in Greece, 232.

Sipha flava (on sugar-cane), in Br. Guiana, 369; natural enemies of, in Porto Rico, 39, 40, 289; ant associated with, 289; susceptibility of varieties of sugar-cane to, 39; not transmitting mosaic, 40.

Siphocoryne avenae, auct. (see Rhopalosiphum prunifoliae).

Siphocoryne indobrassicae (see Rhopalosiphum pseudobrassicae). siphonophorae, Pachyneuron.

Sirex cyaneus, fungus associated with, in timber in Britain, 191.

Sirex gigas, fungus associated with. in timber in Britain, 191, 632.

Sirex noctilio, establishment of Rhyssa persuasoria against, on pines in New Zealand, 58.

Sisal, pests of, in Tanganyika, 113.

Sisal Weevil (see Scyphophorus acubunctatus).

Siteroptes, on couch grass in Queensland, 494.

Sitka Spruce (see Picea sitchensis). Sitodiplosis mosellana (on wheat), in Br. Columbia, 60; in Germany, 619; in Japan, 670; in Sweden, 230, 231, 620; bionomics of, 230, 670; natural enemies of, 231, 620, 670.

Sitodrepa panicea, Opilo domesticus predacious on, in Germany, 662: in stored ginger in Jamaica, 491: damaging books, 706; oviposition

of, 164, 165.

Sitona, measures against, on lucerne in Russia, 76.

Sitona callosa, on lucerne in Russia. 76.

Sitona crinita, food-plants of, in Russia, 76, 432.

Sitona gressoria, on lupins in Germany, 248.

Sitona grisea, on lupins in Germany,

Sitona inops, on lucerne in Russia,

Sitona lineata, on lucerne in Czechoslovakia, 657; on peas and clover in Vancouver, 293.

Sitona longula, on lucerne in Russia,

Sitophilus (see Calandra).

Sitotroga cerealella, on field maize, 643; in stored cereals, 116, 218, 389, 414, 668; bionomics of, measures against, 116, 668; 218; tests of fumigants against, 389, 414; use of eggs of, for rearing Trichogramma, 73, 103, 235, 356, 545, 725; factors concerned in finding of eggs of, by T. evanescens, 150.

sjaelandicus, Corymbites.

sjöstedti, Athalia.

Sloe, Coccids on, in Germany and Russia, 251, 481.

smaragdina, Oecophylla. smenus, Glyptomorpha.

smithi, Lachnosterna (Phytalus). smithiae, Pterocomma (Clavigerus).

Smynthurus viridis, in Australia, 84, 449; natural enemy and control of, on leguminous plants,

Soap, spraying with, 62, 149, 169, 246, 587, 663; in mixed sprays, 21, 26, 32, 36, 45, 46, 62, 72, 88, 100, 114, 131, 150, 191, 237, 241, 260, 270, 309, 335, 350, 355, 361, 376, 404, 412, 421, 422,

431, 433, 467, 484, 522, 536, 542, 549, 559, 563, 594, 604, 615, 646, 653, 663, 682, 690, 701, 704, 707, 710, 711, 713, 716, 733, 755, 762, 768, 771; effects of, on formation of soluble arsenic from lead arsenate, 21; contact performance of types of, 45; emulsifying oils, 82, 167, 185, 187, 246, 501, 653; formulae containing, 36, 46, 65, 82, 100, 131, 149, 167, 187, 191, 246, 260, 270, 335, 350, 355, 404, 412, 431, 467, 501, 559, 563, 564, 604, 615, 646, 653, 701, 711, 755, 762, 771; plants injured by, 704.

Soap, Coconut-oil, as a spreader for derris and cubé, 646; contact

performance of, 45.

Soap, Cotton-oil, by-product of, against cotton red spider, 471.

Soap, Fish-oil, in mixed sprays, 148, 291, 550, 646, 750; as an emulsifier, 398, 484; formulae containing, 398, 550, 646.

Soap, Linseed-oil, as an emulsifier,

484.

Soap, Seal-oil, as an emulsifier, 290. Soap-bark Tree (see Quillaia saponaria). socius, Zelus.

Soda (see Sodium Carbonate).

Soda, Caustic (see Sodium Hydroxide).

Sodium Arsenate, in baits for locusts, 698; in bait-spray for Dacus oleae, 700.

Sodium Arsenite, in baits, 108, 139, 170, 182, 219, 410, 441, 540, 574, 682, 685, 686, 698; unsatisfactory in baits for Cydia, 506; in bait-sprays, 71, 108, 179, 619; dusting with, 609,682; aeroplane dusting with, 681, 682; spraying with, 152, 169, 682; trees and timber treated with, against Coleoptera, 453, 514; formulae containing, 108, 152, 169, 170, 182, 410, 453, 514, 574, 619, 681, 682, 685, 686, 698; question of danger of, to cattle, 609, 682; effects of, on parasites of Antestia, 179.

Sodium Benzoate, in baits for

Cydia, 137, 506. Sodium Carbonate (Soda), insecti-

cide spray formulae containing, 221, 245, 246, 335, 701, 713.

Sodium Chloride (see Salt).

Sodium Cyanide, fumigants containing, against leaf-cutting ants, 533; against Vesperus, 699; against wireworms, 383, 630, 631; and magnesium sulphate, 630, 631; and sulphuric acid, for generating hydrocyanic acid gas (q.v.), 18, 176, 306, 542, 593, 702.

Sodium Ethyl Xanthate, unsatisfactory as a soil fumigant against *Tipula aino*, **54**.

Sodium Fluoaluminate (see Cryolite).

Sodium Fluoride, in baits, 51, 160, 224, 387, 713; in bait-sprays, 50, 51; mixed with stored seeds against weevils, 642; for treating timber against Lyctus, 190, 514; formulae containing, 50, 160, 190, 224, 387, 514, 642, 713; causing foliage injury, 51; other fluorine compounds compared with, 387.

Sodium Fluosilicate, in baits, 37, 75, 153, 202, 270, 387, 393, 509, 520, 566, 608, 609, 639, 686, 699; in bait-sprays, 108, 267; dusting with, 145, 226, 239, 431, 433, 477, 686, 691; carriers for, 226, 433, 572; spraying with, 145, 433; mixed with stored seeds against weevils, 642; treatment of timber with, against pests, 190, 514; tests of toxicity of, to insects, 205, 226, 387, 572, 608, 609; castor injured by, 205; formulae containing, 37, 75, 153, 190, 202, 226, 267, 387, 393, 477, 514, 566, 609, 639, 642, 686, 699; compared with sodium fluoride, 387.

Sodium Hydroxide (Caustic Soda), as a repellent for bark-eating caterpillars, 602; mixed spray formulae containing, 246, 412, 755; wetting and spreading properties of mixtures of, with various fatty acids, 320.

Sodium Lauryl Sulphate, as a contact spray, 286; as a spreader for nicotine-bentonite, 732, 733; as an emulsifier for oils, 739; Gardinol and Lethalate containing, 45, 615.

Sodium Nitrate, effect of spraying with, on plum, 62.

Sodium Oleate, tests of, in baits for Cydia, 135; contact performance of, 45; other spreaders for nicotine thiocyanate compared with, 704; affecting oil deposits of sprays, 280.

Sodium Oleyl Sulphate, use of, in dusts against pea Aphid, 327, 740: as a spreader, 554.

Sodium Phosphate, increasing toxicity of lead arsenate to larvae of *Popillia*, 324.

Sodium Selenate, 66.

Sodium Silicate, uses of, for removing spray residues from apples, 318, 402, 511; and lead arsenate, causing foliage injury to apple, 402.

Sodium Sulphoricinate, emulsifier for pyrethrum containing, 412.

Soil, fauna of, in Ukraine, 200, 368; technique of sampling, for wireworms, 630; effects of character of: on wireworms, 72, 200, 201, 356, 424, 425, 426, 429, 430, 494, 569, 630; on other insects, 40, 72, 83, 88, 157, 158, 193, 194, 200, 201, 210, 220, 223, 230, 251, 267, 276, 294, 356, 366, 367, 392, 393, 399, 420, 422, 423, 429, 430, 434, 442, 494, 626, 634, 751, 757.

sokolowi, Telenomus.

solana, Empoasca.

solani, Agromyza (Liviomyza); Macrosiphum; Phenacoccus.

solanifolii, Macrosiphum (Illinoia). Solanum, Anthonomus eugenii on, in U.S.A., **64, 328;** effect of species of, on predacious Pentatomids, **13.**

Solanum arundo, Coccids on, in Italian Somaliland, 757.

Solanum carolinense, Leptinotarsa decembineata fed on, 13, 14.

Solanum dulcamara, Leptinotarsa decemlineata on, 13, 14, 415; likely to facilitate spread of L. decemlineata in Europe, 415.

Solanum melongena (see Egg-plant). Solanum tuberosum (see Potato).

Solar Heat, reactions of insects to, 6, 604; against pests in stored products, 8, 118, 408. Solar Oil, 472.

Solenobia pineti, on Pinus cembra in Austria, 664.

Solenopsis fugax, in Russian Union, 107, 305; fostering Aphids, 107; predactious on Aspidiotus perniciosus, 305.

Solenopsis geminata var. maniosa, bionomics and control of, in California, 588.

Solenopsis geminata rufa, erroneously recorded as damaging rubber insulation in Netherlands Indies, 262.

Solidago, weevils on, in Connecticut, 15; Epiblema spp. on, in Ontario, 510.

solidus, Platypus.

solitarius, Apanteles; Spilocryptus. Solomon Islands, Pyroderces paroditis on coconut in, 497; Liothrips urichi introduced from Fiji into, against Clidemia hirta, 703.

solstitialis, Amphimallus (Amphi-

mallon, Rhizotrogus).

Solubea pugnax, food-plants of in Louisiana, 218.

Somaliland, British, new Noctuid

on Sorghum in, **405.** Somaliland, Italian (see Africa,

Italian East).
somniaria, Ellopia.

sonchi, Ensina.

Sonchus arvensis, new Coccid on, in Java, 607; Trypetids on, in Ukraine, 618, 619.

Sonchus oleraceus, destruction of, against Hercothrips fasciatus in

California, 19, 129.

Sonronius, gen. n., key to N. American species of, 78.

Sooty Moulds, insects associated with, 454, 587, 753, 757.

Sophora japonica, Aphis gossypii on, in China, 245.

sophorae, Brassolis.

Sophronia humerella, measures against, on lavender in France, 301.

Sorbus, identity of woolly Aphid on,

in Maine, 484.

Sorbus americana, mite on, in Br. Columbia, 171.

Sorbus aucuparia, mite on, in Br. Columbia, 171; Chionaspis salicis on, in Finland, 201; Otiorrhynchus hirticornis on, in Germany, 344; Argyresthia conjugella on, in Sweden and Switzerland, 299; Rhagoletis cerasi not observed in fruits of, 156.

sordidus, Cosmopolites.

sorghi, Aphis.

sorghiella, Celama.

sorghiellae, Apanteles.

Sorghum, Toxoptera graminum on, in Argentina, 289, 290; pests of, in India, 55, 408, 531; pests of, in Italy, 528; Sesamia vuteria on, in Morocco, 500, 501; pests of, in Porto Rico, 39; new Noctuid on, in Br. Somaliland, 405; pests of, in Sudan, 513; pests of, in U.S.A., 63, 144, 560;

Peregrinus maidis and virus diseases of, 55, 531; resistance of varieties of, to Blissus leucopterus, 63; (stored), treatment of, against weevils in Tanganyika, 642; (broom corn), legislation dealing with treatment of, against Pyrausta nubilalis in U.S.A., 750. Sorghum Molasses (see Molasses).

Sorghum Webworm (see Celama

sorghiella).

sovor, Diabrotica.

Sour-sop (see Anona muricata).

Soviet Green, spraying with, against

Cydia pomonella, 479.

Sow Thistle (see Sonchus oleraceus). Soy Bean, pests of, in Japan, 208, 264, 310, 516, 605, 673; Cydia glycinivorella on, in Manchuria, 54, 310, 452, 606; Melolontha melolontha on, in Rumania, 721; Tetranychus on, in Ukraine, 257; not attacked by Blissus leucopterus in U.S.A., 78; calcium arsenate in soil injurious to, 328. Soy-bean Flour or Meal, uses of, in

sprays, **125**, **489**, **490**, **542**. Soy-bean Oil (see Oil, Soy-bean). Soy-bean Pod Borer (see *Cydia*

glycinivorella).

soya, Profeltiella (Contarinia). Soxhlet Method, 258, 412, 739.

spadix, Codiosoma.

Spain, miscellaneous pests in, 695, 697-701; beneficial insects and biological control in, 695, 696, 697, 698, 701; dosage tables for fumigating Citrus with HCN in, 702; Lepidosaphes beckii originally imported into Palestine from, 636.

Spalangia (parasite of Dacus ferrugineus), introduced into Ceylon from Malaya, 259.

Spalangia afra, parasite of Trypetids in Zanzibar, 492.

Spaniopterus crucifer, attempted introduction of, into Mauritius against Aspidiotus destructor, 176.

Sparganothis sulfureana, bionomics and control of, on cranberry in New Jersey, **562**, **563**.

sparsus, Tanymecus.

Spathius, parasite of Hylesinus aculeatus in New Jersey, 394.

Spathius canadensis, 394; bionomics of, in Connecticut, 14.

speciosa, Diabrotica; Orphulella.

spectabilis, Dendrolimus.

Specularius erythrinae, gen et sp. n., distribution of, in seeds of Erythrina spp., 292.

L

speiseri, Acanthoplus. speyeri, Mycophila.

Sphaerakis mayri, parasite of Bruchus brachialis in N. Carolina, 24, 580.

sphaerogaster, Caloglyphus (Isoglyphus).

Sphegigaster, parasite of Phytomyza ilicicola in Maryland, 541.

Sphenophorus minimus, on wheat in Ohio, 130.

Sphenoptera gossypii (on cotton), parasites of, in India, 89, 204; in Nigeria, 667; measures against, in Sudan, 512.

Sphex torridus, predacious on sugarcane borers in Madagascar, 67.

Sphingomorpha pudens, damaging fruit in Sierra Leone, 239.

spicatum, Acrosternum.

Spiders, producing silk, **68**; destroying insects, **87**, **309**, **557**.

Spilocryptus solitarius, parasite of Aglaope infausta in Spain, 697. Spilonota (see Eucosma).

spilotus, Anthonomus.

Spindle Oil, 478, 481.

spinipennis, Agrilus.

spinipes, Maleuterpes; Tarsonemus. spinolae, Lygus.

spinosus, Dectes.

Spintherus linearis, parasite of Apion spp. in Russia, 308.

spinulifera, Eriophyes. spiraecola, Aphis.

spirifex, Tarsonemus.

spissicornis, Coleophora. splendana, Cydia (Carpocapsa).

Spodoptera exempta (see Laphygma). Spodoptera mauritia, Laphygma exempta recorded as, in Hawaii, 692; on cereals and grasses in India, 240, 360; in Malaya, 89;

India, 240, 360; in Malaya, 89; parasites of, 89, 241; measures against, 360.

Spondias lutea, new weevil on, in Brazil, 405.

spondias, Hemiliopsis.

Sprays, types of equipment for applying, 134, 216, 343, 387, 440, 483, 536, 643, 682; question of volume delivery of, 46; use of aeroplanes for applying, 19, 748; factors affecting retention of deposits of, 321, 370; dusts compared with, 443, 467, 522, 564, 647, 726, 734; programmes for, in orchards in Illinois, 652. spretus, Corymbites (Selatosomus).

Spruce (*Picea*), pests of, in Canada, **77**, **103**, **215**, **293**, **330**, **487**, **534**, **538**; pests of, in Czechoslovakia,

215, 657, 715; bark-beetles on, in Estonia, 462; Chrysobothris chrysostigma on, in Finland, 518; pests of, in Germany, 154, 189, 344, 345, 664, 762; Chermes spp. on, in Japan, 437, 438; pests of, in Sweden, 634, 635; C. abietis on, in Switzerland, 459; pests of, in U.S.A., 80, 270, 293, 321, 330, 369, 441, 487; unsuitable foodplant for Dendrolimus spectabilis, 672; oil emulsions causing injury to, 270; effects of soil treatment with lead arsenate on, 441; pests of timber of, 91, 354.

Spruce, Black (see *Picea mariana*). Spruce, Oriental (see *Picea orien-*

talis).

Spruce, Red (see *Picea rubra*).
Spruce, Serbian (see *Picea omorika*).

Spruce, Sitka (see *Picea sitchensis*). Spruce Budworm (see *Harmologa fumiferana*).

Spruce Gall Aphis (see Chermes abietis).

Spruce Sawfly (see Diprion polytomum).

spuria, Gossyparia.

sputator, Agriotes.

squamosus, Hypomeces.

Squash, Camptocladius macleayi on, in New South Wales, 209; pests of, in U.S.A., 17, 647, 726, 739; used for rearing Encarsia formosa, 623.

Squash Bug (see Anasa tristis).
Squash Vine Borer (see Melittia

satyriniformis).

S.S.S. Spreader, in sprays, **546**, **547**. SS₃, as a spreader for sprays, **548**. stabulans, Lydella; Muscina.

Stachytarpheta jamaicensis, planted to favour Scoliids in Mauritius, 317.

Stantonia carpocapsae, sp. n., parasite of Cydia pomonella in Kentucky, 330.

Stantonia lamprosemae, sp. n., parasite of Lamprosema indicata in Cuba, 330; in Porto Rico, 330.

Staphylococcus causing disease in Lepidiota caudata in Queensland, 267.

Starch, content of, in ash bark in relation to *Hylesinus fraxini*, **631**; content of, in timber and its depletion in relation to Lyctids, **190**, **635**, **707**, **763**, **764**; in bait for Lepismids, **37**.

Starlings, destroying noxious insects 180, 255.

Stathmopoda melanochra, introduced into New Zealand against Erio-

coccus coriaceus, 753.

Statistical Methods, application of, to entomological investigations, 94, 142, 165, 196, 357; book on, 688.

Stauroderus scalaris, ploughing to destroy egg-pods of, in Siberia, 353.

Stauropus alternus, on tea in India, 149.

Steam, uses of, against pests, 326, 477, 539, 626, 750.

Steel Brushes, against locusts, 685. steinitzi, Aphanogmus.

stellata, Lyda: Trypanea.

Stemona tuberosa, preparation and insecticidal value of extracts of. 206

Stenaraeoides octocinctus, parasite of Dendrolimus spectabilis in Japan,

Stenobracon (see Glyptomorpha).

Stenoma catenifer, on avocado in Brazil, 502.

Stenomalus, parasite of Geomyza tripunctata in Britain, 382.

stenoperitrematus, Paratetranychus (Eurytetranychus).

stenoptera, Meliana flammea.

Stephanitis pyri, measures against, on apple and pear in Cyprus, 260. Stephanitis rhododendri, on rhodo-

dendron in Britain, 350.

Stephanoderes hampei (on coffee), S. seriatus mistaken for, in Brazil, 319; measures against, in Ceylon, 359, 360; failure to establish Prorops nasuta against, in Netherlands Indies, 337; probably unable to breed in dry market coffee, 359.

Stephanoderes seriatus, bionomics of, on coffee in Brazil, 319.

Sterculia spp., Dysdercus spp. on, in Africa, 365, 374, 512, 642; Lamiid on, in Nyasaland, 114. sterculiae, Tepperia.

Stereonychus fraxini, on Fraxinus excelsior in Czechoslovakia, 658.

Stereum sanguinolentum (and fungus resembling), Sirex spp. associated with, in Britain, 632.

sternata, Tiphia.

Stethorus punctillum, predacious on mites in Holland, 465; killed by tar distillate sprays, 465.

Stethorus punctum, predacious on Paratetranychus pilosus in Connecticut, 724.

Stethorus rotundatus, predacious on Tetranychus telarius in Formosa, 674.

Sthenias grisator, habits and control of, on Erythrina in India, 149.

Stiamus brachyurus, on sugar-cane in Madagascar, 67.

Stiboscopus thoracicus, parasite of Apanteles americanus in Cuba, 171.

sticticalis, Loxostege.

sticticraspis, Diatraea (Argyria).

Stictocephala festina, on potato in Hawaii, 214; on cotton and narcissus in U.S.A., 279, 288.

Stictodiplosis moringae, on Moringa in India, 55, 603; bionomics and control of, 603.

Stigmaeus floridanus, on pineapple in Formosa, 674.

stigmaterus, Microdus (Bassus). Stigmatrachelus concinnus, on cassava in Madagascar, 455.

Stigmus americanus, in New Jersey, 394.

Stilpnotia salicis, in Austria, 589; in Belgium, Germany and Holland, 576; in Hungary, 576, 589; in Russia, 432; in U.S.A., 16, 17, 576, 589; on poplar and willow, 15, 17, 432, 589; parasites and biological control of, 576, 589, 590; other measures against, 432.

stipella, Borkhausenia.

stironotus, Pseudanogmus.

Stobaera tricarinata, on narcissus in U.S.A., 288.

platani, Stomacoccus measures against, on plane in California, 18. Stomaphis ulmicola, sp. n.,

Ulmus propingua in Japan, 452. Stomatoceras imbili, sp. n., parasite of Hypsipyla robusta in Queensland, **502.**

Stomatoceras sulcatiscutellum, parasite of Nephantis serinopa in Cochin, 452.

Stomatodexia diadema (parasite of Diatraea), parasitised by Thysanus dipterophaga in Br. Guiana, 491.

Stomatorrhina lunata (egg parasite of Nomadacris septemfasciata), in Angola, 688; in Nyasaland, 114. Stomatothrips flavus, on cotton in

S. Carolina, 278.

Stored Products, keys to pests of, in Central Europe, 196; books on control and breeding of pests of, **347, 468.**

Storks, doubtful danger of poisoned locusts to, 180.

Strategus, measures against, on coconut in St. Lucia, 121.

Straw, barriers of, for Cirphis

unipuncta, 723. Strawberry, pests of, in Britain, 103, 383, 384, 612, 613; Anthonomus rubi on, in Estonia, 462; Harpalus rufipes on, in Finland, 201: pests of, in Germany, 194, 521, 660; pests of, in Japan, 604, 771; Lymantria dispar on, in Jugoslavia, 759; Haltica oleracea on, in Poland, 255; Coleopterous pests of, in Tasmania, 208; pests of, in U.S.A., 18, 23, 45, 270, 288, 291, 324, 328, 399, 443, 545, 735, 736; regulations dealing with treatment and movement of, against Popillia japonica in U.S.A., 384; susceptibility of varieties of, to Anthonomus rubi, 383; Dacus cucurbitae experimentally developing on, 668; Capitophorus spp. and virus diseases of, 104, 291, 383.

Strawberry (Fruit), problem of insecticide residues on, 383, 735; HCN unsuitable as a fumigant

for, **307.**

Strawberry Aphis (see Capitophorus fragariae).

Strawberry Crown Borer (see Tylo-

derma fragariae). Strawberry Crown Moth (see Aegeria

bibionipennis).

Strawberry Leaf-roller (see Ancylis comptana).

Strawberry Root Weevil (see Otior-rhynchus ovatus).

Strawberry Weevil (see Anthonomus bisignifer).

strenuana, Epiblema.

Stretchia plusiaeformis, bionomics and control of, on gooseberry in Colorado, 281.

striata, Anastrepha.

striatalis, Diatraea (see D. venosata).

striativentris, Opius. striatulus, Euscelis.

striatus, Acromyrmex (Atta).

strobi, Chermes (Pineus); Phlepsius. strobilella, Cydia (Laspeyresia).

strobilohius, Chermes (Cnaphalodes).

Stromatium fulvum, measures against, in timber and furniture in Egypt, **591**; parasite of, **592**. strumarius, Pachytychius.

Sturmia, parasite of Eterusia virescens in India, 149.

Sturmia bimaculata (see S. in-conspicua).

Sturmia elzneri, parasite of Lepidoderma albohirtum in Queensland, 151.

Sturmia inconspicua, parasite of Diprion pini in Holland, 348; parasite of Spodoptera mauritia in India, 241; parasite of Lymantria dispar in Jugoslavia, 760.

Sturmia inconspicuella, hosts of, in

India, **765.**

Sturmia inconspicuoides (parasite of Noctuids), in India, 55; in Queensland, 151.

Sturmia scutellata, parasite of Celerio euphorbiae in Czechoslovakia, 529; parasite of Lymantria dispar in Jugoslavia, 760.

Styrax benzoin, incidence of ambrosia beetles on poisoned and girdled,

in Malaya, 453.

Subcoccinella vigintiquatuorpunctata, tests of adhesiveness of pyrethrum dusts on, 197.

subferruginea, Hypocassida. subnigrescens, Mermis. substriatus, Blapstinus.

Subterranean Clover (see Trifolium subterraneum).

succincta, Epicampocera; Patanga.

succineus, Longitarsus.

Succingulum mistum, parasite of Euxoa segetum in S. Rhodesia, 90. Succinic Acid, esters of, attracting

Cydia pomonella, 137.

Sudan, Anglo-Egyptian, pests of cotton etc. in, 511-513; miscellaneous pests in, 513, 514; locusts in, 6, 111, 112, 198, 679.

Sudan Cotton Bollworm (see Diparopsis castanea).

sudanensis, Hercothrips. suffodiens, Platypus.

Sugar, new mite in, in Formosa, 436; not eaten by Gryllulus domesticus, 38; types of, used for feeding insects, 424, 561; in baits, 37, 51, 57, 125, 137, 142, 182, 202, 224, 506, 520, 526, 544, 639, 672; in bait-sprays, 50, 51, 267, 325, 376, 526; use of, in sprays against thrips, 274, 398, 399, 489, 552, 553. (See Syrup.)

Sugar Pine (see Pinus lambertiana). Sugar-cane, Diatraea saccharalis on, in Argentina, 390; pests of, in Australia, 151, 584; pests of, in Brazil, 82, 502; Alissonotum impressicolle on, in Burma, 294; Pyrilla on, in Ceylon, 8; pests of, in China, 243, 498; pests of, in Formosa, 263, 294, 673, 768,

INDEX, 937

769; pests of, in Br. Guiana, 221, 369, 491; pests of, in Hawaii, 454, 493, 585, 589; pests of, in India, 8, 55, 241, 243, 261, 294, 408, 498, 532, 602, 628; Podothrips lucasseni on, in Java, 457; pests of, in Loochoo Islands, 264, 437, 516; pests of, in Madagascar, 66, 67; pests of, in Mauritius, 176, 316, 317, 454; pests of, in Natal, 411: Diatraea saccharalis on, in Peru, 390: Chionaspis tegalensis on, in Réunion, 501; pests of, in U.S.A., 28, 137, 326, 390, 577; pests of, in W. Indies, 10, 39, 40, 43, 82, 121, 221, 222, 223, 269, 288, 289, 317, 370, 404, 448, 490, 752; susceptibility of varieties of, to pests, 10, 39, 151, 221, 263, 264, 294, 317, 490, 498, 501, 769; locusts and grasshoppers damaging, 67, 411, 584, 769; termites attacking, 243, 261, 411; insects and diseases of, 39, 40, 386, 411, 454, 577; mite and Jassid possibly associated with stem galls on, 585; carbon bisulphide causing injury to, 437.

Sugar-cane Froghopper (see Tomas-

pis saccharina).

Sugar-cane Mealybug (see Pseudo-coccus boninsis).

Sugar-cane Wireworm (see Cono-

derus exsul).

Sugitania maculifera, bionomics of, on Camellia in Japan, 311.

suis, Haematopinus. sulcata, Navomorpha.

sulcatus, Navomorpna.
sulcatus, Otiorrhynchus (Brachyr-

rhinus). sulciventris, Rhoecocoris. sulfureana, Sparganothis.

Sulphite Lye, as an adhesive for cryolite sprays, 25; as an emulsi-

fier, 371, 506, 614, 708, 738.
Sulphur, 651; against Jassids, 140, 558, 565, 609, 610; apparently not reducing parasitism of Typhlocyba pomaria, 558; against Lygus pratensis, 279, 565, 710; against Psallus seriatus, 136, 277; against Malacosoma americana, 542; ineffective against Heliothis armigera, 327; against mites, 15, 20, 72, 108, 114, 160, 246, 298, 334, 335, 471, 488, 521, 648, 690, 724, 748; against thrips, 55, 222, 355, 644, 727; against papaya whitefly, 727; dusting with, 15, 20, 55, 72, 108, 136, 140, 246, 327, 355, 542, 565,

609, 644, 690, 710, 727; aeroplane dusting with, 748; determination of particle size of dusts of, 739; in mixed dusts, 65, 125, 129, 142, 276, 277, 279, 327, 334, 350, 397, 443, 542, 544, 556, 565, 609, 644, 645, 726, 734, 748, 752, 757; uses and tests of, as a fumigant, 37, 188, 190, 579, 633, 714: and arsenic, fumigants against ants containing, 533: spraying with, 15, 16, 125, 298, 521, 558, 690, 724; wetting agents and adhesives for, 542, 750; in mixed sprays, 15, 16, 25, 26, 72, 79, 114, 149, 150, 222, 334, 443, 489, 490, 566, 648, 724, 725, 727, 728, 732, 733; nicotine and anabasine sulphates adsorbed on, 479; formulae containing, 25, 125, 140, 142, 149, 276, 277, 298, 327, 334, 335, 443, 489, 490, 544, 645, 724, 728, 734, 752, 757; soil treatment with, ineffective against pests, 327, 461, 645; causing injury to plants, 16, 26, 79, 125, 216, 556; other insecticides compared with, 279, 471, 565, 645.

Sulphuric Acid, and cyanides, for generating hydrocyanic acid gas (q.v.), 53, 176, 306, 542.

Sulphurous Slag, generating hydrogen sulphide (q.v.), 160.

Sumach Gall Aphis (see Melaphis chinensis).

sumatrana, Brontispa (see Plesispa reichei).

Sumatrol, toxicity of, to insects, 616, 632; toxicarol precursor containing, 616, 631.

Sunflower, Astylus atromaculatus on, in Argentina, 705; new mite on, in Central Asia, 530; Heliothis armigera on, in N. Rhodesia, 366; grain mites attacking stored seeds of, 96.

Sunflower Seed Oil (see Oil, Sunflower Seed).

Sun Spots, correlation of insect outbreaks with, 486, 604.

Sunn Hemp (see Crotalaria juncea).

superbiens, Thalessa.

Superphosphate, and carbolic powder, dusting with, against mites, **85**; and lime, soil treatment with, against wireworms, **631**.

superstitiosus, Dysdercus.

surinamensis, Oryzaephilus; Pyenoscelus.

suspensa, Anastrepha. susurrans, Zenillia (Carcelia). suturalis, Eleodes; Lochmaea.

Swede, pests of, in Britain, 296, 709. Sweden, forest pests in, 299, 611, 634; list of insects found in fruit-tree bands in, 689; Contarinia medicaginis on lucerne in, 232; wheat gall-midges in, 154, 230, 231, 620; pests in building timbers in, 155, 440, 624; beneficial insects in, 155, 231, 232; summaries of plant quarantines in, 134, 692.

Swedish Fly (see Oscinella frit). Sweet Clover, pests of, in U.S.A.,

127, 287.

Sweet Pea, Macrosiphum onobrychis transmitting vein-mosaic of clover to, in New Jersey, 12; Heliothis phloxiphaga on, in Vancouver, 293.

Sweet Potato, pests of, in Ceylon, 358; pests of, in Formosa, 207, 674; Aserica japonica on, in Japan, 605; Pseudococcus kenyae on, in Kenya, 592; Cylas formicarius on, in Loochoo Islands, 669; pests of, in Louisiana, 286; new weevil on, in S. Rhodesia, 106; Helopeltis bergrothi on, in Sudan, 513; pests of, in W. Indies, 145, 222, 223, 269, 446; susceptibility of types of, to Euscepes batatae, 447; tubers of, as baits for C. formicarius, 269.

Sweet Potato Weevil (see Cylas

formicarius).

Switzerland, Pentatomids on cereals in, 91; forest pests in, 164, 299, 459; Leptinotarsa decembineata in, 70, 91, 120; miscellaneous pests in, 156, 689; orchard pests in, 99, 156, 163, 299, 300, 715; vine pests in, 299; beneficial insects in, 92, 100, 156; standardisation of insecticides etc. in, 155; legislation against introduction of L. decembineata into Germany from, 91.

Sword Bean (see Canavalia ensi-

formis).

Syagrus calcaratus (on cotton), in Belgian Congo, 315; in Nigeria, 667; susceptibility of varieties of cotton to, 667.

Sybra ordinata, on cotton in For-

mosa, **309.**

Sycamore, pests of, in France, 316. sycophanta, Calosoma.

Sylepta derogata (on cotton), in China, 498; in Belgian Congo, 315; in Formosa, 309; in

Nigeria, 667; in Nyasaland, 114; birds destroying, in Punjab, 180; in Sudan, 513.

sylvaticus, Aspidiotus.

Symitha mangifera, sp. n., on mango in Seychelles, **405.**

Sympiesis ancylae, parasite of Sparganothis sulfureana in New Jersey, 563.

Synacanthotermes zanzibarensis, damaging timber in Tanganyika, 642.

Synanthedon opalescens (see Aegeria). Synaphaeta guexi, drought affecting damage to poplar by, in California, 650.

Synonycha grandis (predacious on Aphids), bionomics and utilisation of, in Formosa, 768, 769.

Syntomaspis druparum, measures against, on apple in Maine, **333.** Syntomaspis eurytomae, biology of,

Syntomosphyrum indicum, imported into Fiji from New South Wales against Dacus spp., 702.

sypnoides, Tolna.

Syprestia carinosa, bionomics of, on Anona in Brazil, 115.

syriacus, Retithrips. syrites, Ceuthorrhynchus.

Syromastes marginatus, in Switzerland, 92.

Syrphus, predacious on Chermes nordmannianae in Germany, 661. Syrphus balteatus, predacious on Eriosoma lanigerum in Japan, 771.

Syrup, in baits: for ants, 588; for Cydia, 137, 168, 506; for Lepismids, 37, 520; types of, in mopping mixtures against Anthonomus grandis, 645.

Systaena taeniata, measures against, on potato in Colorado, 643.

Systates nocivus, sp. n., on coffee in Nyasaland, 405.

Systoechus ctenopterus, bionomics of, in Italy, 163.

Systole, parasite of Lixus scabricollis in Italy, 528.

T.

T-gas, fumigation with, against Lepismids, 37; ignition temperatures of, 313.

tabaci, Thrips.

tabashii, Elachertus.

Tabidia aculealis, on sweet potato in Ceylon, 358.

tabidus, Trachelus.

Tachina aethiopica, parasite of Laphygma exempta in S. Rhodesia,

Tachina cinerascens, parasite of Loxostege sticticalis in Russia. 94: aberrant male of, 94.

Tachina civilis, bionomics of, in Russian Union, 303.

Tachina fallax, parasite of Spodop-tera mauritia in India, 241.

Tachina larvarum, parasite of Euxoa segetum in Italy, 528; parasite of Lymantria dispar in Jugoslavia, 760; parasite of Lasiocampid in Korea, 264.

taeniata, Systaena. taeniopus, Chlorops.

Taeniothrips alliorum, parasite of, in Japan and Loochoo Islands, 142.

Taeniothrips gladioli (see T. simplex). Taeniothrips hawaiiensis, associated with pineapple fields in Hawaii, 213.

Taeniothrips inconsequens, on plum in Britain, 191; on pear in Br. Columbia, 59; on prune in Oregon, 29, 291; measures Oregon, 29, 291; against, 29, 191, 291.

Taeniothrips simplex (Gladiolus Thrips), in Bermuda, 626; in 65, 273, 398, U.S.A., 552: possibly damaging iris, 273; measures against, 65, 398, 552.

Taeniothrips xanthius, on orchids

in Japan, 673.

Tahitian Chestnut (see Inocarpus edulis).

taiwanensis, Carpoglyphus.

takahashii, Nipponaphis (Thoracaphis).

Takamaka (see Calophyllum inophyllum).

Talc, trees banded with, against ants, 83; repellent to Empoasca **736, 740, 766**; as an adhesive for derris spray, **674.**

Talinum, Loxostege bifidalis on, in

Brazil, 333.

Tallow, Gibbium psylloides attacking, in Germany, 519.

Tamarind, Psychids on, in Madras, 499.

tamsuyensis, Melanotus.

Tanacetum vulgare, Calocoris norvegicus on, in Germany, 234.

Tanaomastix abnormis, parasite of Pseudococcus citri in Spain, 701.

Tanganyika Territory, coffee pests in, 68, 113, 178, 179; cotton pests in, 112, 113, 610, 642; wild food-plant of Diparopsis castanea in, 199; new Bruchid on Erythrina in, 292; Cicadulina mbila and streak disease of maize in, 756; miscellaneous pests in, 112, 113, 118, 642; locusts and grasshoppers in, 112, 113, 642, 683; termites in, 113, 642; Trypetids in, 492; Pseudococcus kenyae not occurring in, 592; beneficial insects in, 112, 179, 492, 610, 611; summary of plant quarantines in, 229.

Tangerine (see Orange, Mandarin).

Tanglefoot (see Adhesives).

Tansy (see Tanacetum vulgare). Tanymecus palliatus, birds destroying, in Austria, 255; in Germany, 719; in Russian Union, 237, 432; on beet, 237, 255; on apple and tree seedlings, 432,

719; bionomics of, 237.

Tanymecus sparsus, on Sorghum in

Sudan, 513.

Tar, maize seed and tea treated with, against pests, 362, 602; and lime, soil treatment with,

against wireworms, 631.

Tar Distillates, as dormant sprays against orchard pests, 16, 25, 62, 63, 99, 234, 235, 250, 272, 343, 350, 351, 465, 466, 482, 558, 562, 659, 662, 663, 718, 724, 738; action of, on eggs of Psylla mali, 482, 659; ineffective against Membracid on apple, 269, 574; destroying predacious enemies of Paratetranychus pilosus, 16, 100, **464, 724**; use of, against *Xylo*trechus on coffee, 56; against vine pests, **521**, **701**, **758**; against pests of other plants, 270, 301, 658, 711; soil treatment with, against Rhagoletis cerasi, 300, 347; repellent action of, 56, 235, 351; emulsifiers for, 301, 738; formulae for, 25, 63, 234, 272, 298, 301, 347, 350, 482, 558, 562, 574, 701, 711, 738; and Bordeaux mixture, 234, 235; and lime-sulphur, 234, 235; and quicklime, 701; and mineral oils, 272, 298, 343, 466, 482, 562, 738; causing injury to fruit trees and lavender, 97, 98, 273, 301, 351, 663; types of, 25, 188, 234, 250, 301, 482; effect of freezing on,

188; composition, properties and standardisation of, 97, 98, 99, 156, 300; mineral oil sprays compared with, 16, 62, 270, 466, 482, 562, 663; nicotine compared with, 62, 738.

taragamae, Apanteles.

Taraxacum, Tetranychus on, in Ukraine, 257.

Taraxacum kok-saghyz, pests of, in Ukraine, 72, 106, 618, 619.

Taraxacum magalorrhizon, pests of, in Ukraine, **106, 618, 619.**

Tarichium jachontovi, infesting Coccinellids in Central Asia, 239.

Tarnished Plant Bug (see Lygus pratensis).

Taro (see Colocasia).

Taro Leafhopper (see Megamelus proserpina).

tarsalis, Cephalonomia; Habrolepoidea.

Tarsonemus, on Fatsia in Germany, 660.

Tarsonemus bancrofti, on sugar-cane in U.S.A., 28.

Tarsonemus latus, measures against, on delphinium in New Hampshire, 20.

pallidus, measures Tarsonemus against, on strawberry in Britain, 383; shading favouring infestation of chrysanthemum by, in U.S.A., 396.

spinipes, Tarsonemus associated with stem galls on sugar-cane in Hawaii, 585.

Tarsonemus spirifex, on oats in Britain, 709.

Tarsostenus univittatus, predacious on Minthea rugicollis in Malaya,

Tartar Emetic, in baits for ants, 219; in bait-spray for Trypetids, 325; spraying with, against gladiolus thrips, 552, spreaders and adhesives for, 552, 553; Citrus not affected by mist sprays of, 325; determination of antimony in spray residues of 695; other insecticides compared with, **552.**

Tartaric Acid, in baits for Cydia, 135.

Tasmania, apple pests in, 8-10. 208; Coleopterous strawberry pests in, 208; Penthaleus major on peas in, 582; grasshoppers in, 584; Halotydeus destructor not established in, 582; beneficial insects and biological control in. 10, 208.

tasmaniae, Aphodius. tasmanica, Apanteles. tatei, Tetrastichus. tavaresi, Aphis.

taxicornis, Labidostomis.

Taxus brevifolia, new Coccid on, in U.S.A., **226.**

tayabanus, Pseudococcus.

Tea, pests of, in Ceylon, 240, 337, 360, 639; Tettigoniid on, in China, 244; pests of, in India, 148, 149, 601, 693; pests of, in Nyasaland, 114; new Curculionid on, in Uganda, 612; Aspidiotus perniciosus not developing on, 305; termites damaging, 149, 240, 337; susceptibility of varieties of, to Tetranychus bioculatus, 694; Paris green injections causing injury to, 338.

Tea Tortrix (see Homona coffearia). Teak, pests of, in Burma and India, 338, 339, 764, 765; Phassus signifer on, in Formosa, 517.

Teaseed Oil (see Oil, Teaseed).

Technomyrmex albipes, habits of, in Mauritius, 204.

tecta, Aspidomorpha. Tectona grandis (see Teak).

tectonae, Nemeritis. tectus, Ptinus.

tegalensis, Chionaspis.

telarius, L., Tetranychus. telarius, auct., Tetranychus (see T.

tiliarius).

Telenomus, Asiatic species of, 89; parasite of Ischnodemus saccharivorus in Loochoo Islands, 437; parasite of Aelia rostrata in Spain,

Telenomus acrobates, parasite of Chrysopids in Italy, 189.

Telenomus almanzori, sp. n., parasite of Hylesia nigricans in Argentina, 772.

Telenomus ashmeadi, parasite of Chlorochloa sayi in Montana, 44.

Telenomus attaci, sp. n., parasite of Attacus atlas in Malaya and Siam, 89.

Telenomus beneficiens, parasite of Diatraea venosata in Asia, 89.

Telenomus cyrus, sp. n., parasite of Nezara viridula in Java, 89.

Telenomus dignoides, sp. n., parasite of Scirpophaga intacta in Java, 89.

Telenomus dignus, parasite of Schoenobius bipunctifer in China and Java, **89.**

Telenomus nawai (parasite of Noctuids), introduced into Egypt from Fiji, 212; in Hawaii, 586.

Telenomus ochus, sp. n., parasite of Oreta carnea in Sumatra, 688.

Telenomus olynthus, sp. n., parasite of Pyralid in Sumatra, 688.

Telenomus periparetus, sp. n., parasite of Geometrid in Sumatra, 688. Telenomus remus, sp. n., parasite of

Spodoptera mauritia in Malaya,

Telenomus rowani (parasite of Schoenobius bipunctifer), in China,

109; in Java, 89.

Telenomus sokolowi, bionomics of, parasitic on Pentatomids in Switzerland, 92.

Telenomus tirathabae, in Asia, 89.

Telenomus triptus, sp. n., parasite of Scotinophara coarctata in Malaya,

Telenomus urios, sp. n., parasite of Hidari irava in Malaya, 688.

Telenomus usipetes, sp. n., parasite of Hapalia machaeralis in Burma, 688.

Teleonemia lantanae (attacking Lantana), introduction of, into Australia, 86, 212; in Fiji, 86, 212; introduced into Samoa, 212; bionomics of, 86, 87; technique for transporting, 87.

Telephone Cables (see Electric

Cables).

Telephone Poles (see Timber).

Telephorus, new Nematode experimentally attacking, in Denmark,

Temperature, relations of locusts and grasshoppers to, 6, 54, 72, 105, 146, 147, 410, 416, 679, 756; relations of other insects to, 4, 27, 47, 54, 61, 72, 77, 78, 88, 118, 157, 158, 159, 165, 174, 178, 192, 193, 195, 224, 225, 230, 231, 235, 236, 253, 257, 274, 283, 284, 290, 309, 322, 336, 349, 351, 378, 401, 406, 407 421, 423, 425, 428, 430, 434, 455, 456, 457, 458, 460, 467, 472, 474, 475, 486, 487, 515, 517, 528, 532, 545, 546, 569, 595, 596, 603, 637, 668, 671, 732; relations of mites to, 74, 96, 97, 150, 151, 159, 160, 193, 379, 380, 381, 396, 465, 470, 539, 656.

tenebricosus, auct., Otiorrhynchus

(see O. clavipes).

Tenebrio molitor, oviposition of, 164, 165; tests of contact agents on, 45; sorption of hydrocyanic acid by, 567.

Tenebrio obscurus, tests of methyl bromide against, in stored cereals

in U.S.A., 746.

tenebrioides, Zabrus.

Tenebrionids, preparation of larval specimens of 357.

tenebrionis, Capnodis.

Tenebroides mauritanicus, tests of trichlorethylene against, in stored grain in France, 414; in cassava tubers in Madagascar, 455; in copra and rice in Malaya, 31.

tenellus, Eutettix.

tenera, Acrocercops.

Tennessee, Chalepus dorsalis in, 549 : Heliothis armigera in 509. tenthredinum, Diplostichus.

tenuimaculatus, Adoretus.

Tenuipalpus, possibly transmitting disease of orange in Argentina, 581.

Tenuipalpus obovatus, on tea in India, 149.

tenuis, Engytatus (Cyrtopeltis); Eriobhves.

Tephrosia spp., pests of, in Java, 110,607; pests of, in Porto Rico, 224; as a cover crop against Diaprepes, 751; insecticidal properties and constituents of, 29, 90, 204, 224, 361, 749, 751.

Tephrosin, Mundulea sericea prob-

ably containing, 36.

Tepperia sterculiae, bionomics of, on Brachychiton in New South Wales, 404.

Terastia meticulosalis, on Erythrina in India, 149.

terebrans, Apate.

formosanus, Termes damaging banana in Formosa, 263.

Terminalia catappa, Coccids on, in Haiti, 41; Anomala xanthopleura on, in Loochoo Islands, 516.

terminifera, Chortoicetes.

Termites, attacking living plants, 110, 113, 149, 186, 187, 219, 240, 243, 261, 263, 285, 337, 411, 413, 454, 512, 642; damaging grass, 48, 113, 449; attacking timber, 17, 55, 193, 220, 223, 240, 285, 314, 449, 514, 642, 706, 724, 748; method of testing susceptibility of timbers to, 449; damaging books, 223, 706; measures against, 17, 113, 149, 188, 220, 223, 338, 389, 449, 512, 514, 724, 748; surveys of bionomics of, 240, 314; natural enemies of, 142, 143; Pyrophorus bellamyi reared on, 493; destroying Xyleutes ceramicus, classification and new species of, 219, 413.

Terpinyl Acetate, in baits for Cydia, 135, 506.

terrenus, Bruchus (Bruchidius).

terryi, Pseudococcobius.

Tessaratoma javanica, on Schleichera trijuga in Ceylon, 359.

Tessaromma undatum, on Eucalyptus in New Zealand and Australia, 754.

tessellata, Paroxyna.

tessellatus, Prociphilus; Eucalymnatus (Lecanium).

testacea, Malacosoma neustria; Melipona (Trigona).

testaceipes, Lysiphlebus; Mimela (Anomala).

testudinatus, Periphyllus.

testudonigra, Brachyplatys.

testulalis, Maruca.

tetensi, Dasyneura.

Tetracha chilensis (see Megacephala carolina).

Tetrachlorobenzene, bands treated with, against *Cydia* and *Anthonomus*, **613.**

Tetracium rectisporium, infesting Coccid in Natal, 378.

Tetramethyl Thiurum Bisulphide and Monosulphide (see Thiurum Sulphides).

Tetraneura ulmi (see Byrsocrypta gallarum).

Tetranychina, on Oxalis violacea in Formosa, 674.

Tetranychus, of Central Asia, 470, 530; on Cephaelis ipecacuanha in Formosa, 674; on pear and Camellia in Japan, 53, 438; food-plants of, in Russian Union, 257, 470, 471; acquiring virus of beet curly-top, 445; bionomics and control of, 470, 471; effect of essential oils on, 249, 250.

Tetranychus althaeae (see T. telarius).

Tetranychus bimaculatus (see T telarius).

Tetranychus bioculatus, measures against, on tea in India, 693, 694.

Tetranychus longipenis, sp. n., on apple in Central Asia, **530**.

Tetranychus pacificus, measures against, on vines and fruit trees in U.S.A., 131, 488, 748.

Tetranychus scabrisetus, sp. n., foodplants of, in Central Asia, **530**.

L., Tetranychus in telarius, Australia, 246, 334; on vines in Bessarabia, 71; in Britain, 349, 383, 712; food-plants of, in Formosa and Korea, 674; on castor in India, 55; ecology of, on Citrus in Palestine, 405; tentatively recorded on cassava in Réunion, 706; on shade trees in Russia, 665; in U.S.A., 286, 328, 396, 553, 648, 704; on beans, 334, 704; on greenhouse and ornamental plants, 286, 349, 553, 648, 704, 712; shade favouring infestation of chrysanthemum by, 396; on strawberry, 328, 383; natural enemies and biological control of, 71, 674, 712; other measures against, 71, 246, 286, 328, 334, 383, 553, 648, 665, 704, 712.

Tetranychus telarius, auct. (see T.

tiliarius).

Tetranychus tiliarius, on Tilia in Finland, 201.

Tetranychus turkestani, sp. n., on cotton, etc., in Central Asia, 470, 472, 530; measures against, 472.

Tetranychus virginis, sp. n., on fruit trees and sunflowers in Central Asia, 530.

Tetrapanax papyrifera, Pseudococcus citri on, in U.S.A., 138.

Tetrastichodes (see Tetrastichus).

Tetrastichus, parasite of Ceratitis spp. in E. Africa, 492; parasite of Gracilaria theivora in India, 149; parasite of Diaprepes in St. Lucia, 403.

Tetrastichus brontispae, utilisation of, against Brontispa in Nether-

lands Indies, 337.

Tetrastichus bruzzonis, hosts and utilisation of, in Italy, **528.**

Tetrastichus carinatus (parasite of Mayetiola destructor), competition of, with Platygaster herricki, 121. Tetrastichus dacicida, parasite of Ceratitis spp. in Tanganyika,

492.
Tetrastichus flavovarius, associated with Eumarchalia gennadii in

Crete, **71.**

Tetrastichus giffardi (parasite of Trypetids), in E. Africa, 492; introduced into Brazil from Hawaii, 627; rearing of, 628.

Tetrastichus giffardianus (parasite of Trypetids), in Africa, 492; in Hawaii, 84; attempted introduction of, into other countries, 58, 212.

Tetrastichus marylandensis, parasite of Aphis maidis in Porto Rico, 39.

Tetrastichus pyrillae, parasite of Pyrilla in India, 243.

Tetrastichus rapo, hyperparasite of Brevicoryne brassicae in Ukraine, 303.

Tetrastichus schoenobii, parasite of Schoenobius bipunctifer in China,

Tetrastichus silvaticus, sp. n., parasite of Malacosoma disstria in N. America, 227.

Tetrastichus tatei, sp. n., parasite of Gynaikothrips uzeli in Porto Rico, 41.

Tetrastichus turionum, establish-Rhyacionia ment of. against buoliana in U.S.A., 745.

Tettigonia viridissima, in Cyprus, 108.

Tettigoxenos orientalis, parasite of Nephotettix in Japan, 439.

teutonicus, Cylindroiulus londinensis. texana, Anomis (see A. luridula).

texanus. Chelonus.

Texas, Anastrepha ludens in, 325; cotton pests in, 142, 227, 274, 275, 276, 277, 278, 328; Celama sorghiella on maize and Sorghum in, 279, 560; Acrobasis caryae on pecan in, 138; vegetable pests in, 28, 397; Conopid parasite of honey bees in, 133; beneficial insects and biological control in, 227, 275, 276, 280, 328; insects introduced into Australia from, against Xanthium pungens, 450.

Textiles, pests of, and their control, 227, 228, 400, 514, 519, 770.

Thalassodes quadraria, on Schleichera trijuga in Ceylon, 359.

Thalessa superbiens, parasite of Semanotus japonicus in Japan, 516.

Thallium Salts, in baits for ants, 11, 219, 222, 223, 588; injuring bark of Inga, 223.

Thamnotettix, synonymy of, 78.

Thamnotettix nigrifrons, on narcissus in U.S.A., 288.

Thamnotettix prasinus, type genus, 78.

Thamnotettix (Cicadula) quadrinotatus, **78.**

Thamnurgides (see Poecilips).

thatcheri, Cinara.

Thaumetopoea, distribution of European species of, 518.

Thaumetopoea pinivora (Pine Processionary Caterpillar), in Denmark, 518.

Thaumetoboea processionea, measures against, on oaks and vines in Germany, 664.

theae, Oscinis.

Thecodiplosis brachyntera, on pines in Germany, 312.

Thecodiplosis mosellana (see

Thegilis theobromae, sp. n., on cacao in Brazil, 106.

theivora, Gracilaria.

themei, Diabrotica (see D. pallipes). theobromae, Thegilis.

Theresia (Paratheresia) brasiliensis, parasite of *Diatraea* in Brazil, 82.

Theresia claripalpis (parasite of Diatraea saccharalis), in Peru and Argentina, 391; introduction of, into U.S.A., 391.

Thermobia aegyptiaca, damaging books in Egypt, 706.

Thermobia domestica, bionomics and control of, in houses in Germany and U.S.A., 37, 224, 387.

thermoptera, Titanoceros.

Theronia atalantae, parasite of Lepidoptera in China and Korea, 245, 264; hosts of, in Jugoslavia,

Thespesia spp., relation of cotton pests to, 43, 55, 199, 367, 374,

446, 766.

Thiocyanates, against Coccids, 172, 272; against thrips, 508, 553, 554, 727; against other insects, 131, 690, 740; against mites, 16, 328, 553, 554, 712; dusts containing, 508, 740, 741; spraying with, 16, 131, 172, 272, 553, 554, 690, 712, 727; spreaders for, 554, 727; and oil emulsions, 16, 172, 272; formulae containing, 172, 272, 712, 740; causing injury to plants, 554, 712; other insecticides compared with, 553, (See also Nicotine Thio-690. cyanate.)

Thiodiphenylamine, against Coleoptera, 323, 328, 443, 551, 566, 647; against Cydia spp., 62, 79, 149, 401, 443, 449, 510, 544, 647; against other Lepidoptera, 9, 327, 736; against Rhagoletis spp., 16, 490; against plum sawfly, 615; against Thysanoptera, 644; tests of toxicity of, to Hippodamia and fish, 390, 439; dusting with, 544, 566, 644; talc as a carrier for, 566, 644;

formulae for, in sprays, 16, 149, 443, 490, 510, 551, 615, 736; adhesives and spreaders for, 16, 443, 490, 551, 615; and sulphurs, 443, 490; causing injury to plants, 150, 328; analysis of, 695; compared with other insecticides, 79, 150, 328, 646, 736.

Thiuram Sulphides, in spray formulae as insecticides and repellents, 443, 551, 752.

Thomasiniana oculiperda, on roses in Britain, 297.

thomsoni, Piestopleura.

Thonalmus chevrolati, destroying sugar-cane pests in Porto Rico, 289.

Thoracaphis takahashii (see Nipponaphis).

thoracaphis, Prospaltella.

thoracicus, Stiboscopus; Triaspis.

thraso, Eantis.

Thripobius hirticornis, gen. et sp. n., parasite of Retithrips spp. in Africa, 610.

Thripoctenus brui (parasite of Thysanoptera), bionomics of, in Japan and Loochoo Islands, 142, 143; introduction of, into Hawaii, 143.

Thripoctenus maculatus, bionomics of, in India, 88.

Thrips abdominalis, possibly a vector of pineapple yellow spot in Hawaii, 213; parasite of, in Lapap and Locchoo Islands 142.

Japan and Locchoo Islands, **142.** *Thrips fuscipennis*, bionomics and control of, in Britain, **711.**

Thrips major, on chestnut and beech in Britain, 711.

Thrips nigropilosus, shade favouring, on chrysanthemum in U.S.A., 396.

Thrips panicus, on cotton in S. Carolina, 278.

Thrips tabaci, studies on populations of, in Bermuda, 626; on tobacco in Czechoslovakia, 657; favouring infestation of cotton by Prodenia litura in Egypt, 458; in Germany, 39; in Hawaii, 143, 213, 692, 752; in Japan and Loochoo Islands, 142; in Tanganyika, 113; in U.S.A., 285, 489, 553; susceptibility varieties of carnations to, 489,711; on cucumber, 553; on cyclamen, on onion, 39, 113, 143, 285, 626, 752; relation of, to pineapple yellow spot, 143, 213, natural 692; enemies biological control of, 142, 143,

626; other measures against, 285, 489, 553, 626.

Thrips, Banana Rust (see Scirto-thrips signipennis).

Thrips, Bean (see Hercothrips fas-ciatus).

Thrips, Cacao (see Selenothrips rubrocinctus).

Thrips, Florida Flower (see Frank-liniella cephalica bispinosa).

Thrips, Gladiolus (see Taeniothrips simplex).

Thrips, Greenhouse (see Heliothrips haemorrhoidalis).

Thrips, Iris (see Bregmatothrips iridis).

Thrips, Long-winged (see Scirto-thrips longipennis).

Thrips, Pear (see Taeniothrips inconsequens).

Thuja, pests of, in U.S.A., 126, 392, 441; effects on, of soil treatment with lead arsenate, 441.

Thurberia thespesioides, eradication of, against Anthonomus grandis thurberiae in Arizona, 329.

thurberiae, Anthonomus grandis.

 $thur be riella,\ Bucculatrix.$

thyridopterigis, Habrocytus.

Thyridopteryx ephemeraeformis, bionomics and control of, in U.S.A., 173, 543.

Thyrsocnema (see Sarcophaga).

Thysanoptera, of Kauai (Hawaii), 692; and plant diseases, 213, 318, 349, 581, 692; predacious on other insects, 670; predacious on mites, 71, 712, 724; natural enemies of, 10, 41, 88, 273, 610, 626; classification and new species of, 318, 329, 457, 490.

Thysanus dipterophaga, hyperparasite of Diatraea in Br. Guiana, 491.

tibialis, Chaetocnema; Phanerotoma. tigrinus, Cleonus (Cyphocleonus). Tilia (see Lime, Tilia).

tiliarius, Tetranychus.

Timber, Coleoptera in, 17, 34, 91, 155, 190, 191, 195, 214, 233, 301, 354, 379, 381, 450, 463, 514, 530, 591, 599, 635, 662, 707, 726, 763; termites in, 17, 55, 193, 219, 220, 240, 285, 314, 449, 514, 642, 724, 748; other pests of, 149, 191, 218; key to pests of, in Central Europe, 196; survey of pests of, in Sweden, 440, 624; possible value of X-rays for detecting insects in, 190.

Timbo, Bordeaux mixture not reducing efficiency of, against Epilachna varivestis, 270.

Timocratica albella, on Casuarina in

Brazil, 145.

Timothy Grass (see Phleum prat-

ense).

Tineola biselliella, 460; bionomics of, in Japan, 770; not used for testing moth-proofing solutions in U.S.A., 227; characters of eggs of, 196.

Tiphia, new species of, in Japan

and Korea, 108.

Tiphia asericae, liberation of, against Aserica castanea in U.S.A., 325.

Tiphia popilliavora, utilisation of, against Popillia japonica in U.S.A., 324, 545, 723, 727; Korean strain of, 324.

Tiphia sternata, liberation of, against Aserica castanea in U.S.A., 325.

Tiphia vernalis, utilisation of, against Popillia japonica in U.S.A., 324, 545, 552, 723, 726; adult feeding habits affecting parasitism by, 552.

Tipula, on cereals in Britain, 296. Tipula aino, measures against, on rice in Japan, 53, 771; on wheat,

54.

Tipula paludosa, new Nematode experimentally attacking, 50.

tipulae, Leptacis.

Tiracola plagiata, on castor and Citrus in Ceylon, 359; Trichospilus pupivora reared on, 359.

Tirathaba trichogramma, parasites of, on coconut in Fiji, 334.

tirathabae, Apanteles; Echthromorpha; Telenomus.

Tischeria complanella, on oak in

Finland, 201.

titanis, Lachnosterna (Holotrichia). Titanoceros thermoptera, predacious on Ochrogaster contraria in Australia, 449.

Tlanoma (see Chaetocnema).

Tmetocera (see Eucosma). Toads (see Bufo spp.).

Toads, Horned (see Lizards).

Tobacco, Myzus persicae on, in S. Africa, 177; Frankliniella paucispinosa on, in Argentina, 581; Phthorimaea operculella on, in Brazil, 145, 320; pests of, in greenhouses in Canada, 537, 623; Thrips tabaci on, in Czechoslovakia, 657; Nysius cymoides on in Cyprus, 108; pests of, in India, 55, 88, 242, 606; pests of,

in Indo-China, 185, 205; Macrosiphum solanifolii on, in Italy, 759; book on pests of, and their control in Poland, 72; Phaulacridium gemini on, in Queensland, 494; pests of, in S. Rhodesia, 106, 377; pests of, in Sumatra, 603; pests of, in Tanganyika, 642; Barathra brassicae on, in Ukraine, 159; pests of, in U.S.A., 132, 133, 144, 183, 393, 726, 744; pests of, in W. Indies, 40, 490; susceptibility of varieties of, to Myzus persicae, 177; insects and virus diseases of, 88, 242, 377, 578, 581, 759; scorched by barium fluosilicate, 133; suitability of, for rearing Encarsia formosa, 623.

Tobacco (Stored), pests of, and their control, 328, 363, 396, 490, 523; (seed) Orphinus aethiops

in, in Java, 434.

Tobacco (as an insecticide), dusting with, 65, 498, 652; as a diluent for dusts, 65, 133, 652; fumigation with, 274; spraying with, 72, 88, 131, 221, 241, 376, 422, 659, 663, 693, 716; duration of activity of sprays of, 422; against pests in soil, 209, 645; formulae containing, 131, 221, 241, 652, 693; and lime, 498; and soap, 72, 88, 376, 422, 663, 716; and soap-bark powder, 221.

Tobacco Capsid (see Engytatus

tenuis).

Tobacco Flea-beetle (see Epitrix parvula).

Tobacco Moth (see Ephestia elutella).

Tobacco Stem Borer (see Phthorimaea heliopa).

Tobago, summary of plant pest legislation in, 491.

Tolna sypnoides, damaging fruit in Sierra Leone, 239.

Tomaspis, ecology and natural enemies of species of, in Brazil and W. Indies, 369.

Tomaspis saccharina (Sugar-cane Froghopper), in Grenada, 752; in Trinidad, 10, 370, 404; varieties of sugar-cane resistant to, 10; measures against, 10, 404; question of biological control of, 370.

Tomaspis tristis, natural enemies of, in Brazil, **370.**

Tomato, pests of, in Argentina, 290, 581; pests of, in Australia,

103, 210, 318, 334, 496; Epicauta spp. on, in Brazil, 290; pests of, in Br. Isles, 349, 710, 712; pests of, in greenhouses in Canada, 537, 623; Epilachna vigintioctopunctata on, in Ceylon, 358; mites on, in Cyprus, 108; Nezara viridula on, in Egypt, 457; pests of, in Hawaii, 213, 214, 752; pests of, in India, 409; Capsid on, in Libya, 260; Phyllocoptes on, in New Zealand, 45, 103; Dacus cucurbitae on, in Philippines, 181; thrips on, in Sierra Leone, 240; Bemisia gossypiperda on, in Sudan, 513; Bemisia little attacked by Heliothis armigera in eastern Transvaal, 364; pests of, in U.S.A., 13, 19, 20, 117, 127, 217, 327, 397, 488, 553, 580, 644, 744, 747; Epicauta spp. on, in Uruguay, 290; susceptibility of varieties of, to D. cucurbitae, 213; insects and virus diseases of, 260, 318, 349, 581, 759; relation of, to beneficial insects, 13, 623; insecticides causing injury to, 210, 554, 712.

Tomato Leaf-miner (see Agromyza

solani).

Tomato Mite (see *Phyllocoptes lyco-persici*).

Tomato Pin Worm (see *Phthorimaea lycopersicella*).

Tomato Stem Borer (see Phthorimaea plaesiosema).

tomentosus, Rhynchites.

Tomostethus juncivorus, on Juncus in Japan, 672.

Toon Shoot Borer (see *Hypsipyla robusta*).

Torches, against Promecotheca, 495. torquata, Haltica.

torquens, Contarinia (see C. nas-turtii).

torridus, Sphex.

tortricis, Anachaetopsis.

Tortrix, bionomics and control of species of, attacking fruit trees in Caucasus, 477.

Tortrix argyrospila, measures against, on fruit trees in Br. Columbia and New York, 144, 574.

Tortrix diversana, on apple and pear in Caucasus, 477.

Tortrix excessana, bionomics and control of, on stone fruits in New Zealand, 57, 58.

Tortrix heparana, on apple and pear in Caucasus, 477, 478.

Tortrix lecheana, on apple and pear in Caucasus, 477.

Tortrix longicellana, bionomics and control of, on apple in Korea, 243. Tortrix occidentalis, on onion in

Tanganyika, 642.

Tortrix piceana, on Pinus cembra in Austria, 664.

Tortrix podana, on field and stored apples in Britain, 612; gas storage, a protection against, 612.

Tortrix postvittana, bionomics and control of, on fruit trees etc. in Tasmania and New Zealand, 8–10, 57, 58.

Tortrix pronubana, on ornamental plants in Britain, 349, 350.

Tortrix reticulana, on apple and pear in Caucasus, 477, 478.

Tortrix rosaceana, bionomics and control of, on dewberry in Utah, 141.

Tortrix rosana, on myrtle and cranberry in Holland, 299.

Tortrix viridana, tests of baits against, on oak in Germany, 50.
Tortrix xylosteana, on pear in Japan, 604; measures against, on apple in Korea, 243.

Toxicarol, forms and precursor of, and their toxicity to insects, 616,

631, 632.

Toxoptera aurantii (on Citrus), in Cyprus, 706; probably in Libya, 260; bionomics of, in Palestine, 379, 380; tests of insecticides on, in Tanganyika, 35; measures against, 380, 707.

Toxoptera graminum (on cereals), natural enemies of, in Argentina and Uruguay, 30, 31, 289, 290; in U.S.A., 577; on flax, 289, 290; transmitting sugar-cane mosaic, 577; measures against, 290.

Toxoptera piricola, measures against, on pear in Japan, 604.

Trachea secalis, on cereals in Holland, 299.

Trachelus tabidus, on wheat in Ohio, 130.

trachynotus, Meteorus.

Trachyspermum copticum, experiment in infestation of, by insects and mites, 250.

Tradescantia, Blapstinus substriatus fed on, 27.

Tragischoschema wahlbergi, on cotton etc. in Nyasaland, 114.

Trama troglodytes, ant associated with, on rubber-producing plants in Ukraine, 106, 107.

tranquebaricus, Apoderus.

transversa, Acontia.

Traps, for Coleoptera, 403, 625, 719, 727; for *Gryllotalpa*, 432; for Trypetids, 58, 108, 407, 697, 699, 700; description of, 719.

Trenches, against Hybalus cornifrons, 757; against locusts, 611; against termites, 338; against weevils on clover, beet, etc., 75, 132, 236, 256, 520, 722.

Trialeurodes abutilonea, on beans in

Virginia, 580.

Trialeurodes porosus, sp. n., on Zizyphus spina-christi in Egypt, 440.

Trialeurodes vaporariorum (Greenhouse Whitefly), utilisation of Encarsia formosa against: in Australia, 449, 582; in Canada, 623; in New Zealand, 582; food-plants of, in relation to E. formosa, 623.

Trialeurodes variabilis (Papaya Whitefly), Coccinellid predacious on, in Florida, 727; sulphur

dusts against, 727.

Trialeurodes williamsi, sp. n., on ferns in greenhouses in Britain, 772.

Triaspis thoracicus, importation and liberation of, against Bruchids in U.S.A., 65, 329, 580.

Tribolium, on field maize in S. Carolina, 643; in flour mills and stored grain in U.S.A., 130, 389; tests of methyl bromide against, 389.

Tribolium castaneum, in stored grain in Algeria and France, 413, 414; in butter-nuts imported into Holland from Dutch Guiana, 529; in stored ginger in Jamaica, 491; associated with copra and stored rice in Malaya, 31; in stored cassava tubers in Madagascar, 455; in flour warehouses in U.S.A., 322; susceptibility of, to low temperatures, 322; uses and tests of fumigants against, 31, 322, 413, 414, 571, 633; other measures against, 31, 32, 322; derris dusts ineffective against, 183.

Tribolium confusum, population studies of, in Java, 311; in stored cassava tubers in Madagascar, 455; in stored grain and flour in U.S.A., 116, 321, 322, 326, 389, 390; susceptibility of, to low temperatures, 322; studies on oviposition of, 164, 165; uses and tests of fumigants

against, 116, 321, 322, 326, 389, 390, 567, 575, 741; other measures against, 116, 322, 326; tests of homologues of Paris green on, 30.

Tribolium ferrugineum (see T. castaneum).

n in in

Tribolium navale, auct. (see T. castaneum).

Tribulus terrestris, effect of feeding on, on Schistocerca gregaria, 628. tricarinata, Stobaera.

Trichacis remulus, imported into U.S.A. against Mayetiola destructor, 329.

Trichiocampus viminalis, on poplar in Connecticut, 17.

Trichlorethylene, tests of, as a fumigant, 414.

Trichogramma (parasite of Lepidopterous eggs), in Barbados, 222; in Burma and India, 765; in Japan, 671; in Madagascar, 67; in Tasmania, 10; in U.S.A., 14, 16, 725; field utilisation of, and its value, 14, 16, 222, 725, 765; susceptibility of, to cold, 725; parasite of Sialis, 102; problems and status of forms of, 102, 119, 398, 744.

Trichogramma australicum, 67; parasite of Chilo simplex in Japan, 671; local host for mass-breeding of, in Mauritius, 176.

Trichogramma dendrolimi (dendrolimusi), hosts of, in Japan, 671, 672.

Trichogramma evanescens, parasite of Noctuids in Japan, 605, 671; rearing and utilisation of, against Lepidoptera in Russian Union, 73, 235, 236, 356; longevity and productivity of, 545; factors concerned in host-finding by, 150; considered correct name for T. minutum, 398; T. semblidis distinct from, 102.

Trichogramma japonicum (parasite of Lepidopterous eggs), hosts and artificial rearing of, in Japan, 515, 669, 670, 671, 672; effect of temperature and humidity on, 515, 670.

Trichogramma minutissimum (see T. minutum).

Trichogramma minutum (parasite of Lepidopterous eggs), in Barbados, 448; in India, 409, 765; in Peru, 627; in Porto Rico, 289; in U.S.A., 398, 744; rearing of, on Corcyra cephalonica, 409; utilisation of, against forest pests,

765: suggested utilisation of, against Ephestia elutella, 524; T. minutissimum the summer phase of, 744; T. evanescens considered correct name for, 398.

Trichogramma pretiosum, hosts of, in U.S.A., 744; var. nigra a phase of, 744.

Trichogramma pretiosum var. nigra (see T. pretiosum).

Trichogramma semblidis, status, hosts and dimorphism of, in Britain, 102.

trichogramma, Tirathaba.

Trichomalus trujilloi, sp. n., parasite of Cydia molesta in Uruguay,

Trichomma enecator, parasite of Cydia molesta in Italy, 154.

Trichopoda pennipes, predacious on Nezara viridula in Florida, 727.

Trichopsidea oestracea, bionomics of, in Australia, 629.

Trichospilus pupivora (parasite of Nephantis serinopa), bionomics and utilisation of, in Ceylon, 359; in Cochin, 452; laboratory host of, 359.

tricinctus, Closterocerus.

tricoloripes, Hemiteles. tricostata, Eleodes.

Tridacus (see Dacus).

tridentata, Eutetrapha (Saperda).

Triethanolamine Oleate, properties and uses of, in sprays, 45, 174, 388, 652, 704, 739.

trifasciata, Coccinella.

Trifidaphis phaseoli, measures against, on cotton in S. Carolina, 645; ants associated with, on rubber-producing plants Ukraine, **106, 107.**

Trifolium (see Clover).

Trifolium agrarium, Nodonota puncticollis on, in New York, 288.

Trifolium alexandrinum, Prodenia litura on, in Palestine, 638.

Trifolium hybridum, Apion seniculum on, in Latvia, 256; pests of, in U.S.A., 12, 287; Macrosiphum onobrychis transmitting vein-mosaic to, 12.

Trifolium subterraneum, pests of, in Victoria, 84, 85.

Trigona (see Melipona).

Trigonogastra, parasite of Oscinis theae in India, 149.

Trigonura hicoriae, parasite of Magdalis spp. in Connecticut, 15.

trilineata, Hyperaspis; Lema. trimaculata, Macropsis.

Trinidad, cotton pests in, 66; miscellaneous pests in, 10; sugarcane pests in, 10, 370, 404; beneficial insects and biological control in, 10, 370; summary of plant pest legislation in, 491; insecticides for use in, 533.

Trinodes hirtus, bionomics of, in

Japan, 515.

Trionymus peregrinus, sex ratio and

status of male of, 105.

Trionymus sacchari (on sugar-cane), in Bombay, 261; in Madagascar, ants associated with, in Natal, 411; in W. Indies, 39, 269; susceptibility of varieties of sugar-cane to, 39; Sorghum, 39.

Tripolitania (see Libya).

triptus, Telenomus.

tripunctata, Drosophila; Geomyza (Balioptera).

triquetra, Homalodisca.

Trirhithrum (see Ceratitis).

Tristania, Coloborhombus fasciati-

pennis on, in Java, 295. tristis, Anasa; Eurytoma; Hetero-nychus; Lema; Tomaspis. tristriatus, Eriophyes.

tritici, Contarinia; Frankliniella. Triumfetta, Dysdercus spp. on, in N. Rhodesia and Uganda, 366, 373.

trivialis, Antestia. trochala, Procometis.

troglodytes, Trama.

Trogoderma granarium, review of data on, in stored grain, 501; danger of establishment of, in France and Fr. Colonies, 501.

Trogoderma versicolor, tests methyl bromide against, in stored

cereals in U.S.A., 746. Trophocampa (see Casinaria).

Tropidacris latreillei, on coconut in Br. Guiana, 369.

trujilloi, Trichomalus.

tryoni, Dacus (Chaetodacus) (see D. ferrugineus); Diachasma (Opius).

Trypanea stellata, bionomics and control of, on Taraxacum koksaghyz in Ukraine, 619.

Trypetids (Fruit-flies), studies on, in India, 602; legislation dealing with control of, in New South Wales, 267.

Tsuga, Otiorrhynchus ovatus on, in nurseries in New York, 392; lead arsenate causing injury to,

tuberculatus, Peritetranychus. tuberosula, Platygaster.

949 ÍNDEX.

tucumanus, Ipobracon.

Tulip, Anuraphis tulipae on, in Britain, 350; new mite on imported, in California, 650.

tulibae. Anuraphis (Aphis):

Eviophyes.

tumulosus, Ligyrus.

Tung-oil Tree (see Aleuvites fordi). Tunisia, Lindorus lophanthae predacious on Coccids in, 410.

Turf, Anomala horticola damaging, in Britain, 462; Lamellicorns damaging, in France, 229; Aphodius howitti damaging, in New South Wales, 209; Cirphis unipuncta damaging, in Nova Scotia, 441; termites damaging, in Tanganyika, 113; insects damaging, in U.S.A., 27, 398, 588, 589, 726, 727, 729.

turionana, Rhyacionia. turionum, Tetrastichus.

turkestani, Tetranychus (Eotetrany-

Turkey, Hylotrupes bajulus in building timbers in, 530; Bostrychids boring in lead sheathing of cables summary of plant in, 232; quarantines in, 229.

Turkeys, destroying grasshoppers and crickets in Utah, 128.

Turmeric, pests of, in Ceylon, 359.

Turnip, Hispa armigera on, in China, termite damaging, in Florida, 285; Cirphis unipuncta on, in Nova Scotia, 441; Athalia sjöstedti on, in Tanganyika, 113; Schistocerca gregaria experimentally fed on, 410; methyl bromide not impairing germination of seeds of, 389.

Turnip Beetle, Red (see Entomoscelis adonidis).

turnus, Papilio glaucus.

Turpentine, fumigation of seed coffee with, against Stephanoderes hampei, 359.

Tychius flavus, measures against, on

lucerne in Russia, 76.

Tychius quinquepunctatus, on peas in Czechoslovakia, 657.

Tyloderma fragariae, bionomics and control of, on strawberry in Kentucky, 736.

Typha latifolia, Noctuid on, in

Czechoslovakia, 345.

typhae, Phragmatiphila (Nonagria). Typhlocyba, new Crabro predacious on, in Ohio, 369.

froggatti (australis, Typhlocyba Frogg.) (on apple), in Australia, 208, 494; in New Zealand, 56, 208; on blackberry. 495: population studies on, 56: parasites and biological control of, 57,

Typhlocyba pomaria (on apple), in Nova Scotia, 534; bionomics of, in U.S.A., 57, 556, 557-559, 646; measures against, 534, 557, 558, 559, 646.

typhlocybae, Aphelopus. typographus, Ips.

Tyria jacobaeae, introduced into Australia against ragwort, 450; Harpobittacus predacious on, 450.

Tyroglyphus dimidiatus (see Tyro-

phagus putrescentiae).

Tyroglyphus farinae, measures against, infesting cheese in Ontario, 539; in stored cereals in Russian Union, 95, 96, 97; bionomics of, 96, 97, 539.

Tyroglyphus krameri (see

hardia).

Tyroglyphus lintneri, Tyrophagus putrescentiae recorded U.S.A., 286.

Tyroglyphus longior (see Tyrophagus

putrescentiae).

Tyroglyphus mycophagus (see Caloglyphus)

Tyroglyphus ovatus (see Aleuro-

glyphus).

Tyroglyphus putrescentiae (see Tyrophagus).

Tyrophagus noxius, in stored cereals etc. in Russian Union, 95, 96, 97; bionomics of, 97.

Tyrophagus (Tyroglyphus) putrescentiae, on mushrooms in Britain, Holland and U.S.A., 286, 297, 656, 712; on potatoes and cereals in Russia, 95, 422; measures against, 656, 712.

putrescentiae Tyrophagus castellanii, on mushrooms Britain, 297.

U.

Ufens, parasite of Diaprepes in St. Lucia, 403.

Uganda, cotton pests in, 83, 372-375, 376; coffee pests in, 151, 372, 610; new weevil on tea in, 612; locusts in, 112, 372, 679 Trypetids in, 492; beneficial insects in, 374, 375, 492, 610.

ugandanus, Dysdercus; Paraplesius.

uhleri. Horistonotus.

Sarcophaga (Thyrsoculiginosa, nema).

Eriosoma (Schizoneura); ulmi. Tetraneura (see Lepidosaphes; Byrsocrypta gallarum).

ulmi, auct., Oligonychus (Meta-tetranychus) (see Paratetranychus pilosus).

ulmicola, Atanycolus; Stomaphis.

Ulmus (see Elm).

Ultrawet, increasing efficiency of derris dust, 285; uses of, in sprays, 285, 548, 725; properties and composition of, 285, 548.

Uncaria gambir, Lepidopterous pests of, in Sumatra, 688.

undalis, Hellula (Oebia).

undata, Oncometopia.

undatum, Tessaromma.

undecimnotata, Semiadalia.

undulata, Phyllotreta.

unicolor, Byturus; Chlorotettix; Hockeria; Rhogas; Scopaeothrips.

unicostata, Monosteira.

unifasciata, Belvosia; Comperiella. unionalis, Margaronia (Glyphodes). unipuncta, Cirphis (Leucania).

uniseta, Voriella.
United States of America, cereal pests in, 78, 121, 135, 285, 326, 329, 367, 577, 580, 696, 729, 730; Citrus pests in, 41, 131, 324, 325, 722; cotton pests in, 133, 134, 229, 274, 275, 276, 277, 278, 279, 327, 328, 329, 384, 580, 686, 722; pests of forage crops in, 23, 135, 216, 326, 327, 580, 696; pests of forest and shade trees in, 215, 293, 325, 326, 329, 330, 369, 400, 441, 484, 487, 542, 543, 546, 547, 551, 576, 580, 581, 588, 589, 696, 745, 749; pests of greenhouse and ornamental plants in, 20, 21, 78, 273, 288, 328, 386, 396, 541, 550, 692, 704; mushroom pests in, 286, 328, 579; orchard pests in, 137, 173, 320, 323, 343, 389, 442, 450, 484, 542, 554, 555, 560, 696, 722, 731, 732, 733, 734, 745; pests of small fruits in, 324, 328; pecan pests in, 323, 324, 722; book on pests of subtropical fruits in, 722; sugar-cane pests in, 28, 137, 326, 577; pests of field and stored tobacco in, 132, 133, 328, 396; pests of stored grain and flour in, 64, 118, 130, 321, 326, 362, 389, 746; pests in other stored products and houses in, 224, 227, 228, 284, 367, 387, 400, 568, 746; vegetable pests

in, 21, 26, 140, 285, 326, 327, 328, 397, 565, 580, 651, 704, 731, 739; vine pests in, 131, 173, 323, 324, 722; insects feeding on purslane in, 218; Anabrus simplex in, 44, 128, 580, 609, 686; lists and new species of Aphids in, 119, 247, 304, 369, 496, 696, 772; Aserica castanea in, 215, 324, 325; grasshoppers in, 123, 282, 326, 580, 588, 676, 680, 681, 686; Lymantria dispar and Nygmia phaeorrhoea in, 14, 216, 332, 560, 589, 646, 723; Popillia japonica in, 14, 23, 108, 216, 324, 384, 441, 443, 444, 543, 545, 551, 552, 723, 726, 727, 751; insects and plant diseases in, 78, 173, 288, 325, 328, 386, 551, 577, 749; risk of spread of insects by wind in, 215, 216; beneficial insects and biological control in, 1, 41, 78, 121, 133, 137, 215, 323, 324, 325, 328, 329, 410, 450, 576, 580, 588, 589, 729, 730, 744, 745; fungi infesting Magicicada septendecim in, 81, 395; plant pest legislation in, 133, 134, 228, 229, 750; organisation economic entomology in, 23, 136; literature of economic entomology in, 367, 772; list of publications on insecticides in, 48; patents relating to pest control in, 48, 304, 624; introduction of beneficial insects into other countries from, 57, 154, 329, 575, 594, 595; Schistoceros hamatus imported into England from, 191. (See also under the various States.)

univittatus, Tarsostenus.

Uragoga ipecacuanha (see Cephaëlis). Urania Green, spraying with, against Cydia pomonella, 479, 658; and lime, 479; causing injury to apple, 658; other insecticides compared with, 479. Urapteroides astheniata, bionomics of, on Endospermum malaccense in Malaya, 110.

Urena lobata, not attacked by Platyedra gossypiella in Antigua, 446; Earias fabia on, in Java, 766; new weevil on, in St. Lucia, 405.

urichi, Liothrips.

urios, Telenomus (Aholcus). Urnisa, in W. Australia, 584. urozonus, Eupelmus.

Urtica dioica, Psylliodes attenuata on, in Poland, 719.

Uruguay, Bostrychids boring in lead sheathing of cables in, 232; miscellaneous pests in, 290, 582; beneficial insects in, 290, 582; new Cynipid parasite of Aphidius platensis in, 290.

usambarica, Antestia; Hyperaspis.

usambica, Dirphya.

usipetes, Telenomus.

Utah, beet pests in, 127, 128, 283, 743; miscellaneous pests in, 126-128, 141, 326; lists and new species of Aphids in, 119, 247, 369, 496, 696, 772; grasshopper survey in, 48; beneficial insects in, 127, 141; birds and lizards destroying insects in, 128, 397.

utahensis, Cinara. utilis, Pseudaphycus. uzeli, Gynaikothrips.

V.

vacciniana, Rhopobota (see R. naevana).

vaccinii, Dasyneura; Frankliniella; Mineola.

Vaccinium spp., insects on, in Germany, 69, 344, 345, 461; Cryptocephalus incertus on, in Massachusetts, 646. (See Blueberry and Cranberry.)

Vacuum Fumigation, 24, 84, 388,

541, 593, 597, 750.

vaga, Schistocerca.

Valanga nigricornis, damaging tobacco in Sumatra, 603.

Vanduzea segmentata, measures against, on chrysanthemum and egg-plant in Louisiana, 395; in Mexico, 395.

Vanessa (Pyrameis) cardui japonica, on Arctium lappa in Japan, 673. Vanessa io geisha, on hops in Japan,

Vanilla, Synthetic, in baits for Dacus, 58.

vaporariorum, Trialeurodes. varia, Cuphocera; Idioderma.

variabilis, Hypera (Phytonomus); Lacon; Mylabris; Sericothrips; Trialeurodes.

varians, Neoproba (Cyrtopeltis) (see Engytatus geniculatus).

variegana, Argyroploce.

variegata, Adonia. variegatus, Meloë.

variolosa, Poecilonota. varipes, Omphale.

varius, Chlorophorus ; Megatoma.

varivestis, Epilachna.

vastatrix, Phylloxera (see P. vitifoliae).

Vateria indica, gum of, in banding mixture against ants, 93.

Vatsol, use of, in removing spray residues from cherries, 141.

Vegetable Black, as a warning colour in baits, 682.

Vegetable Marrow, Heliothis armigera on, in S. Africa, 364.

Vegetable Weevil (see Listroderes obliquus).

vehemens, Acicnemis. velata, Epicauta. velutinus, Cerambyx. venosata, Diatraea.

ventralis, Exochomus; Rhizobius. ventricosus, Pediculoides. venustissima, Cheletomorpha.

Veratrine, dust containing, against Bupalus piniarius, 313.

verbasci, Anthrenus.

vermiculus, Lepidosaphes. Vermont, Diprion polytomum on spruce in, 487; new parasite of

Malacosoma disstria in, 227.

vernalis, Tiphia. vernata, Paleacrita.

versicolor, Meteorus; Trogoderma.

versuta, Graphocephala.

Verticillium, infesting Gasterocercodes brasiliensis in Brazil, **504.** Verticillium coccorum, infesting

Coccids in Germany, 101.

vesicatoria, Lytta. vesicularis, Eupelmella.

Vespa orientalis, measures against,

in Britain, 108.

Vesperus, study on, 699.

vestigialis, Euxoa (Agrotis). vestitus, Hylesinus (Chaetoptelius). vesuviana, Carpomyia.

Vetch Bruchid (see Bruchus brachialis).

Vetches (Vicia spp.), Pachytychius strumarius on, in Morocco, 641; Tortricid on, in Quebec, 60; Bruchus brachialis on, in U.S.A., 23, 580; planted among cereals against Blissus leucopterus, 78.

vibex, Cassida.
Vicia (see Vetches).
Vicia faba (see Beans).
vicina, Cydonia; Oxya.
vicinus, Homorocoryphus;

icinus, Homorocoryphus; Scapteriscus.

Victoria, Aonidiella aurantii on Citrus in, 85; pests of subterranean clover in, 84, 85; miscellaneous pests in, 449, 582; Cydia spp. in orchards in, 168, 448; new thrips on rose in, 318;

locusts and grasshoppers in, 583, 584, 676; biological control of noxious insects and plants in, 84, 169, 448, 450; Harpobittacus destroying Tyria jacobaeae in, 450; restrictions on movement of vines from, against Phylloxera, 608.

victoria, Brachymeria (Chalcis).

vigilans, Senotainia.

vigintioctopunctata, Epilachna.

vigintipunctata, Melosoma.

vigintiquatuorpunctata, Subcoccinella.

Vigna (see Cowpeas). vilella, Platyedra.

Villa hottentotta, parasite of Euxoa segetum in Italy, **528.**

villosa, Ochrosidia.

viminalis, Trichiocampus.

Vine, Grape, Haltica ampelophaga on, in Algeria, 371; pests of, in Australia, 111, 335, 608; Macropsis trimaculata on, in Canada, 173; Nysius cymoides on, in Cyprus, 108; Anomala vitis on, in Czechoslovakia, 657; pests of, in France, 415, 619; pests of, in Germany, 4, 154, 419, 521, 601, 661, 664; Rhipiphorothrips cruentatus on, in India, 87: Pseudococcus citri and sooty fungus on, in Italy, 757; Phylloxera vitifoliae on, in Japan, 560; Polychrosis botrana on, in Jugoslavia, 720; Anomala chamaeleon on, in Korea, 767; Sphingid on, in Libya, 201; Pseudococcus vitis on, in Malta, 469; pests of, in Palestine, 406, 638; pests of, in Rumania, 71, 721; Labidostomis taxicornis on, in Sicily, 70; pests of, in Spain, 700, 701; pests of, in Switzerland, 299; pests of, in U.S.A., 19, 23, 127, 131, 173, 285, 286, 323, 608, 722, 748; Dacus oleae associated 161; susceptibility of with, varieties of, to pests, 88, 560; of winter review measures against pests of, 521; insecticides causing injury to, 285, 323, 701; absorption of selenium by, 131. (See Grapes.)

Vine Moths (see Clysia ambiguella and Polychrosis botrana).

Vinegar, in baits for *Cydia* and *Ceratitis*, **137**, **168**, **407**, **699**, **700**; types of, **168**, **407**.

vinula, Dicranura (Cerura). violentus, Rhynocoris,

Violet, pests of, in Britain, 350, 589; unfavourable food-plant for Tetranychus, 471.

Virachola (Deudorix) bimaculata, bionomics of, on coffee, etc., in

Sierra Leone, **239.** Virachola (Deudorix) lorisona, on

coffee in Tanganyika, 113.
virescens, Charagia; Eterusia

(Heterusia); Heliothis.

virgata, Ferrisiana (Ferrisia, Pseudococcus).

Virginia, cabbage insects in, 13; pests of leguminous plants in, 327, 549, 580, 609; maize pests in, 144, 304, 730; orchard pests in, 271, 272, 506, 507, 556; pests of field and stored tobacco in, 328, 363, 393, 744.

virginiensis, Chalcophora.

virginis, Tetranychus (Apotetranychus).

virguncula, Porthesia.

viridana, Tortrix. viridanus, Myllocerus.

viridanus, Myllocerus. viridescens, Eupteromalus.

viridis, Agrilus; Cassida; Chermes; Coccus (Lecanium); Doloessa;

Smynthurus. viridissima, Palomena; Tettigonia.

viridius, Chlorotettix. viridula, Nezara.

Virus Diseases, of solanaceous plants, 32, 47, 69, 77, 88, 89, 242, 260, 318, 342, 349, 377, 379, 495, 527, 564, 581, 615; of other plants, 55, 68, 78, 89, 104, 143, 167, 173, 208, 213, 239, 242, 291, 349, 358, 383, 385, 386, 419, 439, 496, 531, 532, 581, 615, 666, 667, 692, 756; relation of Aphids to, 32, 47, 68, 104, 239, 291, 342, 377, 379, 383, 495, 527, 564, 615 relation of other insects to, 55, 68, 77, 78, 89, 143, 167, 173, 208, 213, 242, 260, 318, 349, 358, 385, 386, 419, 439, 496, 532, 581, 666, 667, 692, 756; possible relation of mites to, 581; book on, 68. (See Curly-top, Mosaic, Rosette and Fiji Diseases.)

viteana, Polychrosis.

viticida, Fidia.

vitifoliae, Phylloxera.

Vitis vinifera (see Vine, Grape).

vitis, Anomala; Chionaspis; Epitrimerus; Phyllocoptes; Pseudococcus (Dactylopius); Pulvinaria. vitrea, Chalioides.

vitripennis, Brachymeria; Hyaliodes; Mormoniella,

vittata, Cassida; Diabrotica (see D. melanocephala); Phyllotreta. vittatum, Phaulacridium.

vittatus, Pantophthalmus; Prepodes. Volck Oil, 484, 546.

TOTAL Anthony

vorax, Anthrenus; Dermestes.

Voria ruralis, parasite of Polia oleracea in Italy, 528.

Voriella uniseta, parasite of Tortrix postvittana in Tasmania, 10.

vulgaris, Asaphes (Isocrates); Chrysopa; Dilophus (see D. febrilis); Melolontha (see M. melolontha); Zenillia (Phryxe). vulnerans, Doratifera.

vulnerator, Pristomerus. vulpinus, Dermestes.

vuteria, Sesamia.

W.

wahlbergi, Chilocorus; Tragischoschema.

walkeri, Pseudococcus. wallacei, Crossotarsus.

Walnut, new mite on, in Central Asia, 530; Cydia pomonella on, in Europe and Palestine, 192, 500, 658; pests of, in U.S.A., 18, 20, 650, 722, 747; injured by insecticides, 747; Lymantria dispar not surviving on, 759; Lyctus in timber of, 190.

walshi, Eurytoma.

Washington, cranberry pests in, 171, 172; miscellaneous pests in, 60, 326, 327, 489; orchard pests in, 174, 488, 544; imported parasites and biological control of Stilpnotia salicis in, 589, 590. Washingtonia robusta, Homaledra sabalella on, in Florida, 22.

Water, plants and timber immersed in, against pests, 273, 764; rôle of, in insect hibernation, 304.

Water, Hot, treatment of plants and planting material with, 104, 273, 314, 383, 384, 437, 488; other uses of, against pests, 31, 539, 608.

Water Gas Tar Oil, 25.

Watercress, *Phaedon cochleariae* on, in Britain, **382**; sawfly on, in China, **497**.

waterhousei, Longitarsus.

Watermelon, new mites on, in Central Asia, 530; Empoasca fabae on, in Bermuda, 626; pests of, S. Carolina, 644.

Wattle Bagworm (see Acanthopsyche

junodi).

Wax Emulsions, increasing efficiency of oil sprays against *Chrysomphalus obscurus*, **18**; and barium fluosilicate, experiments in painting shade trees with, against borers, **543**.

Wax Moth, Large (see Galleria

mellonella).

Wax Moth, Small (see Achroia grisella).

Wesmaëlius concinnus, liberation of, against Chermes pini in Australia, 449.

West Indian Blister Mite (see

Eriophyes gossypii).

Weymouth Pine (see Pinus strobus). Wheat, Aphids on, in Argentina, 30, 31, 289, 290; pests of, in Australia, 169, 584; pests of, in Britain, 102, 382, 463, 527; pests of, in Canada, 59, 121, 170, 441; Mayetiola destructor on, in Cyprus, 108; pests of, in Czechoslovakia, 657, 658, 659; Nezara viridula on, in Egypt, 457; Cephus pygmaeus on, in Estonia, 462; pests of, in Germany, 51, 521, 619; pests of, in India, 241, 628; pests of, in Japan, 54, 670; pests of, in Loochoo Islands, 264, 516; Lema tristis on, in Manchuria, 767; pests of, in Morocco, 500, 501, 641; little attacked by Chlorops taeniopus in Poland, 518; Aelia rostrata on, in Portugal. 697; pests of, in Portugal, 697; pests of, in Russian Union, 76, 160, 422, 424, 425, 426, 427, 428, 617, 618; Hybalus cornifrons on, in Sicily, 757; Pentatomids on, in Spain, 697, 698; gall-midges on, in Sweden, 154, 230, 231, 620; Pentatomids on, in Switzerland, 91, 92; pests of, in U.S.A., 26, 44, 78, 79, 121, 122, 123, 126, 129, 130, 135, 140, 170, 326, 329, 545, 569, 577, 728; damaged by grasshoppers and locusts, 44, 140, 169, 426, 584, 628; varieties and time of sowing of, in relation to pests, 122, 126, 326, 427, 618, 658, 659, 698, 728; type of injury to, by Pentatomids, 51, 92, 521; trap-strips of, against Chlorops taeniopus, 518; planted with beet against wireworms, injured by soil 383, 462; fumigants, 54.

Wheat (Stored), pests of, and their control, 90, 95, 96, 125, 326, 362, 414, 415, 644, 668, 746; effect of fumigants on, 389, 390, 414, 415;

absorption of methyl bromide by, 390.

Wheat Gall-midges (see Contarinia tritici and Sitodiplosis mosellana). Wheat Stem Maggot (see Meromyza

americana).

Wheat Stem Sawfly (see Cephus cinctus).

Wheat Stem Sawfly, Black (see

Trachelus tabidus).

Whiteflies, of Japan, 496; natural enemies of, 41, 296, 337, 449, 491, 582, 623, 727, 748; and plant diseases, 89, 240, 242, 413, 455, 666, 667, 706; classification and new species of, 413, 440, 772.

Whitefly, Citrus (see Dialeurodes

citri).

Whitefly, Greenhouse (see Trialeurodes vaporariorum).

Whitefly, Papaya (see Trialeurodes

variabilis).

Wicker, Leptidea brevipennis damaging, in Czechoslovakia, 658.

williamsi, Frankliniella; Trialeu-

vodes.

Willow (Salix), Stilpnotia salicis on, in Austria, 589; pests of, in Britain, 380, 483, 612, 613; weevil on, in Czechoslovakia, Phassus signifer on, in 658 : Formosa, 517; Ceresa bubalus on, in France, 343; pests of, in Germany, 344, 345, 461, 664, 665, 717; mites on, in Holland, 465; S. salicis on, in Hungary, 589; Pristiphora on, in Japan, 771; Anomala chamaeleon on, in Korea, 767; Melasoma saliceti on, in Russian Union, 429; pests of, in U.S.A., 19, 225, 589, 744; (timber), Coleoptera in, 354, 483, **707.** (See Wicker.)

Willow Aphis, Large (see Lachnus

salignus).

Wind, relation of locust flights to, 139, 146, 678; relations of, to flight and spread of other insects, 157, 215, 216, 284, 307, 308, 314, 378, 406, 423, 455, 474, 547, 606, 607, 661, 709, 752, 753.

Wine Rackings, lees of, in baits for

Wine Rackings, lees of, in baits for Cydia, 168.

Winter Moth (see Operophtera brumata).

Winthemia, parasite of Argyroploce

illepida in Fiji, 497.

Winthemia quadripustulata (datanae), parasite of Lepidoptera in U.S.A., 330, 723; bionomics of, 330. Wireworms (Elaterids), problem of, in Britain, 383, 462, 630; predacious on other insects, 493, 737; soil conditions in relation to, 72, 200, 201, 356, 424, 425, 426, 429, 430; movement of, in soil, 425, 426, 630; technique of sampling soil for, 630; flooding as a means of controlling, 569; cultural measures against, 383, 462; soil fumigation against, 236, 431, 630, 747; survey of measures against, 496; preparation of specimens of, 357.

Wisconsin, Camnula pellucida in, 681; Macrosiphum onobrychis

on peas in, 740.

Wissadula rostrata, Dysdercus on, in Uganda, 374.

woglumi, Aleurocanthus.

woodi, Acarapis.

Woodlice, measures against, on mushrooms in U.S.A., 328.

Wool, insects attacking, 413, 568. (See Textiles.)

Woolly Apple Aphis (see Eriosoma lanigerum).

Woolrex Oil, 82.

Wyoming, Paratrioza cockerelli on potato in, 578.

X.

X-rays, possible value of, for detecting insects in timber, 190; Lyctus not killed by, 190.

Xanthium pungens, insects introduced into Australia against, 450.

xanthius, Taeniothrips.

Xanthodes graellsi (see Acontia). xanthodes, Dacus (Chaetodacus). xanthomelaena, Galerucella (see G. luteola).

xanthophilum, Monalonion.

Xanthopimpla, parasite of Clania minuscula in China, 245.

xanthopleura, Anomala. xanthostylum, Apion.

Xenylla corticalis, measures against, on mushrooms in Holland, 656.

Xerophilaphis scorzonerae, sp. n., bionomics of, on rubberproducing plants in Ukraine, 106, 107.

Xestobium rufovillosum, bionomics of, in timber in Britain, 34, 190, 380, 707; characters of, 34.

Xiphydria prolongata, fungus associated with, in Britain, 632.

Xorides albipictus, parasite of Eutetrapha tridentata in Connecticut, 15.

Xyleborinus saxeseni (see Xyleborus xylographus).

Xyleborus, on pecan in Florida, 587.

Xyleborus arquatus, on Berberis

nepalensis in Ceylon, 359.

Xyleborus coffeae, tentatively recorded on coffee in Réunion, 706.

Xyleborus dispar, on currant in

Finland, 201.

Xyleborus fornicatus (on tea), in Ceylon, 360; measures against, in India, 148; on Crotalaria, 149.

Xyleborus germanus, Ceratostomella ulmi cultured from, on elm in New Jersey, 325.

Xyleborus mancus, on Styrax in

Malaya, 453.

Xyleborus movstatti, measures against, on avocado in Fiji, 497. Xyleborus xylographus, on Acer

pseudoplatanus in France, 316. Xyleutes ceramicus, bionomics of, on

teak in Burma, 338, 339. Xyleutes persona, larval stage of, on

Cassia spp. in Burma, 340. Xylia dolabriformis, pests of, in

Burma, **339.**

Xylobiops basilare, Ceratostomella ulmi cultured from, on elm in New Jersey, 325.

xylographus, Xyleborus.

Xylomyges eridania, susceptibility of, to nicotine, 332.

Xylonomus sapporensis, parasite of Callidium rufipenne in Japan,

Xylosandrus (see Xyleborus).

xylosteana, Tortrix (Cacoecia).

Xyloteles (see Xylotoles).

Xyloterus lineatus, on Pinus sylvestris in Estonia, 462.

Xylotoles spp., bionomics and control of, in New Zealand, 266,

Xylotrechus antilope, bionomics of, on cork oak in Morocco, 46.

Xylotrechus quadripes, measures against, on coffee in Mysore, 55.

Xystrocera festiva, on coffee Indo-China, 261; bionomics and control of, in Java, 261.

Xystus (see Charips).

Y.

Yam, Aspidiotus harti on, in Cayman Islands, 269; Pseudococcus kenyae on, in Kenya, 592.

vamamai, Antheraea.

Yeast, in food for Rhagoletis pomonella, 561.

Yellow Flat Disease of Lilies (see Rosette Disease).

Yellow-headed Fireworm (see Peronea minuta).

yponomeutae, Discochaeta.

ypsilon, Agrotis (Rhyacia).

Yucca, Lepidosaphes beckii on, in Palestine, 636. yuccae, Puto.

Ζ.

Zabrus tenebrioides, on cereals in Czechoslovakia, 657.

Zanzibar, parasites of Trypetids in, 492; natural enemies of Coccids in, 611; plant pest legislation in, 111.

zanzibarensis, Synacanthotermes.

Zatropis incertus, associated with cotton pests in Mexico, 275.

zehntneri, Elasmus.

Zelkova, Agrilus spinipennis on, in Japan, 265.

Zelus socius, predacious on Eutettix tenellus in Utah, 128.

Zenillia caesar, parasite of Pyrausta nubilalis in U.S.A., 730.

Zenillia ceratomiae, parasite of Desmia funeralis in California, 286. Zenillia distincta, parasite of Protoparce in Jamaica, 490.

Zenillia mitis, parasite of Pyrausta nubilalis in Poland, 623.

Zenillia nemea, parasite of Pieris vapae in Britain, 342.

Zenillia pullata, bionomics of, in Russian Union, 303.

Zenillia susurrans, parasite of Dioryctria abietella in Czechoslovakia, 529.

Zenillia vulgaris, parasite of Pieris spp. in Britain, 342.

Zephyranthes candida, Brithys crini on, in Formosa, 52.

Zeuxia cinerea, parasite of Cleonus mendicus in Italy, 527.

Zeuxia nigripes, parasite of Lixus junci in Morocco, 417.

Zeuzera coffeae, on castor in Ceylon, 359; on tea in India, 149.

Zeuzera pyrina, measures against, in Cyprus, 260; in Czechoslovakia, 658; in Malta, 469; factors affecting infestation by, in Russia, 75; in forests, 75, 658; on fruit trees, 75, 260, 469.

Arsenate, spraying with, against orchard pests, 16, 79, 174, 401, 489, 556, 724; adhesive and spreader for, 16, 489; formulae containing, 16, 401, 489, 724; and lime, 401, 489, 724; and lime-sulphur, 489; inverted spray mixtures of oils and, 174; and sulphur, 724; and zinc sulphate, 724; other insecticides compared with, 79, 401, 556.

Zinc Arsenite, spraying with, against bean and potato beetle, 327, 643; and lime-sulphur, formula containing, 643; cryolite, 327. compared with

Zinc Carbonate, as an insecticide, 248.

Zinc Chloride, for treating timber against Lyctus, 190, 514.

Zinc Oxide, as an insecticide, 248. Zinc Phosphide, in baits for crickets, 457, 528.

Zinc Sulphate, action of, in arsenical sprays, 16, 125, 138, 174, 175, 587; use of, for precipitating sulphur from lime-sulphur, 488; formulae containing, 125, 138, 174, 443, 588, 724.

zinckenella, Etiella.

Zizera (Zizina) otis lysizone, on Indigofera endecaphylla in Java, 110.

Zizyphus, not attacked by Parlatoria zizyphus in Formosa, 263. Zizyphus jujuba, Carpomyia vesuviana on, in India, 765.

Zizyphus spina-christi, new Aleurodid on, in Egypt, 440.

zizyphus, Parlatoria.

Zodion fulvifrons, bionomics of, in U.S.A., 133.

zonatus, Dacus (Chaetodacus).

zonellus, Chilo.

Zonocerus elegans, damaging lucerne in S. Africa, 152; damaging cassava, etc., in Tanganyika, 112, 642; Empusa infesting, 642; measures against, 152, 642.

Zonosemata electa, on Capsicum in

Connecticut, 17.

zoolathridi, Cinara utahensis.

Zootermopsis, in U.S.A., 219.

Zootermopsis angusticollis, imported into Japan in timber Oregon, 55.

zuimushi, Amphimermis.

Zyklon, fumigation with, 38, 591,

NOTICES.

Secretaries of Societies and Editors of Journals willing to exchange their publications with those of the Institute are requested to communicate with the Assistant Director.

The Annual Subscription, in advance, to the Review, Series A (Agricultural), is 30s. post free; Series B (Medical and Veterinary), 15s. post free. Subscriptions to the current Volume received after 30th June in each year will be charged at the published price, viz.: Series A, 36s.; Series B, 18s. Prices of Back Volumes on application.

Orders and Subscriptions should be sent direct to the Assistant Director, Imperial Institute of Entomology, 41, Queen's Gate, London, S.W.7, or through a bookseller.